

## EXTENSION OF CLASSICAL SEQUENCES TO NEGATIVE INTEGERS

BENALI BENZAGHOU

*Faculté de Mathématiques*  
*USTHB BP 32 El-Alia,*  
*16111 Bab-Ezzouar, Alger*

**e-mail:** benrect@wissal.dz

AND

DANIEL BARSKY

*Université Paris 13, Institut Galilée,*  
*LAGA UMR 7539, 99 Av. J.-B.*  
*Clément 93430 Villetaneuse, France*

**e-mail:** barsky@math.univ-paris13.fr

### Abstract

We give a method to extend Bell exponential polynomials to negative indices. This generalizes many results of this type such as the extension to negative indices of Stirling numbers or of Bernoulli numbers.

**Keywords:** Stirling numbers, Bell polynomials.

**2000 Mathematics Subject Classification:** 11B73.

### REFERENCES

- [1] D. Branson, *An extension of Stirling numbers*, *Fib. Quat.* **34** (3) (1996), 213–223.
- [2] L. Comtet, *Analyse combinatoire*, Vol. I and II, Presses Universitaires de France, Paris 1970.

- [3] S. Roman, *The harmonic logarithms and the binomial formula*, J. Combin. Theory, Serie A, **63** (1993), 143–163.

Received 2004  
Revised December 2005