

## MAXIMAL SUBMONOIDS OF MONOIDS OF HYPERSUBSTITUTIONS

ILINKA DIMITROVA AND JÖRG KOPPITZ

*University of Potsdam*

*Institute of Mathematics*

*Am Neuen Palais, 14415 Potsdam, Germany*

**e-mail:** ilinkad@abv.bg; koppitz@rz.uni-potsdam.de

### Abstract

For a monoid  $M$  of hypersubstitutions, the collection of all  $M$ -solid varieties forms a complete sublattice of the lattice  $\mathcal{L}(\tau)$  of all varieties of a given type  $\tau$ . Therefore, by the study of monoids of hypersubstitutions one can get more insight into the structure of the lattice  $\mathcal{L}(\tau)$ . In particular, monoids of hypersubstitutions were studied in [9] as well as in [5]. We will give a complete characterization of all maximal submonoids of the monoid  $Reg(n)$  of all regular hypersubstitutions of type  $\tau = (n)$  (introduced in [4]). The concept of a transformation hypersubstitution, introduced in [1], gives a relationship between monoids of hypersubstitutions and transformation semigroups. In the present paper, we apply the recent results about transformation semigroups by I. Guydzenov and I. Dimitrova ([11], [12]) to describe monoids of transformation hypersubstitutions.

**Keywords:** regular hypersubstitutions, maximal monoids of hypersubstitutions, transformation semigroups.

**2000 Mathematics Subject Classification:** 08B05, 08B15, 20M07, 16Y60.

### REFERENCES

- [1] V. Budd, K. Denecke and S.L. Wismath, *Short Solid Superassociative Type (n) Varieties*, East-West Journal of Mathematics **2** (2) (2001), 129–145.

- [2] Th. Changphas, Monoids of Hypersubstitutions, Dissertation, Universität Potsdam 2004.
- [3] Th. Changphas and K. Denecke, *Full Hypersubstitutions and Full Solid Varieties of Semigroups*, East-West Journal of Mathematics **4** (1) (2002), 177–193.
- [4] K. Denecke and J. Koppitz, *M-Solid Varieties of Semigroups*, Discuss. Math. **15** (1995), 23–41.
- [5] K. Denecke and J. Koppitz, *M-Solid Varieties of Algebras*, Springer Science-Business Media 2006.
- [6] K. Denecke, J. Koppitz and S. Niwczyk, *Equational theories generated by hypersubstitutions of type (n)*, Int. Journal of Algebra and Computation **12** (6) (2002), 867–876.
- [7] K. Denecke, J. Koppitz and S. Shtrakov, *The Depth of a Hypersubstitution*, Journal of Automata, Languages and Combinatorics **6** (3) (2001), 253–262.
- [8] K. Denecke and M. Reichel, *Monoids of hypersubstitutions and M-solid varieties*, Contributions to General Algebra 9, Wien 1995, 117–126.
- [9] K. Denecke and S.L. Wismath, *Hyperidentities and clones*, Gordon and Breach Scientific Publishers, 2000.
- [10] K. Denecke and S.L. Wismath, *Complexity of Terms, Composition, and Hypersubstitution*, Int. Journal of Mathematics and Mathematical Sciences **15** (2003), 959–969.
- [11] V.H. Fernandes, G.M.S. Gomes and M.M. Jesus, *Presentations for Some Monoids of Partial Transformations on a Finite Chain*, Communications in Algebra **33** (2005), 587–604.
- [12] Il. Gyudzhenov and Il. Dimitrova, *On the Maximal Subsemigroups of the Semigroup of All Isotone Transformations with Defect  $\geq 2$* , Comptes rendus de l'Academie bulgare des Sciences **59** (3) (2006), 239–244.
- [13] Il. Gyudzhenov and Il. Dimitrova, *On the Maximal Subsemigroups of the Semigroup of all Monotone Transformations*, Discuss. Math., submitted.
- [14] J.M. Howie, *An Introduction to Semigroup Theory*, Academic Press, London 1976.
- [15] M.W. Liebeck, C.E. Praeger and J. Saxl, *A Classification of the Maximal Subgroups of the Finite Alternating and Symmetric Groups*, Journal of Algebra **111** (1987), 365–383.
- [16] X. Yang, *A Classification of Maximal Subsemigroups of Finite Order-Preserving Transformation Semigroups*, Communications in Algebra **28** (3) (2000), 1503–1513.

Received 23 March 2006  
Revised 13 April 2006