

k -NORMALIZATION AND $(k + 1)$ -LEVEL INFLATION OF VARIETIES

VALERIE CHENG AND SHELLY WISMATH

Math/C.S. Dept., University of Lethbridge
401 University Drive, Lethbridge, Ab., T1K-3M4 Canada

e-mail: wismaths@cs.uleth.ca

Abstract

Let τ be a type of algebras. A common measurement of the complexity of terms of type τ is the depth of a term. For $k \geq 1$, an identity $s \approx t$ of type τ is said to be k -normal (with respect to this depth complexity measurement) if either $s = t$ or both s and t have depth $\geq k$. A variety is called k -normal if all its identities are k -normal. Taking $k = 1$ with respect to the usual depth valuation of terms gives the well-known property of normality of identities or varieties. For any variety V , there is a least k -normal variety $N_k(V)$ containing V , the variety determined by the set of all k -normal identities of V . The concept of k -normalization was introduced by K. Denecke and S.L. Wismath in [5], and an algebraic characterization of the elements of $N_k(V)$ in terms of the algebras in V was given in [4]. In [1] a simplified version of this characterization of $N_k(V)$ was given, in the special case of the 2-normalization of the variety V of all lattices, using a construction called the 3-level inflation of a lattice. In this paper we show that the analogous $(k + 1)$ -level inflation can be used to characterize the algebras of $N_k(V)$ for any variety V having a unary term which satisfies two technical conditions. This includes any variety V which satisfies $x \approx t(x)$ for some unary term t of depth at least k , and in particular any variety, such as the variety of lattices, which satisfies an idempotent identity.

Keywords: k -normal identities, k -normalization of a variety, $(k + 1)$ -level inflation of algebras.

2000 Mathematics Subject Classification: 08A40, 08B15, 08B05.

REFERENCES

- [1] I. Chajda, V. Cheng and S.L. Wismath, *2-Normalization of lattices*, to appear in Czechoslovak Mathematics Journal.
- [2] A. Christie, Q. Wang and S.L. Wismath, *Minimal characteristic algebras for k -normality*, *Scientiae Mathematicae Japonicae* **61** (3) (2005), 547–565.
- [3] G.T. Clarke, *Semigroup varieties of inflations of unions of groups*, *Semigroup Forum* **23** (4) (1981), 311–319.
- [4] K. Denecke and S.L. Wismath, *A characterization of k -normal varieties*, *Algebra Universalis* **51** (4) (2004), 395–409.
- [5] K. Denecke and S.L. Wismath, *Valuations of terms*, *Algebra Universalis* **50** (1) (2003), 107–128.
- [6] E. Graczyńska, *On normal and regular identities*, *Algebra Universalis* **27** (3) (1990), 387–397.
- [7] I.I. Mel'nik, *Nilpotent shifts of varieties*, (in Russian), *Mat. Zametki*, **14** (1973), 703–712; English translation in: *Math. Notes* **14** (1973), 962–966.

Received 11 July 2006

Revised 30 May 2007