

ON THE AUTOTOPISM GROUP OF THE CORDERO-FIGUEROA SEMIFIELD OF ORDER 3^6

WALTER MELÉNDEZ

RAÚL FIGUEROA

University of Puerto Rico, Río Piedras Campus
Mathematics Department, P.O. Box 70377
San Juan, PR 00936 – 8377, USA

e-mail: wflorian75@gmail.com
junioyjulio@gmail.com

AND

MOISÉS DELGADO

University of Puerto Rico, Cayey Campus
Mathematics and Physics Department
205 Calle Antonio R. Barceló
Cayey, PR 00736, USA

e-mail: moises.delgado@upr.edu

Abstract

In [5] M. Biliotti, V. Jha and N. Johnson were able to completely determine the autotopism group of a generalized twisted field as a subgroup of $\Gamma L(K) \times \Gamma L(K)$, where $K = GF(p^n)$ and $\Gamma L(K)$ is the group of non-singular semilinear transformations over K . In this article, we consider the Cordero-Figueroa semifield of order 3^6 , which is not a generalized twisted field, and we prove that its autotopism group is isomorphic to a subgroup of $\Gamma L(K) \times \Gamma L(K)$, where $K = GF(3^6)$.

Keywords: finite presemifield, finite semifield, autotopism, autotopism group.

2010 Mathematics Subject Classification: 12K10, 17A35.

REFERENCES

- [1] A.A. Albert, *Generalized twisted fields*, Pacific J. Math **11** (1961) 1–8.
doi:10.2140/pjm.1961.11.1

- [2] A.A. Albert, *On nonassociative division algebras*, TAMS **72** (1952) 296–309. doi:10.1090/S0002-9947-1952-0047027-4
- [3] A.A. Albert, *Finite division algebras and finite planes*, Proc. Sympos. Appl. Math. **10** (1960) 53–70. doi:10.1090/psapm/010/0116036
- [4] J. Bierbrauer, *Projective polynomials, a projection construction and a family of semifields*, Designs, Codes and Cryptography **79** (2016) 183–200. doi:10.1007/s10623-015-0044-z
- [5] M. Biliotti, V. Jha and N. Johnson, *The collineation groups of generalized twisted field planes*, G. Dedicata **76** (1999) 91–126.
- [6] N. Jhonson, V. Jha and M. Biliotti, *Handbook of finite translation planes* (Boca Raton: Chapman & Hall/CRC, 2007).
- [7] M. Cordero and R. Figueroa, *Towards a characterization of the generalized twisted field planes*, J. Geom. **52** (1995) 54–63. doi:10.1007/BF01406826
- [8] R.S. Coulter, M. Henderson and P. Kosick, *Planar polynomials for commutative semifields with specified nuclei*, Des. Codes Cryptogr. **44** (2007) 275–286.
- [9] L.E. Dickson, *Linear algebras in which division is always uniquely possible*, Trans. Amer. Math. Soc. **7** (1906) 370–390. doi:10.1090/S0002-9947-1906-1500755-5
- [10] R. Figueroa, *A characterization of the generalized twisted field planes of characteristic ≥ 5* , Geom. Dedicata **50** (1994) 205–216. doi:10.1007/BF01265311
- [11] D.R. Hughes and F.C. Piper, *Projective Planes* (NewYork, 1973).
- [12] D. E. Knuth, *Finite semifields and projective planes*, J. Algebra **2** (1965) 182–217. doi:10.1016/0021-8693(65)90018-9
- [13] M. Lavrauw, *On the isotopism classes of finite semifields*, Finite Fields and Their Applications **14** (2008) 897–910. doi:10.1016/j.ffa.2008.05.002

Received 23 March 2016

Revised 17 April 2016