

A RESULT ON PRIME RINGS WITH GENERALIZED DERIVATIONS

FAIZA SHUJAT

*Department of Mathematics, College of Science
Taibah University, Madinah, Saudi Arabia*

e-mail: fullahkhan@taibahu.edu.sa

AND

SHAHOOR KHAN¹

*Government Degree College, Department of Mathematics
Surankote, 185121, Jammu and Kashmir, India*

e-mail: shahoor.khan@rediffmail.com

Abstract

In this paper we investigate the following result. Let R be a prime ring, Q its symmetric Martindale quotient ring, C its extended centroid, I a nonzero ideal of R . If F and G are the two generalized derivation of R such that $(\mathbf{F}(\mathbf{xy}) + \mathbf{G}(\mathbf{yx}))^n - (\mathbf{xy} \mp \mathbf{yx})^n = 0$, for all $x, y \in I$, then either R is commutative or $F(x) = x$, $G(x) = \mp x$ for all $x \in R$ and $n = 1$.

Keywords: prime ring, generalized derivations, quotient ring, extended centroid.

2010 Mathematics Subject Classification: 16W20, 16W25, 16U80, 16N80.

REFERENCES

- [1] C. Faith and Y. Utumi, *On a new proof of Litoff's theorem*, Acta Math. Acad. Sci. Hungar. **14** (1963) 369–371.
<https://doi.org/10.1007/BF01895723>
- [2] C.L. Chuang, *GPI's having coefficients in Utumi quotient rings*, Proc. Amer. Math. Soc. **103** (1988) 723–728.
<https://doi.org/10.1090/S0002-9939-1988-0947646-4>

¹Corresponding author.

- [3] C. Lanski, *An engle condition with derivation*, Proc. Amer. Mathp. Soc. **183** (1993) 731–734.
<https://doi.org/10.1090/S0002-9939-1993-1132851-9>
- [4] I.N. Herstein, *Topics in Ring Theory* (Univ. of Chicago Press, Chicago, 1969).
- [5] K.I. Beidar, W.S. Martindale III and A.V. Mikhalev, *Rings with generalized identities*, Monographs and Textbooks in Pure and Applied Math. **196** (New York, Marcel Dekker, Inc. 1996).
- [6] M.N. Daif and H.E. Bell, *Remarks on derivations on semiprime rings*, Int. J. Math. Math. Sci. **15** (1992) 205–206.
<https://doi.org/10.1155/S0161171292000255>
- [7] N. Jacobson, *Structure of Rings*, Amer. Math. Soc. Colloq. Pub. **37** (Amer. Math. Soc., Providence, RI, 1964).
- [8] S. Khan, F. Shujat and G. Alhendi, *A Result on annihilator condition and generalized derivations of prime rings*, J.P. Journal of Alg. Num. Th. Appl. **43** (2019) 101–110.
<https://doi.org/10.17654/NT043020101>
- [9] S. Huang and B. Davvaz, *Generalized derivations of rings and Banach algebras*, Comm. Algebra **41** (2013) 1188–1194.
<https://doi.org/10.1080/00927872.2011.642043>
- [10] T.S. Erickson, W.S. Martindale III and J.M. Osborn, *Prime non-associative algebras*, Pacific J. Math. **60** (1975) 49–63.
<https://doi.org/10.2140/pjm.1975.60.49>
- [11] T.K. Lee, *Semiprime rings with differential identities*, Bull. Inst. Math. Acad. Sinica **20** (1992) 27–38.
- [12] T.K. Lee, *Generalized derivations of left faithful rings*, Comm. Algebra **27** (8) (1999) 4057–4073.
<https://doi.org/10.1080/00927879908826682>
- [13] V.K. Kharchenko, *Differential identity of prime rings*, Algebra and Logic **17** (1978) 155–168.
<https://doi.org/10.1007/BF01670115>
- [14] V. De Filippis and S. Huang, *Generalized derivations on semi prime rings*, Bull. Korean Math. Soc. **48** (6) (2011) 1253–1259.
<https://doi.org/10.4134/BKMS.2011.48.6.1253>
- [15] W.S. Martindale III, *Prime rings satisfying a generalized polynomial identity*, J. Algebra **12** (1972) 576–584.
[https://doi.org/10.1016/0021-8693\(69\)90029-5](https://doi.org/10.1016/0021-8693(69)90029-5)

Received 1 September 2020

Revised 14 November 2020

Accepted 14 November 2020