

CANTOR EXTENSION OF A HALF LINEARLY CYCLICALLY ORDERED GROUP

ŠTEFAN ČERNÁK

*Department of Mathematics, Faculty of Civil Engineering, Technical University,
Vysokoškolská 4, SK-042 02 Košice, Slovakia*

e-mail: svfkm@tuke.sk

Abstract

Convergent and fundamental sequences are studied in a half linearly cyclically ordered group G with the abelian increasing part. The main result is the construction of the Cantor extension of G .

Keywords: convergent sequence, fundamental sequence, C -complete half lc -group, Cantor extension of a half lc -group.

2000 Mathematics Subject Classification: Primary 06F15; Secondary 20F60.

M. Giraudet and F. Lucas [3] introduced and investigated the notion of a half linearly ordered group (cf. also D.R. Ton [14], J. Jakubík [6], [7]). J. Jakubík [8] defined and studied the notion of a half linearly cyclically ordered group (lc -group) generalizing the notion of a half linearly ordered group.

The author [1] investigated the Cantor extension of an abelian lc -group. We remark that the Cantor extension of lattice ordered groups was studied by C.J. Everett [2].

Let G be a half lc -group such that its increasing part is abelian and its decreasing part is nonempty (thus G fails to be an lc -group). The notions of a convergent sequence and a fundamental sequence are defined in a natural way. If every fundamental sequence in G is convergent in G , then G is said to be C -complete.

In the present paper necessary and sufficient conditions are found under which G is C -complete. Further, we define the notion of a Cantor extension and we prove that every half lc -group has a Cantor extension which is uniquely determined up to isomorphisms leaving all elements of G fixed.

1. l -CYCLICALLY ORDERED SETS AND GROUPS

We recall the definitions and some results concerning l -cyclically ordered sets (cf. Novák and Novotný [10], Novák [9], Quilot [11]) and l -cyclically ordered groups (cf. Rieger [12], Świerczkowski [13], Jakubík and Pringerová [4], [5]).

Definition 1.1. Let M be a nonempty set and T a ternary relation on M such that the following conditions are satisfied:

- (I) if $[x, y, z] \in T$ then $[y, x, z] \notin T$.
- (II) $[x, y, z] \in T$ implies $[y, z, x] \in T$.
- (III) $[x, y, z] \in T, [y, u, z] \in T$ imply $[x, u, z] \in T$.

Then T is said to be a *cyclic order* on M and (M, T) is called a *cyclically ordered set*.

Let T be a cyclic order on M satisfying the condition:

- (IV) if $x, y, z \in M, x \neq y \neq z \neq x$, then either $[x, y, z] \in T$ or $[z, y, x] \in T$.

Then T is said to be an *l -cyclic order* on M and (M, T) is called an *l -cyclically ordered set*.

Several terms are used in papers for the term l -cyclic order. For instance " l -cyclic order" is called "linear cyclic order" in [9], "complete cyclic order" in [11] and simply "cyclic order" in [12] and [13].

Definition 1.2. Let $(H; +)$ be a group and $(H; T)$ an l -cyclically ordered set such that the following condition is fulfilled:

- (V) if $[x, y, z] \in T, u, v \in H$, then $[u + x + v, u + y + v, u + z + v] \in T$.

Then $(H; +, T)$ is said to be an *l -cyclically ordered group* or *lc-group* (*linearly cyclically ordered group*).

We often write H or $(H; T)$ instead of $(H; +, T)$.

Every subgroup of an lc -group is considered as an lc -group under the induced l -cyclic order.

Example 1.3. Let $(L; \leq)$ be a linearly ordered group $x, y, z \in L$. Define the ternary relation T_L on L by putting

$$[x, y, z] \in T_L \text{ if } x < y < z \text{ or } y < z < x \text{ or } z < x < y.$$

Then $(L; T_L)$ is an lc -group. T_L is called the l -cyclic order generated by the linear order \leq on L . Hence every linearly ordered group is an lc -group (under the l -cyclic order generated by its linear order).

Example 1.4. Let K be the group of all reals k such that $0 \leq k < 1$ with the group operation defined as the addition mod 1. Consider the natural linear order \leq and the ternary relation T_1 on K defined in the same way as T_L in 1.3. Then $(K; T_1)$ is an lc -group.

Define the ternary relation T on the direct product $L \times K$ of groups L and K as follows: for elements $u_1, u_2, u_3 \in L \times K$, $u_1 = (x, k_1)$, $u_2 = (y, k_2)$, $u_3 = (z, k_3)$ we put $[u_1, u_2, u_3] \in T$ if some of the following conditions is valid:

- (i) $[k_1, k_2, k_3] \in T_1$;
- (ii) $k_1 = k_2 \neq k_3$ and $x < y$;
- (iii) $k_2 = k_3 \neq k_1$ and $y < z$;
- (iv) $k_3 = k_1 \neq k_2$ and $z < x$;
- (v) $k_1 = k_2 = k_3$ and $[x, y, z] \in T_L$.

Then $(L \times K; T)$ is an lc -group which will be denoted by $L \otimes K$.

The notion of an isomorphism of lc -groups is defined in a natural way.

Theorem 1.5 (Świerczkowski [13]). *Let H be an lc -group. Then there exists a linearly ordered group L such that H is isomorphic to a subgroup of $L \otimes K$.*

Assume that $(H; T)$ is an lc -group. By 1.5, there exists an isomorphism f of H into $L \otimes K$. Let H_o be the set of all $h \in H$ such that there exists $x \in L$ with the property $f(h) = (x, 0)$. Then H_o is a subgroup of H , $H_o = \{0\}$ or $H_o \neq \{0\}$. Let $H_o \neq \{0\}$, $h \in H_o$, $h \neq 0$. There exists $x \in L$ such that $f(h) = (x, 0)$. H_o turns out to be a linearly ordered group if we put $h > 0$ if $x > 0$. The l -cyclic order T_{H_o} on H_o coincides with the l -cyclic order induced by T .

2. CANTOR EXTENSION OF AN ABELIAN lc -GROUP

Let $(H; T)$ be an abelian lc -group. A construction of a Cantor extension of H will be described (cf. [1]) and some results from [1] will be presented.

Definition 2.1. Let (x_n) be a sequence in H and $x \in H$.

- a) We say that (x_n) converges to x (or x is a limit of (x_n)) in H and we write $x_n \rightarrow x$ (or $\lim x_n = x$)
 - (i) if $\text{card } H = 2$ and there exists $n_o \in N$ such that $x_n = x$ for each $n \in N, n \geq n_o$,
 - or
 - (ii) if $\text{card } H > 2$ and for each $\varepsilon \in H, \varepsilon \neq 0$ with the property $[-\varepsilon, 0, \varepsilon] \in T$ there exists $n_o \in N$ such that $[-\varepsilon, x_n - x, \varepsilon] \in T$ for each $n \in N, n \geq n_o$.
- b) The sequence (x_n) is called *fundamental* in H if for each $\varepsilon \in H, \varepsilon \neq 0$ with the property $[-\varepsilon, 0, \varepsilon] \in T$ there exists $n_o \in N$ such that $[-\varepsilon, x_n - x_m, \varepsilon] \in T$ for each $m, n \in N, m, n \geq n_o$.
- c) By a *zero sequence* we understand a sequence (x_n) such that $x_n \rightarrow 0$.
- d) H is called *C-complete* if each fundamental sequence in H is convergent in H .

The set of all fundamental (zero) sequences in H will be denoted by $F_H(E_H)$.

Definition 2.2. Let H_1 be an abelian *lc*-group satisfying the following conditions:

- (a) H_1 is *C*-complete.
- (b) H is a subgroup of H_1 .
- (c) Every element of H_1 is a limit of some fundamental sequence in H .
- (d) Let (x_n) be a sequence in H such that $x_n \rightarrow 0$ in H . Then $x_n \rightarrow 0$ in H_1 .

Then H_1 is said to be a *Cantor extension* of H .

Now we consider two cases: $H_0 \neq \{0\}$ and $H_0 = \{0\}$.

1) Assume that $H_0 \neq \{0\}$. Let $(x_n), (y_n) \in F_H$. Under the natural definition of the operation $+$ on $F_H, (x_n) + (y_n) = (x_n + y_n), F_H$ is a group and E_H is a subgroup of F_H . We form the factor group $H^* = F_H/E_H$. Symbol $(x_n)^*$ will denote the coset of H^* containing the sequence $(x_n) \in F_H$.

Suppose that $(x_n)^*, (y_n)^*, (z_n)^*$ are mutually distinct elements of H^* . Let T^* be the set of all triples $[(x_n)^*, (y_n)^*, (z_n)^*]$ of elements of H^* such that there exists $n_o \in N$ with $[x_n, y_n, z_n] \in T$ for each $n \in N, n \geq n_o$. Then (H^*, T^*) is an *lc*-group.

Let φ be the mapping from H into H^* defined by $\varphi(x) = (x, x, \dots)^*$ for each $x \in H$. Then φ is an isomorphism of the lc -group H into H^* . We identify x and $\varphi(x)$ for each $x \in H$. Then H is a subgroup of H^* and H^* is a Cantor extension of H .

If we denote $(x_n)^* = X$ and $(x_n, x_n, \dots)^* = X_n$, then we have (cf. [1], the proof of Lemma 3.12)

$$(A) \quad X_n \rightarrow X \text{ in } H^*.$$

Lemma 2.3 ([1], Lemma 3.9). *H is C -complete if and only if H_o is C -complete.*

2) Now assume that $H_o = \{0\}$. Then H can be considered as a subgroup of K .

Lemma 2.4 ([1], Lemma 4.2). *If H is a finite subgroup of K , then H is C -complete.*

Lemma 2.5 ([1], Lemma 4.5). *If H is an infinite subgroup of K , then K is a Cantor extension of H .*

The following result is valid in both cases 1) and 2).

Theorem 2.6 ([1], Theorem 4.9). *Let H be an abelian lc -group. Then*

- (i) *there exists a Cantor extension of H ,*
- (ii) *if H_1 and H_2 are Cantor extensions of H , then there exists an isomorphism Φ from the lc -group H_1 onto H_2 such that $\Phi(x) = x$ for each $x \in H$.*

3. HALF lc -GROUPS

The notion of a half lc -group was introduced by Jakubík [8]. Now we recall the definitions and results that will be applied in the next sections.

Let $(G; +, T)$ be a system such that $(G; +)$ is a group and $(G; T)$ is a cyclically ordered set. Assume that $x, y, z \in G$. Denote

$$G \uparrow = \{u \in G : [x, y, z] \in T \Rightarrow [u + x, u + y, u + z] \in T\},$$

$$G \downarrow = \{u \in G : [x, y, z] \in T \Rightarrow [u + z, u + y, u + x] \in T\}.$$

Definition 3.1. Let $(G; +, T)$ be as above. Assume that the following conditions are fulfilled:

- (1) The system T is nonempty.
- (2) If $[x, y, z] \in T$, then $[x + u, y + u, z + u] \in T$ for each $u \in G$.
- (3) $G = G \uparrow \cup G \downarrow$.
- (4) If $[x, y, z] \in T$, then either $\{x, y, z\} \subseteq G \uparrow$ or $\{x, y, z\} \subseteq G \downarrow$.

Then $(G; +, T)$ is said to be a *half cyclically ordered group*.

Let $(G; +, T)$ be a half cyclically ordered group. The definition implies that $G \uparrow$ is a cyclically ordered group. If $G \uparrow$ is an *lc*-group then $(G; +, T)$ is called a *half lc-group* (half linearly cyclically ordered group).

There are elements $x, y, z \in G$ with $[x, y, z] \in T$. This is an immediate consequence of (1).

Again, we often write G or $(G; T)$ instead of $(G; +, T)$.

In the next, let G be a half *lc*-group. $G \uparrow$ ($G \downarrow$) is called the *increasing* (*decreasing, resp.*) part of G .

A subgroup G' of G is said to be a *half lc-subgroup* of G if the *induced* *l*-cyclic order on G' is nonempty.

Each *lc*-group G with $\text{card } G \geq 3$ is a half *lc*-group (with $G \uparrow = G$ and $G \downarrow = \emptyset$). Every linearly ordered group is an *lc*-group. Hence every half linearly ordered group (for the definition cf. [3]) is a half *lc*-group.

The notion of an isomorphism of half *lc*-groups is defined in a natural way.

From the definition 3.1 it follows (cf. [8]):

- (i) If $x, y \in G \downarrow$, then $x + y \in G \uparrow$;
- (ii) If $x \in G \uparrow, y \in G \downarrow$, then $x + y \in G \downarrow$ and $y + x \in G \downarrow$.

4. CANTOR EXTENSION OF A HALF *lc*-GROUP

In what follows, we assume that (G, T) is a half *lc*-group such that $G \uparrow$ is abelian and $G \downarrow \neq \emptyset$. Hence G is neither abelian group nor *lc*-group.

We will use the notation $G \uparrow = H$ and $G \downarrow = H'$.

Definition 4.1. Let (x_n) be a sequence in G and $x \in G$.

- a) We say that (x_n) *converges* to x (or x is a *limit* of (x_n)) in G and we write $x_n \rightarrow x$ (or $\lim x_n = x$) if for each $\varepsilon \in G, \varepsilon \neq 0$ with the property $[-\varepsilon, 0, \varepsilon] \in T$ there exists $n_o \in N$ such that $[-\varepsilon, x_n - x, \varepsilon] \in T$ and $[-\varepsilon, -x + x_n, \varepsilon] \in T$ for each $n \in N, n \geq n_o$.
- b) The sequence (x_n) is said to be *fundamental* if for each $\varepsilon \in G, \varepsilon \neq 0$ with $[-\varepsilon, 0, \varepsilon] \in T$ there exists $n_o \in N$ such that $[-\varepsilon, x_n - x_m, \varepsilon] \in T$ and $[-\varepsilon, -x_m + x_n, \varepsilon] \in T$ for each $m, n \in N, m, n \geq n_o$.
- c) If $x_n \rightarrow 0$ in G , then (x_n) is called a *zero sequence* in G .
- d) G is said to be *C-complete* if every fundamental sequence in G is convergent in G .

Definition 4.2. Let G_1 be a half *lc*-group with the following properties:

- (α) G_1 is *C-complete*;
- (β) G is a half *lc*-subgroup of G_1 ;
- (γ) Every element of G_1 is a limit of some fundamental sequence in G ;
- (δ) Let (x_n) be a sequence in G such that $x_n \rightarrow 0$ in G . Then $x_n \rightarrow 0$ in G_1 .

Then G_1 is said to be a *Cantor extension* of G .

We prove that G has a Cantor extension and this is uniquely determined up to isomorphisms leaving all elements of G fixed.

Denote by $F(E)$ the set of all fundamental (zero) sequences in G . Symbols F_H and E_H have the same meaning as in the section 2.

The following two lemmas are easy to prove.

Lemma 4.3. *Let (x_n) be a sequence in G . Then $x_n \rightarrow x$ in G if and only if $x_n - x \rightarrow 0$ and $-x + x_n \rightarrow 0$ in G . ■*

For a fixed element $n_o \in N$ and a sequence (x_n) in G we apply the notation $x_n^o = x_{n_o+n-1}$ for each $n \in N$.

Lemma 4.4. *Let (x_n) be a sequence in G .*

- (i) $(x_n) \in E$ if and only if there exists $n_o \in N$ such that (x_n^o) is a sequence in H and $(x_n^o) \in E_H$.
- (ii) Let $x \in G$ such that $x_n \rightarrow x$ in G . Then there exists $n_o \in N$ such that either (x_n^o) is a sequence in H (and then $x \in H$) or (x_n^o) is a sequence in H' (and then $x \in H'$).
- (iii) Let $(x_n) \in F$. Then there exists $n_o \in N$ such that either (x_n^o) is a sequence in H (and then $(x_n^o) \in F_H$) or (x_n^o) is a sequence in H' .

Let (x_n) be a sequence in $H, x \in H$. Then

- (iv) $x_n \rightarrow x$ in H if and only if $x_n \rightarrow x$ in G . ■

Let $\varepsilon \in G, \varepsilon \neq 0$. If $[-\varepsilon, 0, \varepsilon] \in T$, then $\varepsilon \in H$. Thus we have:

Lemma 4.5. $E_H \subseteq E$ and $F_H \subseteq F$. ■

Let a be a fixed element of H' . Every element of H' can be expressed in the form $a + x$ for some $x \in H$.

Lemma 4.6. *Let (x_n) be a sequence in $H, x \in H$. Then*

- (i) $x_n \rightarrow x$ in H if and only if $a + x_n \rightarrow a + x$ in G .
- (ii) $x_n \rightarrow x$ in H if and only if $a + x_n + a \rightarrow a + x + a$ in H .
- (iii) $(x_n) \in F_H$ if and only if some of the following conditions is satisfied
 $(a + x_n) \in F, (a + x_n + a) \in F_H, (-a + x_n + a) \in F_H$.

Proof. (i) and (ii) are easy to verification.

(iii): Let $(x_n) \in F_H$. We intend to show that $(a + x_n) \in F$. Assume that $\varepsilon \in G, \varepsilon \neq 0, [-\varepsilon, 0, \varepsilon] \in T$. Then $\varepsilon \in H$ and so $-a - \varepsilon + a \in H$. Since $(x_n) \in F_H, [-a + \varepsilon + a, 0, -a - \varepsilon + a] \in T$ implies that there exists $n_o \in N$ such that $[-a + \varepsilon + a, x_n - x_m, -a - \varepsilon + a] \in T$ for each $m, n \in N, m, n \geq n_o$. Therefore $[-\varepsilon, a + x_n - (a + x_m), \varepsilon] \in T$. From $[-\varepsilon, -x_m + x_n, \varepsilon] \in T$ it follows that $[-\varepsilon, -(a + x_m) + a + x_n, \varepsilon] \in T$. We conclude that $(a + x_n) \in F$.

The converse and remaining cases are similar. ■

Lemma 4.7. *G is C -complete if and only if H is C -complete.*

Proof. Let G be C -complete and let $(x_n) \in F_H$. In view of Lemma 4.5, we get $(x_n) \in F$. Hence there exists $x \in G$ with $x_n \rightarrow x$ in G . Applying Lemma 4.4 (ii) and Lemma 4.4 (iv), we obtain $x \in H$ and $x_n \rightarrow x$ in H . Hence H is C -complete.

Let H be C -complete and let $(x_n) \in F$. From Lemma 4.4 (iii), we infer that there exists $n_o \in N$ such that either $(x_n^o) \in F_H$ or $(x_n^o) \in H'$. Assume that $(x_n^o) \in F_H$. Then $x_n^o \rightarrow x$ in H . With respect to Lemma 4.4 (iv), $x_n^o \rightarrow x$ in G . This yields that $x_n \rightarrow x$ in G . Assume that $(x_n^o) \in H'$. There exists $(h_n^o) \in H$ with $x_n^o = a + h_n^o$ for each $n \in N$. Since $(a + h_n^o) \in F$, Lemma 4.6 (iii) implies that $(h_n^o) \in F_H$. Hence, $h_n^o \rightarrow h$ in H and by Lemma 4.6 (i) $a + h_n^o \rightarrow a + h$ in G . We conclude now that $x_n \rightarrow a + h$ in G and the proof is complete. ■

The following result is an immediate consequence of Lemmas 4.6 and 4.7.

Lemma 4.8. *Let G be a subgroup of a half lc-group G_1 . Then G_1 is a Cantor extension of G if and only if $G_1 \uparrow$ is a Cantor extension of H . ■*

Investigating a Cantor extension of G , two cases are distinguished: $H_o \neq \{0\}$ and $H_o = \{0\}$ (H_o is as in the section 2).

5. THE CASE $H_o \neq \{0\}$

In the whole section we suppose that $H_o \neq \{0\}$. Since H_o is infinite, G is infinite as well.

We form the sets

$$(B) \quad \begin{aligned} a + H^* &= \{a + (x_n)^* : (x_n)^* \in H^*\}, \\ C_h(G) &= H^* \cup (a + H^*). \end{aligned}$$

Assume that $(x_n) \in F_H$. With respect to Lemma 4.6 (iii), we get $(a + x_n + a) \in F_H$ and $(-a + x_n + a) \in F_H$.

We intend to define a group operation $+$ and a ternary relation T^h on $C_h(G)$. Let $(x_n)^*, (y_n)^*, (z_n)^* \in H^*$.

The operation $+$ on $C_h(G)$ is defined to coincide with the operation $+$ on H^* defined in the section 2, i.e., we put

$$(x_n)^* + (y_n)^* = (x_n + y_n)^*.$$

Further, we put

$$\begin{aligned}
(a + (x_n)^*) + (a + (y_n)^*) &= (a + x_n + a + y_n)^*, \\
(x_n)^* + (a + (y_n)^*) &= a + (-a + x_n + a + y_n)^*, \\
(a + (x_n)^*) + (y_n)^* &= a + (x_n + y_n)^*.
\end{aligned}$$

We define the ternary relation T^h on $C_h(G)$ in such a way that T^h coincides with T^* on H^* .

Further, we put

$$[a + (x_n)^*, a + (y_n)^*, a + (z_n)^*] \in T^h \text{ if } [(z_n)^*, (y_n)^*, (x_n)^*] \in T^*.$$

If p, q and r are distinct elements of $C_h(G)$ such that $[p, q, r] \in T^h$, then either $\{p, q, r\} \subseteq H^*$ or $\{p, q, r\} \subseteq a + H^*$.

Lemma 5.1. $(C_h(G); +)$ is a group.

Proof. First, we verify that the operation $+$ is associative. Only three cases are considered. The remaining cases are similar.

$$\begin{aligned}
((a + (x_n)^*) + (a + (y_n)^*)) + (a + (z_n)^*) &= (a + x_n + a + y_n)^* + (a + (z_n)^*) = \\
a + (-a + a + x_n + a + y_n + a + z_n)^* &= a + (x_n + a + y_n + a + z_n)^*, \\
(a + (x_n)^*) + ((a + (y_n)^*) + (a + (z_n)^*)) &= (a + (x_n)^*) + (a + y_n + a + z_n)^* = \\
a + (x_n + a + y_n + a + z_n)^*. &
\end{aligned}$$

Hence,

$$\begin{aligned}
((a + (x_n)^*) + (a + (y_n)^*)) + (a + (z_n)^*) &= (a + (x_n)^*) + ((a + (y_n)^*) + (a + (z_n)^*)). \\
((a + (x_n)^*) + (a + (y_n)^*)) + (z_n)^* &= (a + x_n + a + y_n)^* + (z_n)^* = (a + x_n + a + \\
y_n + z_n)^*, (a + (x_n)^*) + ((a + (y_n)^*) + (z_n)^*) &= (a + (x_n)^*) + (a + (y_n + z_n)^*) = \\
(a + x_n + a + y_n + z_n)^*. &
\end{aligned}$$

$$\begin{aligned}
\text{Thus, } ((a + (x_n)^*) + (a + (y_n)^*)) + (z_n)^* &= (a + (x_n)^*) + ((a + (y_n)^*) + (z_n)^*). \\
((x_n)^* + (y_n)^*) + (a + (z_n)^*) &= (x_n + y_n)^* + (a + (z_n)^*) = a + (-a + x_n + y_n + \\
a + z_n)^*, (x_n)^* + ((y_n)^* + (a + (z_n)^*)) &= (x_n^* + (a + (-a + y_n + a + z_n)^*)) = \\
a + (-a + x_n + a - a + y_n + a + z_n)^* &= a + (-a + x_n + y_n + a + z_n)^*.
\end{aligned}$$

Therefore, $((x_n)^* + (y_n)^*) + (a + (z_n)^*) = (x_n)^* + ((y_n)^* + (a + (z_n)^*))$.

Now, we show that every element of $C_h(G)$ has an inverse in $C_h(G)$. It suffices to consider elements of $a + H^*$. Assume that $a + (x_n)^* \in a + H^*$. Then $a + (-a - x_n - a)^* \in a + H^*$ and it is the inverse to $a + (x_n)^*$ in $C_h(G)$. ■

Lemma 5.2. Let $(x_n)^*, (y_n)^*, (z_n)^* \in H^*$. Then $[(x_n)^*, (y_n)^*, (z_n)^*] \in T^*$ if and only if some of the following conditions is satisfied:

- (i) $[(-a + z_n + a)^*, (-a + y_n + a)^*, (-a + x_n + a)^*] \in T^*$,
- (ii) $[(a + z_n + a)^*, (a + y_n + a)^*, (a + x_n + a)^*] \in T^*$.

Proof. (i): Assume that $[(x_n)^*, (y_n)^*, (z_n)^*] \in T^*$. Hence there exists $n_o \in N$ such that $[x_n, y_n, z_n] \in T$ for each $n \in N, n \geq n_o$. This yields that $[-a + z_n + a, -a + y_n + a, -a + x_n + a] \in T$ for each $n \in N, n \geq n_o$. According to Lemma 4.6 (iii), we have $(-a + z_n + a), (-a + y_n + a), (-a + x_n + a) \in F_H$. We conclude that $[(-a + z_n + a)^*, (-a + y_n + a)^*, (-a + x_n + a)^*] \in T^*$.

The converse and (ii) are similar. ■

Lemma 5.3. *Let $(x_n)^*, (y_n)^*, (z_n)^*, (u_n)^* \in H^*$.*

- (i) *If $[(x_n)^*, (y_n)^*, (z_n)^*] \in T^*$, then $[(x_n)^* + (u_n)^*, (y_n)^* + (u_n)^*, (z_n)^* + (u_n)^*] \in T^*$ and $[(x_n)^* + (a + (u_n)^*), (y_n)^* + (a + (u_n)^*), (z_n)^* + (a + (u_n)^*)] \in T^h$.*
- (ii) *If $[a + (x_n)^*, a + (y_n)^*, a + (z_n)^*] \in T^h$, then $[(a + (x_n)^*) + (u_n)^*, (a + (y_n)^*) + (u_n)^*, (a + (z_n)^*) + (u_n)^*] \in T^h$ and $[(a + (x_n)^*) + (a + (u_n)^*), (a + (y_n)^*) + (a + (u_n)^*), (a + (z_n)^*) + (a + (u_n)^*)] \in T^*$.*

Proof. (i): Assume that $[(x_n)^*, (y_n)^*, (z_n)^*] \in T^*$. The first part of the assertion follows from the fact that H^* is an lc -group. Now, we prove the second part. From Lemma 5.2 (i), we infer that $[(-a + z_n + a)^*, (-a + y_n + a)^*, (-a + x_n + a)^*] \in T^*$. Then $[(-a + z_n + a)^* + (u_n)^*, (-a + y_n + a)^* + (u_n)^*, (-a + x_n + a)^* + (u_n)^*] \in T^*$, $[(-a + z_n + a + u_n)^*, (-a + y_n + a + u_n)^*, (-a + x_n + a + u_n)^*] \in T^*$. Hence $[a + (-a + x_n + a + u_n)^*, a + (-a + y_n + a + u_n)^*, a + (-a + z_n + a + u_n)^*] \in T^h$, i.e., $[(x_n)^* + (a + (u_n)^*), (y_n)^* + (a + (u_n)^*), (z_n)^* + (a + (u_n)^*)] \in T^h$.

The proof of (ii) is analogous. ■

Lemma 5.4. *Let $(x_n)^*, (y_n)^*, (z_n)^*, (u_n)^* \in H^*$.*

- (i) *If $[(x_n)^*, (y_n)^*, (z_n)^*] \in T^*$, then $[(u_n)^* + (x_n)^*, (u_n)^* + (y_n)^*, (u_n)^* + (z_n)^*] \in T^*$ and $[(a + (u_n)^*) + (z_n)^*, (a + (u_n)^*) + (y_n)^*, (a + (u_n)^*) + (x_n)^*] \in T^h$.*
- (ii) *If $[a + (x_n)^*, a + (y_n)^*, a + (z_n)^*] \in T^h$, then $[(u_n)^* + (a + (x_n)^*), (u_n)^* + (a + (y_n)^*), (u_n)^* + (a + (z_n)^*)] \in T^h$ and $[(a + (u_n)^*) + (a + (z_n)^*), (a + (u_n)^*) + (a + (y_n)^*), (a + (u_n)^*) + (a + (x_n)^*)] \in T^*$.*

Proof. (i) Assume that $[(x_n)^*, (y_n)^*, (z_n)^*] \in T^*$. The first assertion holds because of the fact that H^* is an lc -group. Now, we prove the second assertion. The assumption implies that $[(u_n)^* + (x_n)^*, (u_n)^* + (y_n)^*, (u_n)^* + (z_n)^*] \in T^*$ and so $[(u_n + x_n)^*, (u_n + y_n)^*, (u_n + z_n)^*] \in T^*$. Whence $[a + (u_n + z_n)^*, a + (u_n + y_n)^*, a + (u_n + x_n)^*] \in T^h$. Thus $[(a + (u_n)^*) + (z_n)^*, (a + (u_n)^*) + (y_n)^*, (a + (u_n)^*) + (x_n)^*] \in T^h$.

To prove (ii), we proceed in a similar way. ■

From Lemma 5.4 and (B), we infer the validity of the following result:

Lemma 5.5. $C_h(G) \uparrow = H^*$, $C_h(G) \downarrow = a + H^*$ and $C_h(G) = C_h(G) \uparrow \cup C_h(G) \downarrow$. ■

Since T is nonempty, T^h is nonempty as well. Then Lemmas 5.1, 5.3 and 5.5 yield:

Lemma 5.6. $(C_h(G), +, T^h)$ is a half lc -group. ■

Let $x \in H$. Define the mapping ψ from G into $C_h(G)$ by

$$\psi(x) = (x, x, \dots)^*, \quad \psi(a + x) = a + \psi(x).$$

Then ψ is an isomorphism of the half lc -group G into $C_h(G)$. In the next, we identify x and $\psi(x)$ for each $x \in H$. Then G is a half lc -subgroup of $C_h(G)$. Since H^* is a Cantor extension of H , from Lemma 4.8, we conclude.

Theorem 5.7. $C_h(G)$ is a Cantor extension of G . ■

Remark that it is easy to verify that (A) implies $X_n \rightarrow X$ and $a + X_n \rightarrow a + X$ in G .

Theorem 5.8. Let G_1 and G_2 be Cantor extensions of G . Then there exists an isomorphism f from the half lc -group G_1 , onto G_2 such that $f(x) = x$ for each $x \in G$.

Proof. With respect to 4.8, $G_1 \uparrow$ and $G_2 \uparrow$ are Cantor extension of H . By Theorem 2.6, there exists an isomorphism ϕ from $G_1 \uparrow$ onto $G_2 \uparrow$ with $\phi(x) = x$ for any $x \in H$.

Choose an arbitrary element $z \in G_1 \uparrow$. The mapping $f : G_1 \rightarrow G_2$ defined by $f(z) = \phi(z)$ and $f(a + z) = a + \phi(z)$ is an isomorphism of the half lc -group G_1 onto G_2 and $f(a + x) = a + \phi(x) = a + x$ for each $x \in H$. ■

A half lc -group $C_h(G)$ corresponds to an element $a \in H'$. Let $a' \in H'$, $a' \neq a$. Then the half lc -group $(C'_h(G); +', T')$ corresponding to a' can be constructed formally in the same way ($+$, T_h and a are replaced by $+'$, T' and a' , respectively). Therefore, the operations $+$ and $+'$ (relations T^h and T') coincide on G and H^* . From Theorems 5.7 and 5.8, it follows that $C_h(G)$ and $C'_h(G)$ are isomorphic half lc -groups. Moreover, we have:

Lemma 5.9. *A half lc -group $C_h(G) = C'_h(G)$.*

Proof. For each $(x_n)^* \in H^*$ we get $a + (x_n)^* = a' +' (-a' + a + x_n)^*$. Hence, $a + H^* \subseteq a' +' H^*$. Analogously, we get $a' +' H^* \subseteq a + H'$. Therefore, the set $C_h(G) = C'_h(G)$.

Evidently, that relations T^h and T' coincide. Now we show that group operations $+$ on $C_h(G)$ and $+'$ on $C'_h(G)$ coincide.

Let $(x_n)^*, (y_n)^* \in H^*$. Then

$$\begin{aligned} (a + (x_n)^*) + (a + (y_n)^*) &= (a + x_n + a + y_n)^* = (a' - a' + a + x_n + a' - a' + a + y_n)^* = \\ &= (a' +' (-a' + a + x_n)^*) +' (a' +' (-a' + a + y_n)^*); \\ (x_n)^* + (a + (y_n)^*) &= a + (-a + x_n + a + y_n)^* = a' +' (-a' + a - a + x_n + a + y_n)^* = \\ &= a' +' (-a' + x_n + a' - a' + a + y_n)^* = (x_n)^* +' (a' +' (-a' + a + y_n)^*); \\ (a + (x_n)^*) + (y_n)^* &= a + (x_n + y_n)^* = a' +' (-a' + a + x_n + y_n)^* = \\ &= a' +' ((-a' + a + x_n)^* + (y_n)^*). \quad \blacksquare \end{aligned}$$

6. THE CASE $H_o = \{0\}$

In this section, we assume that $H_o = \{0\}$. Then H can be considered as a subgroup of K .

Assume that G is a finite half lc -group. Then H is a finite lc -group. With respect to Lemmas 2.4 and 4.7, we obtain:

Lemma 6.1. *Let G be a finite half lc -group. Then G is C -complete.* ■

Now, assume that G is an infinite half lc -group. Then H is an infinite lc -group.

Let a be a fixed element of H' . We denote

$$\begin{aligned} a + K &= \{a + x : x \in K\}; \\ C_h(G) &= K \cup (a + K). \end{aligned}$$

We will define a group operation $+$ and a ternary relation T^h on $C_h(G)$.

Let $x, y, z \in K$. From Lemma 2.5, we infer that there are fundamental sequences (x_n) and (y_n) in G such that $\lim x_n = x, \lim y_n = y$ in K .

The operation $x + y$ on $C_h(G)$ coincides with $x + y$ on K .

Further, we put

$$\begin{aligned}(a + x) + (a + y) &= \lim(a + x_n + a + y_n), \\ x + (a + y) &= a + \lim(-a + x_n + a + y_n), \\ (a + x) + y &= a + \lim(x_n + y_n).\end{aligned}$$

Limits are taken into account in K . The operation $+$ is correctly defined.

The ternary relation T^h on $C_h(G)$ is defined in the following way:

$$T^h \text{ coincides with } T_1 \text{ on } K.$$

Further, we put

$$[a + x, a + y, a + z] \in T^h \text{ if } [x, y, z] \in T_1,$$

if $p, q, r \in C_h(G)$, $[p, q, r] \in T^h$, then either $\{p, q, r\} \subseteq K$ or $\{p, q, r\} \subseteq a + K$.

It is a routine to verify that the following assertion is true:

Lemma 6.2. $(C_h(G), +, T^h)$ is a half lc -group, $C_h(G) \uparrow = K$, $C_h(G) \downarrow = a + K$. ■

From Lemmas 2.5 and 4.7, it follows:

Lemma 6.3. Let G be an infinite half lc -group. Then $C_h(G)$ is a Cantor extension of G . ■

Let a' and $C'_h(G)$ be as in the Section 5.

Remark 6.4. It is easy to verify that Theorem 5.8 and Lemma 5.9 are valid also in the case $H_o = \{0\}$.

From Lemmas 4.7, 2.5 and Theorem 2.6, we obtain:

Lemma 6.5. Let G be an infinite half lc -group. Then G is C -complete if and only if H is isomorphic to K . ■

Let G be an arbitrary lc -group as in the section 4 (neither $H_o \neq \{0\}$ nor $H_o = \{0\}$ is supposed). From Lemmas 6.1, 6.5 and 2.3, we conclude:

Theorem 6.6. *Let G be a half lc-group such that H is abelian and $H' \neq \emptyset$. Then G is C -complete if and only if some of the following conditions is fulfilled:*

- (i) G is finite;
- (ii) H is isomorphic to K ;
- (iii) $H_o \neq \{0\}$ and H_o is C -complete. ■

By summarizing Theorem 5.7, Lemma 6.3, Theorem 5.8, and Remark 6.4 we get:

Theorem 6.7. *Let G be a half lc-group such that H is abelian and $H' \neq \emptyset$. Then*

- (i) *There exists a Cantor extension of G .*
- (ii) *If G_1 and G_2 are Cantor extensions of G , then there exists an isomorphism from the half lc-group G_1 onto G_2 leaving all elements of G fixed.*

REFERENCES

- [1] Š. Černák, *Cantor extension of an abelian cyclically ordered group*, Math. Slovaca **39** (1989), 31–41.
- [2] C.J. Everett, *Sequence completion of lattice moduls*, Duke Math. J. **11** (1944), 109–119.
- [3] M. Giraudet and F. Lucas, *Groupe à moitié ordonnés*, Fund. Math. **139** (1991), 75–89.
- [4] J. Jakubík and G. Pringerová, *Representations of cyclically ordered groups*, Časopis pěst. Mat. **113** (1988), 184–196.
- [5] J. Jakubík and G. Pringerová, *Radical classes of cyclically ordered groups*, Math. Slovaca **38** (1988), 255–268.
- [6] J. Jakubík, *On half lattice ordered groups*, Czechoslovak Math. J. **46** (1996), 745–767.
- [7] J. Jakubík, *Lexicographic products of half linearly ordered groups*, Czechoslovak Math. J. **51** (2001), 127–138.
- [8] J. Jakubík, *On half cyclically ordered groups*, Czechoslovak Math. J. (to appear).

- [9] V. Novák, *Cuts in cyclically ordered sets*, Czechoslovak Math. J. **34** (1984), 322–333.
- [10] V. Novák and M. Novotný, *On representations of cyclically ordered sets*, Czechoslovak Math. J. **39** (1989), 127–132.
- [11] A. Quilot, *Cyclic orders*, European J. Combin. **10** (1989), 477–488.
- [12] L. Rieger, *On ordered and cyclically ordered groups I., II., III.*, Věstník Král. České spol. Nauk (Czech) **1946**, 1–31; **1947**, 1–33; **1948**, 1–26.
- [13] S. Świerczkowski, *On cyclically ordered groups*, Fund. Math. **47** (1959), 161–166.
- [14] D.R. Ton, *Torsion classes and torsion prime selectors of hl-groups*, Math. Slovaca **50** (2000), 31–40.

Received 7 June 1999
Revised 30 April 1996