

## LEXICO EXTENSION AND A CUT COMPLETION OF A HALF $l$ -GROUP

ŠTEFAN ČERNÁK

*Department of Mathematics, Faculty of Civil Engineering*  
*Technical University*  
*Vysokoškolská 4, SK-042 02 Košice, Slovakia*  
**e-mail:** svfkm@tuke.sk

AND

MILAN DEMKO

*Department of Mathematics, FHPV PU*  
*17 Novembra 1, SK-081 16 Prešov, Slovakia*  
**e-mail:** demko@unipo.sk

### Abstract

The cut completion of an  $hl$ -group  $G$  with the abelian increasing part is investigated under the assumption that  $G$  is a lexico extension of its  $hl$ -subgroup.

**Keywords:** half lattice ordered group,  $l$ -group, cut completion, lexico extension.

**2000 AMS Mathematics Subject Classification:** Primary 06F15; Secondary 06F20.

### 0. INTRODUCTION

The notion of a half  $l$ -group as a generalization of the notion of an  $l$ -group was introduced and studied by M. Giraudet and F. Lucas [4].

R.N. Ball [1] has defined the notion of a cut completion of an  $l$ -group.

In this paper we define the notions of a cut completion and a lexico extension of a half  $l$ -group. We prove a theorem on a cut completion of a half  $l$ -group having an abelian increasing part which can be expressed as a nontrivial lexico extension. A particular case of this theorem is a result of J. Jakubík [5] dealing with a cut completion of an abelian  $l$ -group.

## 1. PRELIMINARIES

Let  $G$  be an abelian  $l$ -group.  $G$  is called a *lexico extension of its  $l$ -subgroup*  $A \neq \{0\}$  if

- (i)  $A$  is a convex  $l$ -subgroup of  $G$ ,
- (ii) if  $0 < g \in G$ ,  $g \notin A$ , then  $g > a$  for each  $a \in A$ .

If  $G$  is a lexico extension of  $A$ , we shall write  $G = \langle A \rangle$ . If  $G = \langle A \rangle$ , then  $A$  is an  $l$ -ideal of  $G$  and (cf. [3] and [2])

- (a)  $A$  is comparable to all convex  $l$ -subgroups of  $G$  (i.e., if  $A'$  is a convex  $l$ -subgroup of  $G$  then either  $A' \subseteq A$  or  $A \subseteq A'$ ).
- (b)  $G/A$  is a linearly ordered group.

Let  $G$  be a group and a partially ordered set. Set

$$G \uparrow = \{g \in G : x \leq y \Rightarrow g + x \leq g + y \text{ for all } x, y \in G\},$$

$$G \downarrow = \{g \in G : x \leq y \Rightarrow g + x \geq g + y \text{ for all } x, y \in G\}.$$

$G \uparrow$  ( $G \downarrow$ ) is called the *increasing* (*decreasing*) part of  $G$ .

$G$  is said to be a *half  $l$ -group* (abbreviated to an *hl-group*) if the following conditions are satisfied (cf. [4]):

- (i) the partial order  $\leq$  on  $G$  is non-trivial,
- (ii) if  $x, y, g \in G$  and  $x \leq y$ , then  $x + g \leq y + g$ ,
- (iii)  $G = G \uparrow \cup G \downarrow$ ,
- (iv)  $G \uparrow$  is an  $l$ -group.

If  $G \uparrow$  is a linearly ordered group, then  $hl$ -group  $G$  will be called a *half linearly ordered group*.

Every  $l$ -group  $G \neq \{0\}$  is a special case of an  $hl$ -group with  $G \downarrow = \emptyset$ .

We denote by  $\mathcal{HL}$  the class of all  $hl$ -groups that fail to be  $l$ -groups.

The following results will be applied in the next.

**Proposition 1.1** (cf. [4]). *Let  $G \in \mathcal{HL}$ . Then*

- (i)  $G \uparrow$  is a subgroup of the group  $G$  and  $G \uparrow$  has index 2,
- (ii)  $G \uparrow$  and  $G \downarrow$  are isomorphic groups and also dually isomorphic lattices,
- (iii) if  $x \in G \uparrow$  and  $y \in G \downarrow$ , then  $x$  and  $y$  are incomparable,
- (iv) the set  $\{g \in G : g \neq 0, 2g = 0\}$  is nonempty.

Let  $G$  be an  $hl$ -group. A subgroup  $A \neq \{0\}$  of  $G$  is called a half  $l$ -subgroup (abbreviated to an  $hl$ -subgroup) if  $A \uparrow = A \cap G \uparrow$  is an  $l$ -subgroup of  $G \uparrow$ . If  $A$  is an  $hl$ -subgroup (proper  $hl$ -subgroup) of  $G$  we use the notation  $A \leq G$  ( $A < G$ ). We say that an  $hl$ -subgroup  $A$  of  $G$  is convex in  $G$  if  $A \uparrow$  is convex in  $G \uparrow$ . A convex  $hl$ -subgroup  $A$  of  $G$  is said to be an  $hl$ -ideal of  $G$  if  $A \uparrow$  is a normal subgroup of  $G$ . According to 1.1  $G \uparrow$  is an  $hl$ -ideal of  $G$ .

Let  $G$  be an  $hl$ -group,  $G \in \mathcal{HL}$  and  $A \uparrow$  an  $hl$ -ideal of  $G$ ,  $A \in \mathcal{HL}$ . We can form the factor group  $\overline{G} = G/A \uparrow$ . For elements  $g_1 + A \uparrow, g_2 + A \uparrow \in \overline{G}$ , we put  $g_1 + A \uparrow \leq g_2 + A \uparrow$  if and only if there exist  $g'_1 \in g_1 + A \uparrow$  and  $g'_2 \in g_2 + A \uparrow$  with  $g'_1 \leq g'_2$ . Then  $\overline{G}$  is a partially ordered set and to each  $g'_1 \in g_1 + A \uparrow$  there exists  $g'_2 \in g_2 + A \uparrow$  such that  $g'_1 \leq g'_2$ . It can be easily verified that if  $A < G$ , then  $\overline{G}$  is an  $hl$ -group with the increasing part  $\overline{G} \uparrow = \{g + A \uparrow : g \in G \uparrow\}$  and decreasing part  $\overline{G} \downarrow = \{g + A \uparrow : g \in G \downarrow\}$ .

If  $A = G$  then  $\overline{G}$  is trivially ordered. Hence  $\overline{G}$  fails to be an  $hl$ -group.

A 1-1 mapping  $\varphi$  from an  $hl$ -group  $G$  onto an  $hl$ -group  $G'$  is called an  $hl$ -isomorphism if  $\varphi$  is a group homomorphism and if  $\varphi|G \uparrow$  is a lattice homomorphism of  $G \uparrow$  onto  $G' \uparrow$ .

## 2. LEXICO EXTENSION OF AN $hl$ -SUBGROUP

Let  $G$  be an  $hl$ -group,  $G \in \mathcal{HL}$  with the abelian increasing part  $G \uparrow$ . Let  $A$  be an  $hl$ -subgroup of  $G$ ,  $A \in \mathcal{HL}$ . If  $G \uparrow$  is a lexico extension of  $A \uparrow$ , then we say that  $G$  is a lexico extension of  $A$  and we express this situation by writing  $G = \langle A \rangle_h$ .

**Lemma 2.1.** *Let  $G = \langle A \rangle_h$ . Then*

- (i)  $A$  is an  $hl$ -ideal of  $G$ ,
- (ii) if  $A < G$ , then  $\overline{G} = G/A \uparrow$  is a half linearly ordered group.

**Proof.** (i) We have to show that  $A \uparrow$  is normal in  $G$ . Since  $A \uparrow$  is a convex  $l$ -subgroup of  $G \uparrow$ ,  $-g + A \uparrow + g$  ( $g \in G$ ) is a convex subset of  $G \uparrow$ . It is a routine to verify that  $-g + A \uparrow + g$  is a subgroup of  $G \uparrow$ . Let  $-g + a_1 + g$ ,  $-g + a_2 + g \in -g + A \uparrow + g$ ,  $g \in G$ , and  $a_1, a_2 \in A \uparrow$ . It is easy to verify that in  $G \uparrow$  we have  $(-g + a_1 + g) \vee (-g + a_2 + g) = -g + (a_1 \vee a_2) + g$  for each  $g \in G \uparrow$ ,  $(-g + a_1 + g) \vee (-g + a_2 + g) = -g + (a_1 \wedge a_2) + g$  for each  $g \in G \downarrow$  and dually. Hence  $-g + A \uparrow + g$  is a sublattice of  $G \uparrow$  for each  $g \in G$ . By summarizing we have that  $-g + A \uparrow + g$  is a convex  $l$ -subgroup of  $G \uparrow$  for each  $g \in G$ . By (a),  $A \uparrow$  and  $-g + A \uparrow + g$  are comparable. The fact that  $G \uparrow$  is abelian implies  $-g + A \uparrow + g = A \uparrow$  for all  $g \in G \uparrow$ . Suppose that  $g \in G \downarrow$  and  $A \uparrow \subseteq -g + A \uparrow + g$ . Let  $a \in A \uparrow$ . Then  $a = -g + a_0 + g$ , where  $a_0 \in A \uparrow$  and hence  $-g + a + g = -2g + a_0 + 2g$ . Since  $2g \in G \uparrow$ , we get  $-g + a + g \in A \uparrow$ . Thus  $-g + A \uparrow + g \subseteq A \uparrow$  for all  $g \in G$ . Therefore,  $A \uparrow$  is normal in  $G$ .

(ii) follows from the property (b) of a lexico extension.  $\blacksquare$

If  $G = \langle A \rangle_h$ , then Lemma 2.1 yields that  $A \uparrow$  is a normal subgroup of  $G$ , but  $A$  need not be normal in  $G$ .

**Examples.** Let  $M$  be the set of all functions  $f : R \rightarrow R$ ;  $f(x) = \pm x + k$ ,  $k \in R$ . If a binary operation on  $M$  is defined as a composition (i.e.,  $fg(x) = f(g(x))$  for all  $x \in R$ ) and a binary relation  $\leq$  on  $M$  is defined pointwise, then  $M$  is a half linearly ordered group, with  $M \uparrow = \{f : f(x) = x + k\}$  and  $M \downarrow = \{f : f(x) = -x + k\}$ . Now, let  $H = \{(f_1, f_2) : f_1, f_2 \in M \uparrow\}$ . For each  $(f_1, f_2), (f'_1, f'_2) \in H$  we put  $(f_1, f_2) \leq (f'_1, f'_2)$  if and only if either  $f_1 < f'_1$  or  $f_1 = f'_1$  and  $f_2 \leq f'_2$ . Then  $H$  is a linearly ordered set that is called the *lexicographic product* of the two linearly ordered sets  $M \uparrow$  and we use the denotation  $H = M \uparrow \circ M \uparrow$ . Analogously we can construct  $K = M \downarrow \circ M \downarrow$ . If a binary operation on  $H$  is defined componentwise, then  $H$  is a linearly ordered group. Therefore,  $G = H \cup K$  is a half linearly ordered group with  $G \uparrow = H, G \downarrow = K$ . Let  $A_1 = \{(id, g) \in G : g \in M \uparrow\}$  ( $id$  is an identity function) and  $A_2 = \{(g_1, g) : g_1 \text{ is a fixed element of } M \downarrow, g \in M \downarrow\}$ . Then  $A = A_1 \cup A_2$  is a half linearly ordered group,  $A \uparrow = A_1, A \downarrow = A_2$ . We have  $G = \langle A \rangle_h$ , but  $A$  fails to be normal in  $G$ . In fact, for all  $f \in M, f \neq id, g_1$  we have  $f^{-1}g_1f \neq g_1$ , thus  $(f, f)^{-1}(g_1, g)(f, f) \notin A_2$  for each  $(g_1, g) \in A_2$ .

Assume that  $G$  is an  $hl$ -group,  $G \in \mathcal{HL}$  and that  $A \in \mathcal{HL}$  is an  $hl$ -ideal of  $G$  such that  $A$  is a normal subgroup of  $G$ . Define a partial order on the factor group  $G/A$  ( and also on the factor group  $G \uparrow / A \uparrow$  ) analogously

as above on  $G/A \uparrow$ . Then  $G/A$  is a lattice ordered group. The mapping  $f : G \uparrow / A \uparrow \rightarrow G/A$  defined by  $f(g+A \uparrow) = g+A, g \in G \uparrow$  is an isomorphism of the lattice ordered group  $G \uparrow / A \uparrow$  onto  $G/A$ .

Suppose that  $G = \langle A \rangle_h$  and that  $A$  is normal in  $G$ . Then, by the property (b) of a lexico extension, we have that  $G/A$  is a linearly ordered group.

**Lemma 2.2.** *Let  $G = \langle A \rangle_h, g_1+A \uparrow, g_2+A \uparrow \in \overline{G}$ , and let  $g_1+A \uparrow < g_2+A \uparrow$ . Then  $g_1 < g_2$ .*

**Proof.** From  $g_1 + A \uparrow < g_2 + A \uparrow$  it follows that either  $g_1, g_2 \in G \uparrow$  or  $g_1, g_2 \in G \downarrow$ . Now, let  $g_1, g_2 \in G \uparrow$ . There exists  $g'_2 \in g_2 + A \uparrow$  such that  $g_1 < g'_2$ . Hence  $g'_2 \in G \uparrow$  and  $g'_2 - g_1 > 0$ . Since  $g'_2 - g_1 \notin A \uparrow$  and  $g'_2 - g_2 \in A \uparrow$ , we get  $g'_2 - g_1 > g'_2 - g_2$ . Hence  $-g_1 > -g_2$  and  $g_1 < g_2$ . Suppose that  $g_1, g_2 \in G \downarrow$ . There exists  $g''_2 \in g_2 + A \uparrow$  with  $g_1 < g''_2$ . Hence  $g''_2 \in G \downarrow, g''_2 - g_1 \in G \uparrow, g''_2 - g_1 \notin A \uparrow$  and  $g''_2 - g_1 > 0$ . Further, we have  $g''_2 - g_2 \in A \uparrow$ . Then  $g''_2 - g_1 > g''_2 - g_2$  implies that  $-g_1 < -g_2$  and so  $g_1 < g_2$ . ■

**Corollary.** *Let  $G = \langle A \rangle_h$ . Then  $G$  is a half linearly ordered group if and only if  $A$  is a half linearly ordered group.*

For the remaining part of this section, we assume that  $G, A$  and  $B$  are  $hl$ -groups from  $\mathcal{HL}$  such that

- (I)  $G \uparrow$  and  $B \uparrow$  are abelian  $l$ -groups,
- (II)  $G = \langle A \rangle_h, A < G$ ,
- (III)  $A$  is an  $hl$ -subgroup of  $B$ ,
- (IV)  $G \cap B = A$ .

According to Proposition 1.1, there exists an element  $a \in A \downarrow$  of order 2.

Form the set

$$H_0 = \{(g, b) : \text{either } g \in G \uparrow, b \in B \uparrow \text{ or } g \in G \downarrow, b \in B \downarrow\}.$$

For elements  $(g_1, b_1), (g_2, b_2) \in H_0$ , we set

$$(g_1, b_1) \equiv (g_2, b_2)$$

if  $g_1 - g_2 \in A \uparrow, b_1 - b_2 \in A \uparrow, g_1 - g_2 = b_2 - b_1$  and if either  $g_1, g_2 \in G \uparrow, b_1, b_2 \in B \uparrow$  or  $g_1, g_2 \in G \downarrow, b_1, b_2 \in B \downarrow$ .

The relation  $\equiv$  is an equivalence. It is clear that the relation  $\equiv$  is reflexive and symmetric. To establish the transitivity, suppose that  $(g_1, b_1) \equiv (g_2, b_2)$ ,  $(g_2, b_2) \equiv (g_3, b_3)$ . We will consider only the following case. Let  $g_1, g_2 \in G \downarrow$ , and  $b_1, b_2 \in B \downarrow$ . Then  $g_3 \in G \downarrow$ , and  $b_3 \in B \downarrow$ . We have  $g_1 - g_2 \in A \uparrow$ ,  $b_2 - b_1 \in A \uparrow$ ,  $g_1 - g_2 = b_2 - b_1$ ,  $g_2 - g_3 \in A \uparrow$ ,  $b_3 - b_2 \in A \uparrow$ , and  $g_2 - g_3 = b_3 - b_2$ . By (I) and (II),  $A \uparrow$  is abelian. Then  $g_1 - g_3 = (g_1 - g_2) + (g_2 - g_3) = (b_2 - b_1) + (b_3 - b_2) = (b_3 - b_2) + (b_2 - b_1) = b_3 - b_1$ . Hence  $g_1 - g_3 \in A \uparrow$  and  $b_3 - b_1 \in A \uparrow$ . Therefore  $(g_1, b_1) \equiv (g_3, b_3)$ .

Denote

$$\begin{aligned} \overline{(g, b)} &= \{(g', b') \in H_0 : (g, b) \equiv (g', b')\}, \\ H &= \{\overline{(g, b)} : (g, b) \in H_0\}. \end{aligned}$$

Let  $\overline{(g_1, b_1)}, \overline{(g_2, b_2)} \in H$ . We put

$$\overline{(g_1, b_1)} + \overline{(g_2, b_2)} = \overline{(g_1 + g_2, b_1 + b_2)}.$$

The binary operation  $+$  on  $H$  is correctly defined;  $\overline{(0, 0)}$  is a neutral element and  $\overline{(-g, -b)}$  is an inverse to  $\overline{(g, b)}$ .

We have

**Lemma 2.3.**  $(H, +)$  is a group. ■

Let  $\overline{(g_1, b_1)}, \overline{(g_2, b_2)} \in H$ . We put

$$\overline{(g_1, b_1)} \leq \overline{(g_2, b_2)}$$

if either  $g_1 < g_2$  and  $g_1 - g_2 \notin A \uparrow$  or  $g_1 - g_2 \in A \uparrow$  and  $g_1 - g_2 \leq b_2 - b_1$ .

The definition implies that either  $g_1, g_2 \in G \uparrow$  or  $g_1, g_2 \in G \downarrow$ . Now we verify that the relation  $\leq$  is correctly defined. Let  $\overline{(g'_1, b'_1)} = \overline{(g_1, b_1)}$ ,  $\overline{(g'_2, b'_2)} = \overline{(g_2, b_2)}$ .

Assume that  $g_1 < g_2$ ,  $g_1 - g_2 \notin A \uparrow$ . Then  $g_1 + A \uparrow < g_2 + A \uparrow$  and  $g_1, g_2 \in G \uparrow$  or  $g_1, g_2 \in G \downarrow$ . Since  $g_1 - g'_1 \in A \uparrow$  and  $g_2 - g'_2 \in A \uparrow$ , we get  $g_1 + A \uparrow = g'_1 + A \uparrow$  and  $g_2 + A \uparrow = g'_2 + A \uparrow$ . With respect to Lemma 2.2, we get  $g'_1 < g'_2$ . Suppose that  $g'_1 - g'_2 \in A \uparrow$ . Then  $g_1 - g_2 = (g_1 - g'_1) + (g'_1 - g'_2) + (g'_2 - g_2) \in A \uparrow$ , a contradiction. Hence  $g'_1 - g'_2 \notin A \uparrow$ .

Assume that  $g_1 - g_2 \in A \uparrow$ ,  $g_1 - g_2 \leq b_2 - b_1$ . Then  $g'_1 - g'_2 = (g'_1 - g_1) + (g_1 - g_2) + (g_2 - g'_2) \leq (b_1 - b'_1) + (b_2 - b_1) + (b'_2 - b_2) = (b_1 - b'_1) + (b'_2 - b_2) + (b_2 - b_1) = (b_1 - b'_1) + (b'_2 - b_1) = (b'_2 - b_1) + (b_1 - b'_1) = b'_2 - b'_1$ . We also have shown that  $g'_1 - g'_2 \in A \uparrow$ .

It is evident that the relation  $\leq$  is reflexive.

Let  $\overline{(g_1, b_1)} \leq \overline{(g_2, b_2)}$ ,  $\overline{(g_2, b_2)} \leq \overline{(g_1, b_1)}$ . Then  $g_1 - g_2 \in A \uparrow$ ,  $g_1 - g_2 \leq b_2 - b_1$  and  $g_2 - g_1 \leq b_1 - b_2$ . Hence  $g_1 - g_2 = b_2 - b_1$  and so  $\overline{(g_1, b_1)} = \overline{(g_2, b_2)}$ .

The antisymmetry is satisfied.

Let  $\overline{(g_1, b_1)} \leq \overline{(g_2, b_2)}$ ,  $\overline{(g_2, b_2)} \leq \overline{(g_3, b_3)}$ .

( $\alpha$ ) Assume that  $g_1 < g_2$ ,  $g_1 - g_2 \notin A \uparrow$ ,  $g_2 < g_3$ , and  $g_2 - g_3 \notin A \uparrow$ . We will consider only the case that  $g_1, g_2 \in G \downarrow$ . Then also  $g_3 \in G \downarrow$  and  $g_1 < g_3$ . Assume that  $g_1 - g_3 \in A \uparrow$ . Then  $g_1 + A \uparrow = g_3 + A \uparrow$ . Since  $g_1 + A \uparrow$  is a convex subset of  $G \downarrow$  and  $g_1 < g_2 < g_3$ , we obtain  $g_2 \in g_1 + A \uparrow$ . Hence  $g_1 - g_2 \in A \uparrow$ , a contradiction.

( $\beta$ ) Assume that  $g_1 - g_2 \in A \uparrow$ ,  $g_1 - g_2 \leq b_2 - b_1$ ,  $g_2 - g_3 \in A \uparrow$ , and  $g_2 - g_3 \leq b_3 - b_2$ . Then  $g_1 - g_3 = (g_1 - g_2) + (g_2 - g_3) \in A \uparrow$ , and  $g_1 - g_3 = (g_1 - g_2) + (g_2 - g_3) \leq (b_2 - b_1) + (b_3 - b_2) = (b_3 - b_2) + (b_2 - b_1) = b_3 - b_1$ .

( $\gamma$ ) Assume that  $g_1 < g_2$ ,  $g_1 - g_2 \notin A \uparrow$ ,  $g_2 - g_3 \in A \uparrow$ , and  $g_2 - g_3 \leq b_3 - b_2$ . We will consider only the case  $g_1, g_2 \in G \downarrow$ . Hence  $g_3 \in G \downarrow$ ,  $g_2 - g_1 > 0$  and  $g_2 - g_1 \notin A \uparrow$ . From this it follows that  $g_2 - g_1 > g_2 - g_3$ ,  $-g_1 < -g_3$  and  $g_1 < g_3$ . Suppose that  $g_1 - g_3 \in A \uparrow$ . Then  $g_1 - g_2 = (g_1 - g_3) + (g_3 - g_2) \in A \uparrow$ , a contradiction.

( $\delta$ ) Suppose that  $g_1 - g_2 \in A \uparrow$ ,  $g_1 - g_2 \leq b_2 - b_1$ ,  $g_2 < g_3$ , and  $g_2 - g_3 \notin A \uparrow$ . The case is analogous to ( $\gamma$ ).

In all cases ( $\alpha$ )-( $\delta$ ) we get  $\overline{(g_1, b_1)} \leq \overline{(g_3, b_3)}$ , i.e the relation  $\leq$  is transitive.

We have shown that the following lemma is valid.

**Lemma 2.4.**  $(H, \leq)$  is a partially ordered set. ■

**Lemma 2.5.** Let  $\overline{(g_1, b_1)}, \overline{(g_2, b_2)}, \overline{(g_3, b_3)} \in H$ ,  $\overline{(g_1, b_1)} \leq \overline{(g_2, b_2)}$ . Then  $\overline{(g_1, b_1)} + \overline{(g_3, b_3)} \leq \overline{(g_2, b_2)} + \overline{(g_3, b_3)}$ .

**Proof.** We will consider only the case that  $g_1, g_2, g_3 \in G \downarrow$ .

Suppose that  $g_1 < g_2$  and  $g_1 - g_2 \notin A \uparrow$ . Then  $g_1 + g_3 < g_2 + g_3$  and  $(g_1 + g_3) - (g_2 + g_3) = g_1 - g_2 \notin A \uparrow$ .

Assume that  $g_1 - g_2 \in A \uparrow$  and  $g_1 - g_2 \leq b_2 - b_1$ . Then  $(g_1 + g_3) - (g_2 + g_3) \in A \uparrow$  and  $(g_1 + g_3) - (g_2 + g_3) = g_1 - g_2 \leq b_2 - b_1 = (b_2 + b_3) - (b_1 + b_3)$ . Therefore  $\overline{(g_1, b_1)} + \overline{(g_3, b_3)} \leq \overline{(g_2, b_2)} + \overline{(g_3, b_3)}$ . ■

Form the sets

$$H \uparrow = \{\overline{(g, b)} : g \in G \uparrow, b \in B \uparrow\}, \quad H \downarrow = \{\overline{(g, b)} : g \in G \downarrow, b \in B \downarrow\}.$$

Then we have

**Lemma 2.6.**  $H = (H \uparrow) \cup (H \downarrow)$ . ■

**Lemma 2.7.**  $H \uparrow$  is an increasing part and  $H \downarrow$  is a decreasing part of  $H$ .

**Proof.** Assume that  $\overline{(g_1, b_1)}, \overline{(g_2, b_2)} \in H$ ,  $\overline{(g_1, b_1)} \leq \overline{(g_2, b_2)}$  and  $\overline{(g_3, b_3)} \in H \downarrow$ . We intend to show that  $H \downarrow$  is a decreasing part of  $H$ , i.e., that  $\overline{(g_3, b_3)} + \overline{(g_2, b_2)} \leq \overline{(g_3, b_3)} + \overline{(g_1, b_1)}$  is valid.

Let  $g_1 < g_2$ ,  $g_1 - g_2 \notin A \uparrow$ . Then  $g_3 + g_1 > g_3 + g_2$ . Suppose that  $(g_3 + g_2) - (g_3 + g_1) \in A \uparrow$ . With respect to Lemma 2.1,  $A \uparrow$  is normal in  $G$ . Thus  $g_2 - g_1 \in -g_3 + A \uparrow + g_3 \subseteq A \uparrow$ . Hence  $g_1 - g_2 \in A \uparrow$ , a contradiction.

Let  $g_1 - g_2 \in A \uparrow$  and  $g_1 - g_2 \leq b_2 - b_1$ . By using the normality of  $A \uparrow$  in  $G$ , we obtain  $g_3 + g_2 - (g_3 + g_1) = g_3 + (g_2 - g_1) - g_3 \in A \uparrow$ . There exist elements  $g'_3 \in G \uparrow$  and  $b'_3 \in B \uparrow$  such that  $g_3 = a + g'_3$ ,  $b_3 = a + b'_3$ . From  $g_2 - g_1 \geq b_1 - b_2$ , it follows  $a + g'_3 - g'_3 + g_2 - g_1 + a \leq a + b'_3 - b'_3 + b_1 - b_2 + a$ ,  $(a + g'_3 + g_2) - (a + g'_3 + g_1) \leq (a + b'_3 + b_1) - (a + b'_3 + b_2)$  and  $(g_3 + g_2) - (g_3 + g_1) \leq (b_3 + b_1) - (b_3 + b_2)$ .

In an analogous way, we show that  $H \uparrow$  is an increasing part of  $H$ . ■

$H \uparrow$  is a group (subgroup of  $H$ ) and a partially ordered set (a partial order is inherited from  $H$ ). Then according to Lemmas 2.5 and 2.7,  $H \uparrow$  is a partially ordered group.

**Lemma 2.8.**  $H \uparrow$  is an  $l$ -group.

**Proof.** It is sufficient to prove that there exists  $\sup\{\overline{(0, 0)}, \overline{(g, b)}\}$  for each  $\overline{(g, b)} \in H \uparrow$ . If  $g \notin A \uparrow$  then  $g > 0$  or  $g < 0$ . Hence  $\overline{(g, b)}$  and  $\overline{(0, 0)}$  are comparable. If  $g \in A \uparrow$  then  $g + b \in B \uparrow$  and  $\overline{(g, b)} = \overline{(0, g + b)}$ . Let



$b' = \sup\{0, g + b\}$  in  $B \uparrow$ . By using the same procedure as in the proof of Lemma 2.4 in [5], we obtain  $\overline{(0, b')} = \sup\{\overline{(0, 0)}, \overline{(g, b)}\}$ . ■

From Lemmas 2.3–2.8 it follows

**Lemma 2.9.**  *$H$  is an  $hl$ -group,  $H \in \mathcal{HL}$ .* ■

Recall that there is  $a \in A \downarrow$ , an element of order 2 (by Proposition 1.1), and that, by (IV),  $A \subseteq B$ . Define the mapping  $\varphi$  of  $G$  into  $H$  by  $\varphi(g) = \overline{(g, 0)}$  if  $g \in G \uparrow$  and  $\varphi(g) = \overline{(g, a)}$  if  $g \in G \downarrow$ . Then  $\varphi$  is an  $hl$ -isomorphism of the  $hl$ -group  $G$  into  $H$ .

If we put  $\psi(b) = \overline{(0, b)}$  for each  $b \in B \uparrow$  and  $\psi(b) = \overline{(a, b)}$  for each  $b \in B \downarrow$ , then  $\psi$  is an  $hl$ -isomorphism of the  $hl$ -group  $B$  into  $H$ .

If  $x \in G \cap B$  then  $\varphi(x) = \psi(x)$ . In fact, if  $x \in (G \cap B) \uparrow = (G \uparrow) \cap (B \uparrow)$ , then  $\varphi(x) = \overline{(x, 0)} = \overline{(0, x)} = \psi(x)$ ; and if  $x \in (G \cap B) \downarrow = (G \downarrow) \cap (B \downarrow)$ , then  $\varphi(x) = \overline{(x, a)} = \overline{(a, x)} = \psi(x)$ .

In the next, we shall identify elements  $g$  and  $\varphi(g)$  for each  $g \in G$  and also  $b$  and  $\psi(b)$  for each  $b \in B$ . Then  $G$  and  $B$  are  $hl$ -subgroups of  $H$ .

**Lemma 2.10.**  $H = \langle B \rangle_h$ .

**Proof.**  $B$  is an  $hl$ -subgroup of  $H$ . We have to prove that  $H \uparrow = \langle B \uparrow \rangle$ . Assume that  $\overline{(g, b)} \in H \uparrow$ ,  $\overline{(0, b')} \in B \uparrow$ ,  $\overline{(0, 0)} \leq \overline{(g, b)} \leq \overline{(0, b')}$ . Then  $g \in A \uparrow \subseteq B \uparrow$  and so  $g + b \in B \uparrow$ ,  $\overline{(g, b)} = \overline{(0, g + b)} \in B \uparrow$ . Hence  $B \uparrow$  is a convex  $l$ -subgroup of  $H \uparrow$ . Let  $\overline{(0, 0)} < \overline{(g, b)} \in H \uparrow$ ,  $\overline{(g, b)} \notin B \uparrow$ . Then  $g \notin A \uparrow$ . Therefore,  $g > 0$  and thus  $\overline{(0, b')} < \overline{(g, b)}$  for each  $\overline{(0, b')} \in B \uparrow$ . ■

By using Lemmas 2.10 and 2.1,  $B$  is an  $hl$ -ideal of  $H$ . Therefore, we can form the factor  $hl$ -group  $\overline{H} = H/B \uparrow$ .

**Lemma 2.11.** *Half  $l$ -groups  $\overline{G}$  and  $\overline{H}$  are  $hl$ -isomorphic.*

**Proof.** Define the mapping  $f : \overline{G} \rightarrow \overline{H}$  by  $f(g + A \uparrow) = g + B \uparrow$ . Let  $g + A \uparrow = g' + A \uparrow$ . Then  $g - g' \in A \uparrow \subset B \uparrow$ . Thus  $g + B \uparrow = g' + B \uparrow$ . Therefore the mapping  $f$  is correctly defined.

Let  $g_1 + A \uparrow, g_2 + A \uparrow \in \overline{G}$ . Then  $f((g_1 + A \uparrow) + (g_2 + A \uparrow)) = f((g_1 + g_2) + A \uparrow) = (g_1 + g_2) + B \uparrow = (g_1 + B \uparrow) + (g_2 + B \uparrow) = f(g_1 + A \uparrow) + f(g_2 + A \uparrow)$ .

Assume that  $g_1, g_2 \in G$  and  $f(g_1 + A \uparrow) = f(g_2 + A \uparrow)$ . From  $g_1 + B \uparrow = g_2 + B \uparrow$ , we infer that either  $g_1, g_2 \in G \uparrow$  or  $g_1, g_2 \in G \downarrow$ . Hence  $g_1 - g_2 \in G \uparrow \cap B \uparrow = A \uparrow$  and so  $g_1 + A \uparrow = g_2 + A \uparrow$ .

Let  $(\overline{g, b}) + B \uparrow \in \overline{H}$ . Assume that  $(\overline{g, b}) \in H \downarrow$ . Hence  $g \in G \downarrow$ . Recall that  $g$  is identified with  $(\overline{g, a})$ . As for  $(\overline{g, a}) - (\overline{g, b}) = (\overline{0, a - b}) \in B \uparrow$ , we have  $(\overline{g, a}) + B \uparrow = (\overline{g, b}) + B \uparrow$ . Therefore  $f(g + A \uparrow) = (\overline{g, b}) + B \uparrow$ . If  $(\overline{g, b}) \in H \uparrow$ , the proof is similar.

We have shown that  $f$  is a group isomorphism of  $\overline{G}$  onto  $\overline{H}$ .

Assume that  $g_1 + A \uparrow, g_2 + A \uparrow \in \overline{G}$  and  $g_1 + A \uparrow \leq g_2 + A \uparrow$ . If  $g_1 + A \uparrow = g_2 + A \uparrow$ , then  $f(g_1 + A \uparrow) = f(g_2 + A \uparrow)$ . Let  $g_1 + A \uparrow < g_2 + A \uparrow$ . By Lemma 2.2,  $g_1 < g_2$ . Hence  $f(g_1 + A \uparrow) = g_1 + B \uparrow < g_2 + B \uparrow = f(g_2 + A \uparrow)$ . The converse is similar. ■

Summarizing the previous results, we have

**Theorem 2.12.** *Let  $A, B$  and  $G$  be hl-groups from  $\mathcal{HL}$  satisfying (I)–(IV). Then there exists an hl-group  $H \in \mathcal{HL}$  such that*

- (i)  $H = \langle B \rangle_h$ ,
- (ii)  $G$  is an hl-subgroup of  $H$ ,
- (iii) hl-groups  $G/A \uparrow$  and  $H/B \uparrow$  are hl-isomorphic.

### 3. CUT COMPLETION OF A LEXICO EXTENSION

Let  $G$  be an hl-group. A subset  $X$  of  $G \uparrow$  is said to be a *cut of  $G \uparrow$*  if  $X$  is an order closed (i.e.,  $g = \bigvee S$  for  $S \subseteq X$  implies  $g \in X$ ) lattice ideal of  $G$  such that  $g + X \neq X \neq X + g$  for any  $0 < g \in G$ . A *cut of  $G \downarrow$*  is defined in the same way. If  $X$  is a cut either of  $G \uparrow$  or of  $G \downarrow$ , then  $X$  is called a cut of  $G$ .

$G \uparrow (G \downarrow)$  is said to be *cut complete* if every cut of  $G \uparrow (G \downarrow)$  has a supremum in  $G \uparrow (G \downarrow)$ .

Remark that if  $Z \subseteq G \uparrow (Z \subseteq G \downarrow)$ , then  $\sup(Z)$  exists in  $G \uparrow (G \downarrow)$  if and only if  $\sup(Z)$  exists in  $G$ , and  $\sup(Z)$  in  $G$  is equal to  $\sup(Z)$  in  $G \uparrow (G \downarrow)$ .

$G$  is called *cut complete* provided that every cut of  $G$  has a supremum in  $G$ .

An *hl*-subgroup  $G'$  of  $G$  will be said to be order dense in  $G$  if for every element  $0 < g \in G$  there exists  $g' \in G'$  with  $0 < g' \leq g$ .

An *hl*-group  $G^C$  is said to be a *cut completion* of  $G$  if the following conditions are satisfied:

- (i)  $G^C$  is cut complete;
- (ii)  $G$  is an order dense *hl*-subgroup of  $G^C$ ;
- (iii) if  $K$  is an *hl*-subgroup of  $G^C$  such that  $G \leq K < G^C$ , then  $K$  is not cut complete.

Since  $G \uparrow$  and  $G \downarrow$  are dually isomorphic lattices, we have

**Lemma 3.1.** *For any *hl*-group  $G$ ,  $G \in \mathcal{HL}$  is cut complete if and only if  $G \uparrow$  is cut complete. ■*

**Lemma 3.2** ([5], Lemma 3.1). *Let  $G$  be an abelian *l*-group,  $G = \langle A \rangle$ ,  $A \neq \{0\}$ . If  $A$  is cut complete, then  $G$  is cut complete. ■*

**Lemma 3.3.** *Let  $H$  and  $B$  be *hl*-groups from  $\mathcal{HL}$  such that  $H \uparrow$  is abelian and  $H = \langle B \rangle_h$ . If  $B$  is cut complete, then  $H$  is cut complete.*

**Proof.** The assumption that  $B$  is cut complete and Lemma 3.1 yield that  $B \uparrow$  is cut complete. From  $H \uparrow = \langle B \uparrow \rangle$ ,  $B \uparrow \neq \{0\}$  and Lemma 3.2, we obtain that  $H \uparrow$  is cut complete. Hence  $H$  is cut complete. ■

**Lemma 3.4.** *Let  $A, B, G$  satisfy the conditions (I)–(IV) and  $H$  be such as in Theorem 2.12. Suppose that  $B = A^C$ . Then  $H = G^C$ .*

**Proof.** In view of Lemma 2.10, we have  $H = \langle B \rangle_h$ .  $A$  is order dense in  $B$  and  $B$  is order dense in  $H$ . This yields that  $A$  is order dense in  $H$ . Thus  $G$  is order dense in  $H$ . From Lemma 3.3, it follows that  $H$  is cut complete. Assume that  $K$  is an *hl*-subgroup of  $H$  such that  $G \leq K < H$ . Then  $K \uparrow$  is an *l*-subgroup of  $H \uparrow$  with  $G \uparrow \leq K \uparrow < H \uparrow$ . In the same way as in the proof of Lemma 3.2 in [5], it can be shown that  $K \uparrow$  fails to be cut complete. Then, by Lemma 3.1,  $K$  is not cut complete. Therefore  $H = G^C$ . ■

**Theorem 3.5.** *Let  $G = \langle A \rangle_h$ , and  $A < G$ . Then*

- (i)  $G^C = \langle A^C \rangle_h$ ,
- (ii) *hl*-groups  $G/A \uparrow$  and  $G^C/A^C \uparrow$  are *hl*-isomorphic.

**Proof.** (i): Put  $B = A^C$ . Let  $H$  be as in Theorem 2.12. Then  $H = \langle B \rangle_h$  holds. Applying Lemma 3.4, we get that  $G^C = \langle A^C \rangle_h$  is valid.

(ii): Immediately follows from Theorem 2.12. ■

#### REFERENCES

- [1] R.N. Ball, *The structure of the  $\alpha$ -completion of a lattice ordered group*, Houston J. Math. **15** (1989), 481–515.
- [2] P. Conrad, *Lattice Ordered Groups*, Tulane University, New Orleans, LA, 1970.
- [3] M.R. Darnel, *Theory of Lattice Ordered Groups*, M. Dekker, Inc., New York 1995.
- [4] M. Giraudet and F. Lucas, *Groupes à moitié ordonnés*, Fund. Math. **139** (1991), 75–89.
- [5] J. Jakubík, *On cut completions of abelian lattice ordered groups*, Czechoslovak Math. J. **50** (125) (2000), 587–602.

Received 27 May 2002  
Revised 27 November 2002