

## ON ABSOLUTE RETRACTS AND ABSOLUTE CONVEX RETRACTS IN SOME CLASSES OF $\ell$ -GROUPS

JÁN JAKUBÍK

*Matematický ústav SAV,*  
*Grešákova 6, 040 01 Košice, Slovakia*  
**email:** kstefan@saske.sk

### Abstract

By dealing with absolute retracts of  $\ell$ -groups we use a definition analogous to that applied by Halmos for the case of Boolean algebras. The main results of the present paper concern absolute convex retracts in the class of all archimedean  $\ell$ -groups and in the class of all complete  $\ell$ -groups.

**Keywords:**  $\ell$ -group, absolute retract, absolute convex retract, archimedean  $\ell$ -group, complete  $\ell$ -group, orthogonal completeness.

**2000 Mathematics Subject Classification:** 06F20, 06F15.

### 1. INTRODUCTION

Retracts of abelian  $\ell$ -groups and of abelian cyclically ordered groups were investigated in [6], [7], [8].

Suppose that  $\mathcal{C}$  is a class of algebras. An algebra  $A \in \mathcal{C}$  is called an absolute retract in  $\mathcal{C}$  if, whenever  $B \in \mathcal{C}$  and  $A$  is a subalgebra of  $B$ , then  $A$  is a retract of  $B$  (i.e., there is a homomorphism  $h$  of  $B$  onto  $A$  such that  $h(a) = a$  for each  $a \in A$ ). Cf., e.g., Halmos [3].

Further, let  $\mathcal{C}$  be a class of  $\ell$ -groups. An element  $A \in \mathcal{C}$  will be called an absolute convex retract in  $\mathcal{C}$  if, whenever  $B \in \mathcal{C}$  and  $A$  is a convex  $\ell$ -subgroup of  $B$ , then  $A$  is a retract of  $B$ .

Let  $\mathcal{G}$  and Arch be the class of all  $\ell$ -groups, or the class of all archimedean  $\ell$ -groups, respectively.

It is easy to verify (cf. Section 2 below) that for  $A \in \mathcal{G}$  the following conditions are equivalent:

- (i)  $A$  is an absolute retract in  $\mathcal{G}$ ;
- (ii)  $A$  is an absolute convex retract in  $\mathcal{G}$ ;
- (iii)  $A = \{0\}$ .

In this note we prove

- ( $\alpha$ ) Let  $A$  be an absolute retract in the class Arch. Then the  $\ell$ -group  $A$  is divisible, complete and orthogonally complete.

By applying a result of [5] we obtain

- ( $\beta$ ) Let  $A \in \text{Arch}$  and suppose that the  $\ell$ -group  $A$  is complete and orthogonally complete. Then  $A$  is an absolute convex retract in the class Arch.

The question whether the implication in ( $\alpha$ ) (or in ( $\beta$ ), respectively) can be reversed remains open.

Let us denote by

Compl - the class of all complete  $\ell$ -groups;

Compl\* - the class of all  $\ell$ -groups which are complete and orthogonally complete.

- ( $\gamma$ ) Let  $A \in \text{Compl}$ . Then the following conditions are equivalent:
  - (i)  $A$  is orthogonally complete.
  - (ii)  $A$  is an absolute convex retract in the class Compl.

As a corollary we obtain that each  $\ell$ -group belonging to Compl\* is an absolute convex retract in the class Compl\*.

We prove that if the class  $\mathcal{C} \subseteq \mathcal{G}$  is closed with respect to direct products and if  $A_i$  ( $i \in I$ ) are absolute (convex) retracts in  $\mathcal{C}$ , then their direct product  $\prod_{i \in I} A_i$  is also an absolute (convex) retract in  $\mathcal{C}$ .

## 2. PRELIMINARIES

For  $\ell$ -groups we apply the notation as in Conrad [1]. Hence, in particular, the group operation in an  $\ell$ -group is written additively.

We recall some relevant notions. Let  $G$  be an  $\ell$ -group.  $G$  is *divisible* if for each  $a \in G$  and each positive integer  $n$  there is  $x \in G$  with  $nx = a$ . A system  $\emptyset \neq \{x_i\}_{i \in I} \subseteq G^+$  is called *orthogonal* (or *disjoint*) if  $x_{i(1)} \wedge x_{i(2)} = 0$  whenever  $i(1)$  and  $i(2)$  are distinct elements of  $I$ . If each orthogonal subset of  $G$  possesses the supremum in  $G$  then  $G$  is said to be *orthogonally complete*.  $G$  is *complete* if each nonempty bounded subset of  $G$  has the supremum and the infimum in  $G$ .

$G$  is *archimedean* if, whenever  $0 < x \in G$  and  $y \in G$ , then there is a positive integer  $n$  such that  $nx \not\leq y$ . For each archimedean  $\ell$ -group  $G$  there exists a complete  $\ell$ -group  $D(G)$  (the Dedekind completion of  $G$ ) such that

- (i)  $G$  is a closed  $\ell$ -subgroup of  $D(G)$ ;
- (ii) for each  $x \in D(G)$  there are subsets  $\{y_i\}_{i \in I}$  and  $\{z_j\}_{j \in J}$  of  $G$  such that the relations

$$\sup\{x_i\}_{i \in I} = x = \inf\{z_j\}_{j \in J}$$

are valid in  $D(G)$ .

Let  $G_1$  be a linearly ordered group and let  $G_2$  be an  $\ell$ -group. The symbol  $G_1 \circ G_2$  denotes the lexicographic product of  $G_1$  and  $G_2$ . The elements of  $G_1 \circ G_2$  are pairs  $(g_1, g_2)$  with  $g_1 \in G_1$  and  $g_2 \in G_2$ . For each  $g_2 \in G_2$ , the pair  $(0, g_2)$  will be identified with the element  $g_2$  of  $G_2$ . Then  $G_2$  is a convex  $\ell$ -subgroup of  $G_1 \circ G_2$ .

**Lemma 2.1.** *Let  $A$  be an  $\ell$ -group,  $A \neq \{0\}$ , and let  $G_1$  be a linearly ordered group,  $G_1 \neq \{0\}$ . Put  $B = G_1 \circ A$ . Then  $A$  fails to be a retract of  $B$ .*

**Proof.** By way of contradiction, suppose that  $A$  is a retract of  $B$ . Let  $h$  be the corresponding retract homomorphism of  $B$  onto  $A$ ; i.e.,  $h(a) = a$  for each  $a \in A$ . There exists  $g_1 \in G_1$  with  $g_1 > 0$ . Denote  $(g_1, 0) = b$ ,  $h(b) = a$ . Further, there exists  $a_1 \in A$  with  $a_1 > a$ . We have  $a_1 < b$ , whence  $h(a_1) \leq h(b)$ , thus  $a_1 \leq a$ , which is a contradiction. ■

Let us denote by  $\mathcal{A}$  the class of all abelian lattice ordered groups. If  $A, G_1$  and  $B$  are as in Lemma 2.1 and  $A, G_1 \in \mathcal{A}$ , then also  $B$  belongs to  $\mathcal{A}$ . Thus Lemma 2.1 yields

**Proposition 2.2.** *Let  $\mathcal{C} \in \{\mathcal{G}, \mathcal{A}\}$  and let  $A$  be an absolute retract (or an absolute convex retract, respectively) in the class  $\mathcal{C}$ . Then  $A = \{0\}$ . ■*

It is obvious that  $\{0\}$  is an absolute (convex) retract in both the classes  $\mathcal{G}$  and  $\mathcal{A}$ .

Let us remark that if  $G_1, B \in \mathcal{G}$  and if  $G_1$  is a retract of  $B$ , then  $G_1$  need not be a convex  $\ell$ -subgroup of  $B$ . This is verified by the following example:

Let  $G_1$  be a linearly ordered group,  $G_1 \neq \{0\}$ . Further, let  $G_2 \in \mathcal{G}$ ,  $G_2 \neq \{0\}$ . Put  $B = G_1 \circ G_2$ . If  $g_1 \in G_1$ , then the element  $(g_1, 0)$  of  $B$  will be identified with the element  $g_1$  of  $G_1$ . Thus  $G_1$  turns out to be an  $\ell$ -subgroup of  $B$  which is not a convex subset of  $B$ . For each  $(g_1, g_2) \in B$  we put  $h((g_1, g_2)) = g_1$ . Then  $h$  is a homomorphism of  $B$  onto  $G_1$  such that  $h(g_1) = g_1$  for each  $g_1 \in G_1$ . Hence  $G_1$  is a retract of  $B$ .

### 3. PROOFS OF $(\alpha)$ , $(\beta)$ AND $(\gamma)$

In this section we assume that  $A$  is an archimedean  $\ell$ -group. Hence  $A$  is abelian.

It is well-known that there exists the divisible hull  $A^d$  of  $A$ . Thus

- (i)  $A^d$  is a divisible  $\ell$ -group;
- (ii)  $A$  is an  $\ell$ -subgroup of  $A^d$ ;
- (iii) if  $g \in A^d$ , then there are  $a \in A$ , a positive integer  $n$  and an integer  $m$  such that  $ng = ma$ .

**Lemma 3.1.** *Assume that  $A$  is an absolute retract in the class Arch. Then the  $\ell$ -group  $A$  is divisible.*

**Proof.** By way of contradiction, suppose that  $A$  fails to be divisible. Thus there are  $a_1 \in A$  and  $n \in \mathbb{N}$  such that there is no  $x$  in  $A$  with  $nx = a_1$ .

Put  $B = A^d$ . In view of the assumption,  $A$  is a retract of  $B$ ; let  $h$  be the corresponding retract homomorphism.

There exists  $b \in B$  with  $nb = a_1$ . Then  $b \notin A$ . Denote  $h(b) = a$ . We have

$$a_1 = h(a_1) = h(nb) = nh(b) = na,$$

which is a contradiction. ■

**Lemma 3.2.** *Assume that  $A$  is an absolute retract in the class Arch. Then  $A$  is a complete  $\ell$ -group.*

**Proof.** By way of contradiction, suppose that  $A$  fails to be complete. Put  $B = D(A)$ . Then  $A$  is an  $\ell$ -subgroup of  $B$  and  $A \neq B$ . Thus there is  $b \in B$  such that  $b$  does not belong to  $A$ .

In view of the assumption,  $A$  is a retract of  $B$ ; let  $h$  be the corresponding retract homomorphism. Put  $h(b) = a$ .

There exists a subset  $\{a_i\}_{i \in I}$  of  $A$  such that the relation

$$b = \bigvee_{i \in I} a_i$$

is valid in  $B$ . Hence  $a_i \leq b$  for each  $i \in I$ . This yields

$$a_i = h(a_i) \leq h(b) = a$$

for each  $i \in I$ . Thus  $b \leq a$ .

At the same time, there exists a subset  $\{a'_j\}_{j \in J}$  of  $A$  such that the relation

$$b = \bigwedge_{j \in J} a'_j$$

holds in  $B$ . Hence  $b \leq a'_j$  for each  $j \in J$ , thus by applying the homomorphism  $h$  we obtain that  $a \leq a'_j$  for each  $j \in J$ . Therefore  $a \leq b$ . Summarizing,  $a = b$  and we arrived at a contradiction. ■

**Lemma 3.3.** *Suppose that  $H$  is a complete  $\ell$ -group. Then there exists an  $\ell$ -group  $K$  such that*

- (i)  $H$  is a convex  $\ell$ -subgroup of  $K$ ;
- (ii)  $K$  is complete and orthogonally complete;
- (iii) for each  $0 < k \in K$  there exists a disjoint subset  $\{x_i\}_{i \in I}$  of  $H$  such that the relation

$$k = \bigvee_{i \in I} x_i$$

is valid in  $K$ .

**Proof.** This is a consequence of results of [5]. ■

**Lemma 3.4.** *Assume that  $A$  is an absolute retract in the class Arch. Then the  $\ell$ -group  $A$  is orthogonally complete.*

**Proof.** In view of Lemma 3.2,  $A$  is complete. Put  $A = H$  and let  $K$  be as in Lemma 3.3. According to the assumption,  $A$  is a retract of  $K$ . Let  $h$  be the corresponding retract homomorphism.

Let  $0 < k \in K$  and let  $\{x_i\}_{i \in I}$  be as in Lemma 3.3. Put  $h(k) = a$ . Then  $a \geq h(x_i) = x_i$  for each  $i \in I$ , whence  $k \leq a$ . Thus the condition (i) of Lemma 3.3 yields that  $k \in A$ . Hence  $K^+ \subseteq A$  and then  $K \subseteq A$ . Therefore  $K = A$  and so  $A$  is orthogonally complete. ■

From Lemmas 3.1, 3.2 and 3.4 we conclude that  $(\alpha)$  is valid.

Let  $G_1, G_2 \in \mathcal{G}$ ; their direct product is denoted by  $G_1 \times G_2$ . If  $g_1 \in G_1$ , then the element  $(g_1, 0)$  of  $G_1 \times G_2$  will be identified with  $g_1$ . Similarly, for  $g_2 \in G_2$ , the element  $(0, g_2)$  of  $G_1 \times G_2$  will be identified with  $g_2$ . Under this identification, both  $G_1$  and  $G_2$  are convex  $\ell$ -subgroups of  $G_1 \times G_2$ .

**Definition 3.5.** (Cf. [2].) Let  $G_1 \in \text{Arch}$ . We say that  $G_1$  has the *splitting property* if, whenever  $H \in \text{Arch}$  and  $G_1$  is a convex  $\ell$ -subgroup of  $H$ , then  $G_1$  is a direct factor of  $H$ .

**Proposition 3.6.** (Cf. [4].) Let  $G_1 \in \text{Arch}$ . Then the following conditions are equivalent:

- (i)  $G_1$  has the splitting property.
- (ii) The  $\ell$ -group  $G_1$  is complete and orthogonally complete.

**Lemma 3.7.** *Let  $H \in \mathcal{G}$  and let  $G_1$  be a direct factor of  $H$ . Then  $G_1$  is a retract of  $H$ .*

**Proof.** There exists  $G_2 \in \mathcal{G}$  such that  $H = G_1 \times G_2$ . For  $(g_1, g_2) \in H$  we put  $h((g_1, g_2)) = g_1$ . Then  $h$  is a retract homomorphism of  $H$  onto  $G_1$ . ■

**Proof of  $(\beta)$ .** Let  $A, B \in \text{Arch}$  and suppose that  $A$  is a convex  $\ell$ -subgroup of  $B$ . Further, suppose that  $A$  is complete and orthogonally complete. In view of Proposition 3.6,  $A$  is a direct factor of  $B$ . Hence according to Lemma 3.7,  $A$  is a retract of  $B$ . Therefore  $A$  is an absolute convex retract in the class Arch. ■

**Lemma 3.8.** *Let  $A \in \text{Compl}$ . Suppose that  $A$  is an absolute convex retract in the class  $\text{Compl}$ . Then  $A$  is orthogonally complete.*

**Proof.** Put  $H = A$  and let  $K$  be as in Lemma 3.3. In view of Lemma 3.3 (i),  $A$  is a convex  $\ell$ -subgroup of  $K$ . Hence according to the assumption,  $A$  is a retract of  $K$ . Now it suffices to apply the same method as in the proof of Lemma 3.4. ■

**Lemma 3.9.** *Let  $A \in \text{Compl}$ . Suppose that  $A$  is orthogonally complete. Then  $A$  is an absolute convex retract in the class  $\text{Compl}$ .*

**Proof.** In view of  $(\beta)$ ,  $A$  is an absolute convex retract in the class  $\text{Arch}$ . It is well-known that the class  $\text{Compl}$  is a subclass of  $\text{Arch}$ . Hence  $A$  is an absolute convex retract in the class  $\text{Compl}$ . ■

From Lemmas 3.8 and 3.9 we conclude that  $(\gamma)$  holds.

**Corollary 3.10.** *Let  $A \in \text{Compl}^*$ . Then  $A$  is an absolute convex retract in the class  $\text{Compl}^*$ .* ■

#### 4. DIRECT PRODUCTS

Let  $A_i$  ( $i \in I$ ) be  $\ell$ -groups; consider their direct product

$$(1) \quad A = \prod_{i \in I} A_i.$$

Without loss of generality we can suppose that  $A_{i(1)} \cap A_{i(2)} = \{0\}$  whenever  $i(1)$  and  $i(2)$  are distinct elements of  $I$ . For  $a \in A$  and  $i \in I$ , we denote by  $a_i$  or by  $a(A_i)$  the component of  $a$  in the direct factor  $A_i$ .

Let  $i \in I$ . Put

$$A'_i = \{a \in A : a_i = 0\}.$$

Then we have

$$(2) \quad A = A_i \times A'_i,$$

$$A'_i = \prod_{j \in I \setminus \{i\}} A_j.$$

Let  $i(0) \in I$  and  $a^{i(0)} \in A_{i(0)}$ . There exists  $a \in A$  such that

$$a_i = \begin{cases} a^{i(0)} & \text{if } i = i(0), \\ 0 & \text{otherwise.} \end{cases}$$

Then the element  $a$  of  $A$  will be identified with the element  $a^{i(0)}$  of  $A_{i(0)}$ . Under this identification, each  $A_i$  turns out to be a convex  $\ell$ -subgroup of  $A$ .

**Lemma 4.1.** *Let  $B$  be an  $\ell$ -group and let  $A$  be an  $\ell$ -subgroup of  $B$ . Suppose that (1) is valid. Let  $i$  be a fixed element of  $I$  and assume that  $A_i$  is a retract of  $B$ ; the corresponding retract homomorphism will be denoted by  $h_i$ . Then for each  $a \in A$  the relation*

$$h_i(a) = a_i$$

*is valid.*

**Proof.** a) At first let  $0 \leq a' \in A'_i$  and  $0 \leq a^i \in A_i$ . Then  $a' \wedge a^i = 0$ , thus

$$0 = h_i(a') \wedge h_i(a^i) = h_i(a') \wedge a^i.$$

Since this is valid for each  $a^i \in A_i$  and  $h_i(a') \in A_i$  we conclude that  $h_i(a') = 0$ . Then  $h_i(-a') = 0$  as well and this yields that  $h_i(a'') = 0$  for each  $a'' \in A'_i$ .

b) Let  $a \in A$ . In view of (2) we have

$$a = a_i + a(A'_i).$$

Thus

$$h_i(a) = h_i(a_i) + h_i(a(A'_i)).$$

According to a),  $h_i(a(A'_i)) = 0$ . Thus  $h_i(a) = a_i$ . ■

**Lemma 4.2.** *Let  $B$  be an  $\ell$ -group and let  $A$  be an  $\ell$ -subgroup of  $B$ . Suppose that (1) is valid and that for each  $i \in I$ ,  $A_i$  is a retract of  $B$ ; the corresponding retract homomorphism will be denoted by  $h_i$ . For  $b \in B$  we put*

$$h(b) = b^1 \in A,$$

*where  $b_i^1 = h_i(b)$  for each  $i \in I$ . Then*



- (i)  $h$  is a homomorphism of  $B$  into  $A$ ;
- (ii)  $h(a) = a$  for each  $a \in A$ .

**Proof.** The definition of  $h$  and the relation (1) immediately yield that (i) is valid. Let  $a \in A$  and  $i \in I$ . Put  $h(a) = a^1$ . We have

$$a = a_i + a(A'_i),$$

thus by applying (i),

$$h(a) = h(a_i) + h(a(A'_i)),$$

$$a_i^1 = h_i(a_i) + h_i(a(A'_i)).$$

Since  $h_i(a_i) = a_i$  and because  $(a(A'_i))_i = 0$ , according to Lemma 4.1, we obtain

$$a_i^1 = a_i \quad \text{for each } i \in I,$$

thus  $a^1 = a$ . ■

**Corollary 4.3.** *Let the assumptions of Lemma 4.2 be valid. Then  $A$  is a retract of  $B$ .* ■

From Corollary 4.3 we immediately conclude

**Proposition 4.4.** *Assume that  $\mathcal{C}$  is a class of  $\ell$ -groups which is closed with respect to direct products. Let  $A_i$  ( $i \in I$ ) be absolute retracts in  $\mathcal{C}$  and let (1) be valid. Then  $A$  is an absolute retract in  $\mathcal{C}$ .* ■

**Proposition 4.5.** *Assume that  $\mathcal{C}$  is a class of  $\ell$ -groups which is closed with respect to direct products. Let  $A_i$  ( $i \in I$ ) be absolute convex retracts in  $\mathcal{C}$  and let (1) be valid. Then  $A$  is an absolute convex retract in  $\mathcal{C}$ .*

**Proof.** Let  $B \in \mathcal{C}$  and suppose that  $A$  is a convex  $\ell$ -subgroup of  $B$ . Then all  $A_i$  are convex  $\ell$ -subgroups of  $B$ . Hence in view of the assumption, all  $A_i$  are retracts of  $B$ . Thus according to Corollary 4.3,  $A$  is a retract of  $B$ . Therefore  $A$  is an absolute convex retract in the class  $\mathcal{C}$ . ■

## 5. AN EXAMPLE

The assertions of the following two lemmas are easy to verify; the proofs will be omitted.

**Lemma 5.1.** *Let  $A$  be an  $\ell$ -group which is complete and divisible. Then*

- (i) *we can define (in a unique way) a multiplication of elements of  $A$  with reals such that  $A$  turns out to be a vector lattice;*
- (ii) *if  $r > 0$  is a real,  $0 < a \in A$ ,  $X = \{q_1 \in Q : 0 < q_1 \leq r\}$ ,  $Y = \{q_2 \in R : r \leq q_2\}$ , then the relations*

$$\sup(q_1 a) = ra = \inf(q_2 a)$$

*are valid in  $A$ ;*

- (iii) *if  $A_1$  is an  $\ell$ -subgroup of  $A$  such that  $A_1$  is complete and divisible, and  $a_1 \in A$ , then for each real  $r$  the multiplication  $ra_1$  in  $A_1$  gives the same result as the multiplication  $ra_1$  in  $A$ . ■*

**Lemma 5.2.** *Let  $A$  be as in Lemma 5.1 and suppose that  $A = \prod_{i \in I} A_i$ . Then all  $A_i$  are complete and divisible; moreover, for each real  $r$ , each  $a \in A$  and each  $i \in I$  we have*

$$(ra)_i = ra_i. \quad \blacksquare$$

Let  $R$  be the additive group of all reals with the natural linear order. We denote by  $\mathcal{C}_{\mathcal{R}}$  the class of all lattice ordered groups which can be expressed as direct products of  $\ell$ -groups isomorphic to  $R$ .

We remark that if  $B \in \mathcal{C}_{\mathcal{R}}$  and if  $A$  is an  $\ell$ -subgroup of  $B$  which is isomorphic to  $R$ , then  $A$  need not be a convex  $\ell$ -subgroup of  $B$ . In fact, suppose that

$$B = \prod_{i \in I} B_i,$$

where each  $B_i$  is isomorphic to  $R$ ; let  $\varphi_i$  be an isomorphism of  $R$  onto  $B_i$ . For each  $r \in R$  put

$$\varphi(r) = (\dots, \varphi_i(r), \dots)_{i \in I},$$

$$A = \varphi(R).$$

$A$  is an  $\ell$ -subgroup of  $B$ ; if  $I$  has more than one element, then  $A$  fails to be convex in  $B$ .

Let  $B$  be as above; suppose that  $A$  is an  $\ell$ -group isomorphic to  $R$  and that  $A$  is an  $\ell$ -subgroup of  $B$ . Let  $0 < a \in A$ . Then  $a_i = a(B_i) \geq 0$  for each  $i \in I$  and there exists  $i(0) \in I$  with  $a_{i(0)} > 0$ . Thus, in view of Lemma 5.1, we have  $(ra)_{i(0)} > 0$  for each  $r \in R$  with  $r \neq 0$ . Further, for each  $a_1 \in A$  there exists a uniquely determined element  $r \in R$  with  $a_1 = ra$ . This yields that the mapping

$$\varphi_{i(0)} : a_1 \mapsto (a_1)_{i(0)}$$

is an isomorphism of  $A$  into  $B_{i(0)}$ .

Let  $b \in B_{i(0)}$ . There exists a unique  $r \in R$  such that

$$b = ra_{i(0)}.$$

Then, in view of Lemma 5.2,  $b = (ra)_{i(0)}$  and hence the mapping  $\varphi_{i(0)}$  is an isomorphism of  $A$  onto  $B_{i(0)}$ .

For each  $b \in B$  we put

$$h(b) = \varphi_{i(0)}^{-1}(b_{i(0)}).$$

Then  $h$  is a homomorphism of  $B$  into  $A$ . For  $a_1 \in A$  the definition of  $\varphi_{i(0)}$  yields

$$h(a_1) = a_1.$$

Thus we obtain

**Lemma 5.3.** *Let  $B \in \mathcal{C}_{\mathcal{R}}$  and let  $A$  be an  $\ell$ -subgroup of  $B$  such that  $A$  is isomorphic to  $R$ . Then  $A$  is a retract of  $B$ . ■*

**Corollary 5.4.** *Let  $A$  be an  $\ell$ -group isomorphic to  $R$ . Then  $A$  is an absolute retract in the class  $\mathcal{C}_{\mathcal{R}}$ . ■*

From Lemma 5.4 and Corollary 4.5 we conclude

**Proposition 5.5.** *Each element of  $\mathcal{C}_{\mathcal{R}}$  is an absolute retract in the class  $\mathcal{C}_{\mathcal{R}}$ . ■*

## REFERENCES

- [1] P. Conrad, *Lattice Ordered Groups*, Tulane University, New Orleans, LA, 1970.
- [2] P. Conrad, *The essential closure of an archimedean lattice-ordered group*, *Duke Math. J.* **38** (1971), 151–160.
- [3] P. Halmos, *Injective and projective Boolean algebras*, p. 114–122 in: “*Lattice Theory*” (R.P. Dilworth, editor), Amer. Math. Soc., Providence, RI, 1961.
- [4] J. Jakubík, *Splitting property of lattice ordered groups*, *Czechoslovak Math. J.* **24** (1974), 257–269.
- [5] J. Jakubík, *Orthogonal hull of a strongly projectable lattice ordered group*, *Czechoslovak Math. J.* **28** (1978), 484–504.
- [6] J. Jakubík, *Retracts of abelian cyclically ordered groups*, *Archivum Math. (Brno)* **25** (1989), 13–18.
- [7] J. Jakubík, *Retracts of abelian lattice ordered groups*, *Czechoslovak Math. J.* **39** (1989), 477–489.
- [8] J. Jakubík, *Retract varieties of lattice ordered groups*, *Czechoslovak Math. J.* **40** (1990), 104–112.

Received 21 October 2002