

ON VARIETIES OF LEFT DISTRIBUTIVE LEFT IDEMPOTENT GROUPOIDS

DAVID STANOVSKÝ*

*Charles University in Prague,
KA MFF UK, Sokolovská 83,
18675 Praha, Czech Republic*

e-mail: stanovsk@karlin.mff.cuni.cz

Abstract

We describe a part of the lattice of subvarieties of left distributive left idempotent groupoids (i.e. those satisfying the identities $x(yz) \approx (xy)(xz)$ and $(xx)y \approx xy$) modulo the lattice of subvarieties of left distributive idempotent groupoids. A free groupoid in a subvariety of LDLI groupoids satisfying an identity $x^n \approx x$ decomposes as the direct product of its largest idempotent factor and a cycle. Some properties of subdirectly irreducible LDLI groupoids are found.

Keywords: left distributivity, left idempotence, right zero band, LDLI groupoids, subdirectly irreducible, free groupoid, lattice of subvarieties.

2000 Mathematics Subject Classification: 08B15, 08B20, 08B26, 20N02.

We consider groupoids (i.e. sets equipped with a binary operation) satisfying the following two identities:

$$(LD) \quad x(yz) \approx (xy)(xz),$$

$$(LI) \quad (xx)y \approx xy.$$

*While working on this paper the author was partially supported by the Grant Agency of the Czech Republic, grants #201/02/0594 and #201/02/0148, and by the Institutional grant MSM113200007.

We call such groupoids *left distributive left idempotent*, shortly LDLI. A groupoid is called *idempotent*, if it satisfies the identity

$$(I) \quad xx \approx x.$$

Note that the well known class of left distributive left quasigroups (see, e.g., [3], [8], and [9]) satisfies left idempotency. Indeed, our results can be applied there.

The purpose of this note is to continue recent investigations of P. Jedlička [4] on LDLI groupoids. We apply his result to compute the lattice of subvarieties of LDLI groupoids satisfying an identity $x^{n+1} \approx x$ for some n , modulo the lattice of subvarieties of LDI groupoids (see Theorem 4). This generalizes a result of T. Kepka [5] who described in a similar way subvarieties of LD groupoids with $x(xy) \approx y$ (such groupoids are called *left symmetric*; they satisfy LI and $x^3 \approx x$).

In Section 2 we show some properties of subdirectly irreducible LDLI groupoids and apply them to get some information about the structure of the lattice of subvarieties satisfying identities $x^{m+n} \approx x^m$.

We use rather standard terminology and notation, for an introduction to universal algebra, see, e.g., [2]. We need the following result of P. Jedlička [4]. Let G be an LDLI groupoid and let ip_G be the smallest equivalence on G containing $\{(a, aa) : a \in G\}$. Then ip_G is a congruence, G/ip_G is idempotent, and ip_G is the smallest congruence such that the corresponding factor is idempotent. Moreover, for any $(a, b) \in ip_G$, $ac = bc$ holds for every $c \in G$. Consequently, any block of ip_G is a subalgebra of G and it is term equivalent to a connected monounary algebra.

1. VARIETIES SATISFYING $x^{n+1} \approx x$

Let us define inductively $x^1 = x$ and $x^n = xx^{n-1}$ for every $n > 1$. (Note that other possible definitions of powers do not really make sense: one can check by induction that any term t in one variable is LI-equivalent to the term x^n , where n is the depth of the rightmost variable in t .) It is easy to prove that in LDLI groupoids the identities $xy \approx x^n y$ and $(x^k)^l \approx x^{k+l-1}$ hold.

We say that a variety \mathcal{V} of groupoids has *exponent* n , if n is the least positive integer such that the identity $x^{n+1} \approx x$ holds in \mathcal{V} . (Of course, such n does not necessarily exist, however, many important varieties, for

instance left n -symmetric left distributive groupoids (see, e.g., [9]), have finite exponent.)

Let C_n denote the groupoid on the set $\{0, \dots, n-1\}$ with the operation $ab = b + 1 \pmod n$. Clearly, C_n are LDLI groupoids.

Lemma 1. *Let G be an LDLI groupoid with $x^{n+1} \approx x$. Then*

- (1) *every block of ip_G is isomorphic to C_k for some $k|n$;*
- (2) *C_n is a homomorphic image of G , if and only if G is isomorphic to the direct product $C_n \times (G/ip_G)$.*

Proof.

(1): is easy.

(2): Choose a projection $g : G \rightarrow C_n$ and put $f(x) = (g(x), x/ip_G)$. Then $f : G \rightarrow C_n \times (G/ip_G)$ is a homomorphism. Since there is a homomorphism $C_k \rightarrow C_l$ iff $l|k$, every block of ip_G is isomorphic to C_n (g restricted to a block of ip_G is a homomorphism). Hence, g is bijective on every block of ip_G , because rotations are the only endomorphisms of C_n , and thus, f is an isomorphism. The other implication is clear. ■

Lemma 2. *Let \mathcal{V} be a subvariety of LDLI groupoids and assume \mathcal{V} has exponent n . Then $C_k \in \mathcal{V}$, iff $k|n$.*

Proof. If k does not divide n , then C_k does not satisfy $x^{n+1} \approx x$. On the other hand, if \mathcal{V} has exponent n , then, according to Lemma 1, ip -blocks of elements of \mathcal{V} are C_k with $k|n$ (indeed, ip -blocks are subgroupoids). Let k_0 be the greatest k such that $C_k \in \mathcal{V}$. If $k_0 < n$, then \mathcal{V} satisfies $x^{k_0+1} \approx x$, a contradiction with minimality of n . Hence, $C_n \in \mathcal{V}$ and thus, $C_k \in \mathcal{V}$ for all $k|n$, because they are homomorphic images of C_n . ■

Let $\mathbf{F}_{\mathcal{V}}(X)$ denote the free groupoid over X in a variety \mathcal{V} . Let \mathcal{I} denote the variety of idempotent groupoids.

Theorem 3. *Let \mathcal{V} be a subvariety of LDLI groupoids and assume \mathcal{V} has exponent n . Then $\mathbf{F}_{\mathcal{V}}(X)$ is isomorphic to $C_n \times \mathbf{F}_{\mathcal{V} \cap \mathcal{I}}(X)$. Consequently, the variety \mathcal{V} is generated by $(\mathcal{V} \cap \mathcal{I}) \cup \{C_n\}$.*

Proof. Since $C_n \in \mathcal{V}$, it is a homomorphic image of $\mathbf{F}_{\mathcal{V}}(X)$. Hence, by Lemma 1, $\mathbf{F}_{\mathcal{V}}(X) \simeq C_n \times H$, where $H = \mathbf{F}_{\mathcal{V}}(X)/ip_{\mathbf{F}_{\mathcal{V}}(X)}$. It is easy to see that $H \simeq \mathbf{F}_{\mathcal{V} \cap \mathcal{I}}(X)$, because ip is the smallest idempotent congruence. ■

A *right zero band* is a groupoid satisfying the identity $xy \approx y$. It is well known that the variety \mathcal{RZB} of right zero bands is minimal (i.e. it is generated by each of its elements).

Theorem 4. *Let L denote the lattice of subvarieties of LDI groupoids, K its sublattice of varieties containing \mathcal{RZB} and N the lattice of positive integer divisors of n . The lattice of subvarieties of the variety of LDLI groupoids satisfying $x^{n+1} \approx x$ is isomorphic to the lattice $(L \times \{1\}) \cup (K \times (N \setminus \{1\}))$ (regarded as a subposet of $L \times N$), sending a variety \mathcal{V} of exponent m to the pair $\Phi(\mathcal{V}) = (\mathcal{V} \cap \mathcal{I}, m)$.*

Proof. First, we check that the mapping Φ is well-defined: the exponent m of a subvariety \mathcal{V} is clearly a divisor of n and since \mathcal{V} contains C_m , it contains a right zero band $(C_m \times C_m)/ip$, where $ip = ip_{C_m \times C_m}$, and thus it contains the whole variety \mathcal{RZB} , because it is minimal. Next, Φ is injective: if \mathcal{V}_1 and \mathcal{V}_2 are distinct varieties of exponent m , then $\mathcal{V}_1 \cap \mathcal{I}$ and $\mathcal{V}_2 \cap \mathcal{I}$ are distinct, because \mathcal{V}_i is generated by $(\mathcal{V}_i \cap \mathcal{I}) \cup \{C_m\}$, $i = 1, 2$. The mapping Φ is onto, a pair (\mathcal{W}, m) is the image of the variety generated by $\mathcal{W} \cup \{C_m\}$. Indeed, let G be an idempotent groupoid in the variety generated by $\mathcal{W} \cup \{C_m\}$ and we show that $G \in \mathcal{W}$. The case $m = 1$ is trivial, so let $m > 1$. By Birkhoff's HSP Theorem (see [1]), there are $H \in \mathcal{W}$, $K \leq H \times C_m^k$ (for some k) and an onto homomorphism $\varphi : K \rightarrow G$. Since ip_K is the smallest idempotent congruence and G is idempotent, there is an onto homomorphism $\psi : K/ip_K \rightarrow G$. Further, $K/ip_K \leq (K \times C_m^k)/ip \simeq K \times (C_m^k/ip)$. However, C_m^k/ip is a right zero band and thus it is in \mathcal{W} . Consequently, G is a homomorphic image of a subgroupoid of a groupoid from \mathcal{W} , thus it is in \mathcal{W} . Finally, Φ clearly preserves the order and it follows from Theorem 3 that also Φ^{-1} preserves the order. Consequently, Φ is a lattice isomorphism. ■

Example 1. B. Roszkowska proved in [7] that the lattice of subvarieties of left symmetric medial idempotent (LSMI) groupoids (those where the identities $x(xy) \approx y$, $xy \cdot uv \approx xu \cdot yv$ and $xx \approx x$ hold) is isomorphic to the lattice of positive integers ordered by divisibility with a top element added. A number n corresponds to the variety based on $w_n(x, y) \approx y$ (relatively to LSMI), where

$$w_n(x, y) = \underbrace{x(y(x(y(\dots))))}_n.$$

Note that right zero bands satisfy $w_n(x, y) \approx y$ iff n is even. Thus, using Theorem 4, it is easy to describe bases of all proper subvarieties of left symmetric left distributive medial groupoids (relatively to LSLDM):

1. $xx \approx x$;
2. $w_n(x, y) \approx y$ and $xx \approx x$, for every n ;
3. $w_n(x, y) \approx y$, for every n even.

(Note that mediality and idempotency imply left distributivity, however, non-idempotent medial groupoids are not necessarily left distributive.)

Example 2. J. Płonka ([6]) investigated idempotent groupoids satisfying

$$\underbrace{x(x(\dots(xy)))}_n \approx y, \quad x(yz) \approx y(xz) \quad \text{and} \quad xz \approx (yx)z.$$

He called them *n-cyclic groupoids*. It is easy to see that they are LDLI and that 1-cyclic groupoids are precisely right zero bands. Płonka proved that the only non-trivial subvarieties of *n-cyclic groupoids* are *m-cyclic groupoids* for $m|n$. One can thus use Theorem 4 to describe the subvarieties of non-idempotent *n-cyclic groupoids*. Every non-trivial one is generated by idempotent *m-cyclic groupoids* and the groupoid C_k , for some divisors m, k of n ; hence there are exactly $q^2 + 1$ such subvarieties, where q is the number of divisors of n .

2. VARIETIES SATISFYING $x^{m+n} \approx x^m$

Theorems 3 and 4 cannot be generalized to varieties satisfying an identity $x^{m+n} \approx x^m$ for $m > 1$. For instance, in an LDLI groupoid G with $x^3 \approx x^2$, every ip_G -block is a *constant groupoid*, i.e. $ab = cd$ for all ip_G -congruent elements a, b, c, d (the corresponding unary algebra is a loop with several “tails” of length 1). However, this variety is not generated by LDI and constant groupoids. Indeed, both LDI and constant groupoids satisfy the identity $xy \approx x(yx)$, while the groupoid

	a	b	c	d
a, b	b	b	c	d
c, d	a	b	d	d

does not, though it is LDLI with $x^3 \approx x^2$. We thus present a weaker result for such varieties.

For a variety \mathcal{V} , let $\mathcal{V}_{m,n}$ denote the subvariety of \mathcal{V} based (relatively to \mathcal{V}) by the identity $x^{m+n} \approx x^m$.

It is well known that a groupoid G is subdirectly irreducible, iff it possesses the smallest non-trivial congruence, and that any variety is generated by its subdirectly irreducible members.

Lemma 5. *Let G be a subdirectly irreducible LDLI groupoid. If C_n is a subalgebra of G for some $n \geq 2$, then G contains no fork (i.e. elements $a \neq b$ with $a^2 = b^2$) and, vice versa, if G contains a fork, then C_n is not a subalgebra of G , for any $n \geq 2$.*

Proof. Put $\alpha = \{(a, b) \in G \times G : a^2 = b^2\}$. It is clear that α is an equivalence, which glues each fork. It is a congruence, because whenever $a^2 = b^2$, we get $at = a^2t = b^2t = bt$ and $(ta)^2 = ta^2 = tb^2 = (tb)^2$.

Put $(a, b) \in \beta$ iff there are k, l such that $a^k = b$ and $b^l = a$. By a similar argument, it is easy to see that β is a congruence, which glues each circle.

Clearly, $\alpha \cap \beta = id_G$, hence either $\alpha = id_G$ or $\beta = id_G$ or both, and thus either G contains no fork, or no circle with two or more elements, or both. ■

Remark. One can prove that a subdirectly irreducible LDLI groupoid G either contains a fork, or there is a prime p and a natural number k such that all ip_G -blocks are circles of length either 1, or p^k (this is proven in [9] for LD left quasigroups, however, it is sufficient to assume in the proof LDLI only). On the other hand, there seems to be no uniformity in the former case. In the following example the ip -blocks have different length of tails and, moreover, one contains a ‘pure fork’ (such that $b \neq b^2 = c^2 \neq c$), while the other don’t. The smallest non-trivial congruence has the only non-trivial block $\{a^2, a^3\}$.

	a	a^2	a^3	b	c	b^2	d
a, a^2, a^3	a^2	a^3	a^3	a^2	a^2	a^3	d
$b, c, b^2 = c^2$	a^3	a^3	a^3	b^2	b^2	b^2	d
d	a^3	a^3	a^3	a^2	a^3	a^3	d

Lemma 6. *Let \mathcal{V} be a subvariety of LDLI groupoids and m, n be positive integers. Then $\mathcal{V}_{m,n}$ is the join of the varieties $\mathcal{V}_{m,1}$ and $\mathcal{V}_{1,n}$.*

Proof. Let G be an LDLI groupoid with $x^{m+n} \approx x^m$. Then ip_G -blocks consist of a circle of length k , where $k|n$, and possibly some “tails” of length at most m (precisely, for any element a out of the circle, a^m lies on the circle). Now, assume that G is subdirectly irreducible. It follows from the previous lemma that either all the circles are of length one, or there are no tails (because whenever a tail joins a circle, there is a fork). Hence, G satisfies either $x^{m+1} \approx x^m$ (tails of length at most m only), or $x^{n+1} \approx x$ (circles only). Now, the claim follows from the fact that any variety is generated by its subdirectly irreducible members. ■

Theorem 7. *Let \mathcal{V} be a subvariety of LDLI groupoids and let k, l, m, n be positive integers. Then*

$$\mathcal{V}_{k,l} \vee \mathcal{V}_{m,n} = \mathcal{V}_{\max(k,m), \text{LCM}(l,n)}$$

and

$$\mathcal{V}_{k,l} \wedge \mathcal{V}_{m,n} = \mathcal{V}_{\min(k,m), \text{GCD}(l,n)}.$$

Proof. For the first equality, use the previous lemma and compute $\mathcal{V}_{k,l} \vee \mathcal{V}_{m,n} = \mathcal{V}_{k,1} \vee \mathcal{V}_{m,1} \vee \mathcal{V}_{1,l} \vee \mathcal{V}_{1,n} = \mathcal{V}_{\max(k,m),1} \vee \mathcal{V}_{1,\text{LCM}(l,n)} = \mathcal{V}_{\max(k,m), \text{LCM}(l,n)}$. The second claim is rather clear. ■

Let \mathcal{V} be a variety of LDLI groupoids such that it satisfies no identity $x^{m+n} \approx x^m$. We show that the mapping $(m, n) \mapsto \mathcal{V}_{m,n}$ is injective on $\mathbb{N} \times \mathbb{N}$. The identity $xz \approx yz$ implies an identity $t \approx s$ iff the depth of the rightmost variable in t equals to the depth of the rightmost variable in s and the two variables are identical. Indeed, all identities of \mathcal{V} have the latter property — otherwise $t(x, \dots, x) \approx s(x, \dots, x)$ were a non-trivial identity in one variable. Hence \mathcal{V} contains all groupoids with $xz \approx yz$, i.e., in fact, unary algebras. It is easy to see that for any m_1, n_1 and m_2, n_2 with $(m_1, n_1) \neq (m_2, n_2)$ there is a unary algebra such that it satisfies exactly one of the identities $x^{m_1+n_1} \approx x^{m_1}$, $x^{m_2+n_2} \approx x^{m_2}$. Hence, $\mathcal{V}_{m_1, n_1} \neq \mathcal{V}_{m_2, n_2}$. Moreover, $\mathcal{V}_{1,1}$ contains \mathcal{RZB} always non-trivial.

However, not every subvariety of LDLI groupoids is equal to $\mathcal{V}_{m,n}$ for some \mathcal{V}, m, n . For instance, consider the variety \mathcal{C} of constant groupoids (satisfying the identity $xy \approx uv$); clearly, $x^3 \approx x^2$ holds in \mathcal{C} . Suppose there is a variety \mathcal{V} such that $\mathcal{V}_{2,1} = \mathcal{C}$. Then $\mathcal{V}_{1,1}$ is a non-trivial idempotent subvariety of \mathcal{C} . However, there is no non-trivial idempotent groupoid in \mathcal{C} , a contradiction.

Problem. As shown above, the claim of Theorem 3 does not work for varieties without exponent. Particularly interesting case is the following: describe the structure of free LDLI groupoids modulo free LDI ones.

REFERENCES

- [1] G. Birkhoff, *On the structure of abstract algebras*, Proc. Cambridge Philos. Soc. **31** (1935), 433–454.
- [2] S. Burris and H.P. Sankappanavar, *A course in universal algebra*, Springer, New York 1981 (and also the (electronic) Millennium Edition 1999).
- [3] R. Fenn and C. Rourke, *Racks and links in codimension two*, J. Knot Theory Ramifications **1** (1992), 343–406.
- [4] P. Jedlička, *On left distributive left idempotent groupoids*, Comment. Math. Univ. Carolinae, to appear.
- [5] T. Kepka, *Non-idempotent left symmetric left distributive groupoids*, Comment. Math. Univ. Carolinae **35** (1994), 181–186.
- [6] J. Płonka, *On k -cyclic groupoids*, Math. Japon. **30** (1985), 371–382.
- [7] B. Roszkowska, *The lattice of varieties of symmetric idempotent entropic groupoids*, Demonstratio Math. **20** (1987), 259–275.

- [8] H. Ryder, *The congruence structure of racks*, *Comm. Algebra* **23** (1995), 4971–4989.
- [9] D. Stanovský, *Left distributive left quasigroups*, PhD Thesis, Charles University in Prague, 2004. Available at
<http://www.karlin.mff.cuni.cz/~stanovsk/math/disert.pdf>

Received 13 August 2004
Revised 30 December 2004