

FOLDNESS OF COMMUTATIVE IDEALS IN BCK-ALGEBRAS

CELESTIN LELE

University of Dschang, Department of Mathematics
BP 67 Dschang, Cameroon

e-mail: lele.clele@yahoo.com

AND

SALISSOU MOUTARI

Laboratory J. A Dieudonné UMR CNRS N° 6621
University of Nice-Sophia Antipolis
Parc Valrose 06108 Nice Cedex 2, France

e-mail: salissou@math.unice.fr

Abstract

This paper deals with some properties of n -fold commutative ideals and n -fold weak commutative ideals in BCK-algebras. Afterwards, we construct some algorithms for studying foldness theory of commutative ideals in BCK-algebras.

Keywords: BCK-algebra, fuzzy point, n -fold commutative ideals, n -fold weak commutative ideals.

2000 Mathematics Subject Classification: 06F35, 13A15, 8A72, 03B52, 03E72.

1. INTRODUCTION

The concept of fuzzy subset was introduced in the middle of the sixties by Zadeh [16]. He defined a fuzzy subset of a set X as a function $A : X \rightarrow [0, 1]$. Based on this definition, Xi [15] introduced in 1991 the notion of fuzzy ideals in BCK-algebras. This work enlightened on the

usefulness of ideals theory in general development of BCI/BCK/BL/MV-algebras. From logical point of view, various ideals correspond to various sets of provable formula, see [2, 3, 4, 10, 11, 12] and the references therein.

The tricky point when studying fuzzy mathematics lies in how to carry out the ordinary concept to the fuzzy case. In other words, how to pick out the rational generalization from the large number of available approaches. The particularity of fuzzy ideals compared to ordinary ideals is that one can not say which one of the BCK-algebra elements belongs (or not) to the fuzzy ideals under consideration.

In this paper we study the foldness theory of commutative ideals in BCK-algebras. This theory can be considered as a natural generalization of commutative ideals. Indeed, given any BCK-algebra X , we use the concept of fuzzy point to characterize n -fold commutative ideals in X .

The remainder of this paper is as follows: In Section 2, we recall some important properties of BCK-algebras and their ideals. In Sections 3 and 4 we give some characterizations of n -fold commutative ideals and n -fold weak commutative ideals. Finally, we construct some algorithms for studying n -fold commutative, n -fold weak commutative ideals and their fuzzification in BCK-algebras.

2. BACKGROUND

For some background information see ([1, 2, 5, 10]). An algebra $(X, *, 0)$ of type $(2, 0)$ is called BCK-algebra iff $\forall x, y, z \in X$ the following conditions hold:

$$\text{BCK-1. } ((x * y) * (x * z)) * (z * y) = 0;$$

$$\text{BCK-2. } (x * (x * y)) * y = 0;$$

$$\text{BCK-3. } x * x = 0;$$

$$\text{BCK-4. } 0 * x = 0;$$

$$\text{BCK-5. } x * y = 0 \text{ and } y * x = 0 \implies x = y.$$

A binary relation \leq can be defined on X by

$$\text{BCK-6. } x \leq y \iff x * y = 0,$$

then (X, \leq) is a partially ordered set with the least element 0.

The following properties also hold in any BCK-algebra ([1, 10, 14, 15]):

1. $x * 0 = x$;
2. $x * y = 0$ and $y * z = 0 \implies x * z = 0$;
3. $x * y = 0 \implies (x * z) * (y * z) = 0$ and $(z * y) * (z * x) = 0$;
4. $(x * y) * z = (x * z) * y$;
5. $(x * y) * x = 0$;
6. $x * (x * (x * y)) = x * y$.

Let $(X, *, 0)$ be a BCK-algebra.

A fuzzy subset of a BCK-algebra X is a function

$$\mu : X \longrightarrow [0, 1].$$

Let ξ be the family of all fuzzy sets in X . For $x \in X$ and $\lambda \in (0, 1]$, $x_\lambda \in \xi$ is a fuzzy point iff

$$x_\lambda(y) = \begin{cases} \lambda & \text{if } x = y, \\ 0 & \text{otherwise.} \end{cases}$$

We denote by $\tilde{X} = \{x_\lambda : x \in X, \lambda \in (0, 1]\}$ the set of all fuzzy points on X and we define a binary operation on \tilde{X} as follows:

$$x_\lambda * y_\mu = (x * y)_{\min(\lambda, \mu)}.$$

It is easy to verify that $\forall x_\lambda, y_\mu, z_\alpha \in \tilde{X}$, the following conditions hold:

$$\text{BCK-1'}. \quad ((x_\lambda * y_\mu) * (x_\lambda * z_\alpha)) * (z_\alpha * y_\mu) = 0_{\min(\lambda, \mu, \alpha)};$$

$$\text{BCK-2'}. \quad [x_\lambda * (x_\lambda * y_\mu)] * y_\mu = 0_{\min(\lambda, \mu)};$$

$$\text{BCK-3'}. \quad x_\lambda * x_\mu = 0_{\min(\lambda, \mu)};$$

$$\text{BCK-4'}. \quad 0_\mu * x_\lambda = 0_{\min(\lambda, \mu)}.$$

Remark 2.1. The condition BCK-5. is not true in $(\tilde{X}, *)$. So the partial order \leq in $(X, *)$ can not be extended to $(\tilde{X}, *)$.

We can also establish the following conditions $\forall x_\lambda, y_\mu, z_\alpha \in \tilde{X}$:

- 1'. $x_\lambda * 0_\mu = x_{\min(\lambda, \mu)}$;
- 2'. $x_\lambda * y_\mu = 0_{\min(\lambda, \mu)}$ and $y_\mu * z_\alpha = 0_{\min(\mu, \alpha)} \implies x_\lambda * z_\alpha = 0_{\min(\lambda, \mu)}$;
- 3'. $x_\lambda * y_\mu = 0_{\min(\lambda, \mu)} \implies (x_\lambda * z_\alpha) * (y_\mu * z_\alpha) = 0_{\min(\lambda, \mu, \alpha)}$ and $(z_\alpha * y_\mu) * (z_\alpha * x_\lambda) = 0_{\min(\lambda, \mu, \alpha)}$;
- 4'. $(x_\lambda * y_\mu) * z_\alpha = (x_\lambda * z_\alpha) * y_\mu$;
- 5'. $(x_\lambda * y_\mu) * x_\lambda = 0_{\min(\lambda, \mu)}$;
- 6'. $x_\lambda * (x_\lambda * (x_\lambda * y_\mu)) = x_\lambda * y_\mu$.

We recall that if A is a fuzzy subset of a BCK-algebra X , then we have the following:

$$(1) \quad \tilde{A} = \{x_\lambda \in \tilde{X} : A(x) \geq \lambda, \lambda \in (0, 1]\}.$$

$$(2) \quad \forall \lambda \in (0, 1], \tilde{X}_\lambda = \{x_\lambda : x \in X\}, \text{ and } \tilde{A}_\lambda = \{x_\lambda \in \tilde{X}_\lambda : A(x) \geq \lambda\}.$$

We have also $\tilde{X}_\lambda \subseteq \tilde{X}$, $\tilde{A} \subseteq \tilde{X}$, $\tilde{A}_\lambda \subseteq \tilde{A}$, $\tilde{A}_\lambda \subseteq \tilde{X}_\lambda$ and one can easily check that $(\tilde{X}_\lambda, *, 0_\lambda)$ is a BCK-algebra.

Definition 2.1 [3]. A nonempty subset I of a BCK-algebra X is called an ideal if it satisfies

1. $0 \in I$;
2. $x * y \in I$ and $y \in I \implies x \in I$.

Definition 2.2 [3]. A fuzzy subset A of a BCK-algebra X is a fuzzy ideal iff

1. $\forall x \in X, A(0) \geq A(x)$;
2. $\forall x, y \in X, A(x) \geq \min(A(x * y), A(y))$.

Definition 2.3. \tilde{A} is a weak ideal of \tilde{X} iff

- 1) $\forall \nu \in Im(A), 0_\nu \in \tilde{A}$;
- 2) $\forall x_\lambda, y_\mu \in \tilde{X}$, such that $x_\lambda * y_\mu \in \tilde{A}$ and $y_\mu \in \tilde{A}$, we have $x_{\min(\lambda, \mu)} \in \tilde{A}$.

Remark 2.2. Any weak ideal \tilde{A} has the following property

$$(x_\lambda * y_\mu = 0_{\min(\lambda, \mu)} \text{ and } y_\mu \in \tilde{A}) \Rightarrow x_{\min(\lambda, \mu)} \in \tilde{A}.$$

Proof. Let $x_\lambda, y_\mu \in \tilde{X}$ such that $x_\lambda * y_\mu = 0_{\min(\lambda, \mu)}$ and $y_\mu \in \tilde{A}$.

$$y_\mu \in \tilde{A} \implies A(y) \geq \mu.$$

Let $A(y) = \alpha$, using Definition 2.3 - 1) we obtain $0_\alpha \in \tilde{A}$.

So $A(0) \geq \alpha$. But $\alpha = A(y) \geq \mu \geq \min(\lambda, \mu)$. Therefore $0_{\min(\lambda, \mu)} \in \tilde{A}$.
Finally, according to Definition 2.3 - 2), we have $x_{\min(\lambda, \mu)} \in \tilde{A}$. ■

A characterization of a weak ideal is given by the following theorem.

Theorem 2.1 [10]. Suppose that A is a fuzzy subset of a BCK-algebra X , then the following conditions are equivalent:

1. A is a fuzzy ideal;
2. $\forall x_\lambda, y_\mu \in \tilde{A}, (z_\alpha * y_\mu) * x_\lambda = 0_{\min(\lambda, \mu, \alpha)} \implies z_{\min(\lambda, \mu, \alpha)} \in \tilde{A}$;
3. $\forall t \in (0, 1]$, the t -level subset $A^t = \{x \in X : A(x) \geq t\}$ is an ideal when $A^t \neq \emptyset$;
4. \tilde{A} is a weak ideal.

3. FUZZY n -FOLD COMMUTATIVE WEAK IDEALS

Throughout this paper, X always means a BCK-algebra and \tilde{X} the set of fuzzy points on X .

Let us denote $(\dots((x*y)*y)*\dots)*y$ by $x*y^n$ and $(\dots((x_\lambda*y_\mu)*y_\mu)*\dots)*y_\mu$ by $x_\lambda * y_\mu^n$ (where y and y_μ occurs respectively n times) with $x, y \in X$, $x_\lambda, y_\mu \in \tilde{X}$.

We recall the following:

Definition 3.1. An nonempty subset I of a BCK-algebra X is called a commutative ideal of X if it satisfies

1. $0 \in I$;
2. $\forall x, y, z \in X, ((x * y) * z \in I \text{ and } z \in I) \implies x * (y * (y * x)) \in I$.

An ideal I of a BCK-algebra X is commutative iff

$$\forall x, y \in X, x * y \in I \implies x * (y * (y * x)) \in I.$$

Lemma 3.1 [10]. For any fuzzy ideal A of X , if $x \leq y$, then $A(y) \leq A(x)$.

Definition 3.2. A BCK-algebra X is n -fold commutative if for any $x, y \in X$, $x * y = x * (y * (y * x^n))$.

Theorem 3.1 [13]. A BCK-algebra X is n -fold commutative iff for any $x, y \in X$, $x * (x * y) \leq y * (y * x^n)$.

Definition 3.3. A nonempty subset I of a BCK-algebra X is an n -fold commutative ideal of X if it satisfies

1. $0 \in I$;
2. $\forall x, y, z \in X$,

$$((x * y) * z \in I \text{ and } z \in I) \implies x * (y * (y * x^n)) \in I.$$

Lemma 3.2 [13]. *An ideal I of a BCK-algebra X is an n -fold commutative ideal iff*

$$\forall x, y \in X, x * y \in I \implies x * (y * (y * x^n)) \in I.$$

Now, we give some characterizations of fuzzy n -fold commutative ideals in BCK-algebras.

Definition 3.4. A fuzzy subset A of X is called a fuzzy n -fold commutative ideal of X if it satisfies

1. $\forall x \in X, A(0) \geq A(x)$;
2. $\forall x, y, z \in X, A(x * (y * (y * x^n))) \geq \min(A((x * y) * z), A(z))$.

Definition 3.5 [10]. \tilde{A} is a commutative weak ideal of \tilde{X} iff

1. $\forall \nu \in Im(A), 0_\nu \in \tilde{A}$;
2. $\forall x_\lambda, y_\mu, z_\alpha \in \tilde{X}$ such that $(x_\lambda * y_\mu) * z_\alpha \in \tilde{A}$ and $z_\alpha \in \tilde{A}$, we have

$$x_{\min(\lambda, \alpha)} * (y_\mu * (y_\mu * x_{\min(\lambda, \alpha)})) \in \tilde{A}.$$

Definition 3.6. \tilde{A} is an n -fold commutative weak ideal of \tilde{X} iff

1. $\forall \nu \in Im(A), 0_\nu \in \tilde{A}$;
2. $\forall x_\lambda, y_\mu, z_\alpha \in \tilde{X}$, if $(x_\lambda * y_\mu) * z_\alpha \in \tilde{A}$ and $z_\alpha \in \tilde{A}$, then

$$x_{\min(\lambda, \alpha)} * (y_\mu * (y_\mu * x_{\min(\lambda, \alpha)}^n)) \in \tilde{A}.$$

Example 3.1. Let $X = \{0, 1, 2, 3, 4\}$ with $*$ defined by the following table:

$*$	0	1	2	3	4
0	0	0	0	0	0
1	1	0	0	0	0
2	2	1	0	1	0
3	3	3	3	0	0
4	4	4	4	4	0

By simple computations one can prove that $(X, *, 0)$ is a BCK-algebra.

Let $t_1, t_2 \in (0, 1]$ and let define a fuzzy subset $A : X \rightarrow [0, 1]$ by

$$t_1 = A(0) = A(1) = A(2) = A(3) > A(4) = t_2.$$

One can easily check that for any $n > 2$,

$$\begin{aligned} \tilde{A} = & \{0_\lambda : \lambda \in (0, t_1]\} \cup \{1_\lambda : \lambda \in (0, t_1]\} \cup \{2_\lambda : \lambda \in (0, t_1]\} \\ & \cup \{3_\lambda : \lambda \in (0, t_1]\} \cup \{4_\lambda : \lambda \in (0, t_1]\} \end{aligned}$$

is an n -fold commutative weak ideal.

Remark 3.1. \tilde{A} is a 1-fold commutative weak ideal of a BCK-algebra X iff \tilde{A} is a commutative weak ideal of X .

Theorem 3.2. *If A is a fuzzy subset of X , then A is a fuzzy n -fold commutative ideal iff \tilde{A} is an n -fold commutative weak ideal.*

Proof.

\implies – Let $\lambda \in \text{Im}(A)$, it is easy to prove that $0_\lambda \in \tilde{A}$;

– Let $(x_\lambda * y_\mu) * z_\alpha \in \tilde{A}$ and $z_\alpha \in \tilde{A}$,

$$A((x * y) * z) \geq \min(\lambda, \mu, \alpha) \text{ and } A(z) \geq \alpha.$$

Since A is a fuzzy n -fold commutative ideal, we have

$$A(x * (y * (y * x^n))) \geq \min(A((x * y) * z),$$

$$A(z)) \geq \min(\min(\lambda, \mu, \alpha), \alpha) = \min(\lambda, \mu, \alpha).$$

Therefore,

$$(x * (y * (y * x^n)))_{\min(\lambda, \mu, \alpha)} = x_{\min(\lambda, \alpha)} * (y_{\mu} * (y_{\mu} * x_{\min(\lambda, \alpha)}^n)) \in \tilde{A}.$$

\Leftarrow – Let $x \in X$, it is easy to prove that $A(0) \geq A(x)$.

– Let $x, y, z \in X$ and let $A((x * y) * z) = \beta$ and $A(z) = \alpha$, then

$$((x * y) * z)_{\min(\beta, \alpha)} = (x_{\beta} * y_{\alpha}) * z_{\alpha} \in \tilde{A} \text{ and } z_{\alpha} \in \tilde{A}.$$

Since \tilde{A} is n -fold commutative weak ideal, we have

$$x_{\min(\beta, \alpha)} * (y_{\alpha} * (y_{\alpha} * x_{\min(\beta, \alpha)}^n)) = (x * (y * (y * x^n)))_{\min(\beta, \alpha)} \in \tilde{A}.$$

Thus $A(x * (y * (y * x^n))) \geq \min(\beta, \alpha) = \min(A((x * y) * z), A(z))$. ■

Proposition 3.1. *In an n -fold commutative BCK-algebra, every weak ideal is an n -fold commutative weak ideal.*

Proof. The proof is straightforward. ■

Corollary 3.1. *In an n -fold commutative BCK-algebra, every fuzzy ideal is a fuzzy n -fold commutative ideal.*

Proposition 3.2. *An n -fold commutative weak ideal is an ideal. But the converse does not hold in general.*

Proof. Let $x_{\lambda}, y_{\mu} \in \tilde{A}$, then

$$x_{\lambda} * y_{\mu} = (x_{\lambda} * 0_{\mu}) * y_{\mu} \in \tilde{A}.$$

Since \tilde{A} is an n -fold commutative weak ideal, we have

$$x_{\min(\lambda,\mu)} = x_{\min(\lambda,\mu)} * (0_\mu * (0_\mu * x_{\min(\lambda,\mu)}^n)) \in \tilde{A}.$$

For the converse, let $X = \{0, 1, 2, 3, 4\}$ with the binary operation $*$ defined by the following table:

*	0	1	2	3	4
0	0	0	0	0	0
1	1	0	1	0	0
2	2	2	0	0	0
3	3	3	3	0	0
4	4	4	4	3	0

Obviously, $(X, *, 0)$ is a BCK-algebra.

Let us define a fuzzy subset $A : X \rightarrow [0, 1]$ by

$$A(0) = 1, \quad A(1) = \frac{1}{2}, \quad A(2) = A(3) = A(4) = \frac{1}{3}.$$

It is easy to check that

$$\begin{aligned} \tilde{A} = & \{0_\lambda : \lambda \in (0, 1]\} \cup \left\{1_\lambda : \lambda \in \left(0, \frac{1}{2}\right]\right\} \cup \left\{2_\lambda : \lambda \in \left(0, \frac{1}{3}\right]\right\} \\ & \cup \left\{3_\lambda : \lambda \in \left(0, \frac{1}{3}\right]\right\} \cup \left\{4_\lambda : \lambda \in \left(0, \frac{1}{3}\right]\right\} \end{aligned}$$

is a weak ideal, but not an n -fold commutative weak ideal because

$$(2_1 * 3_1) * 0_1 = 0_1 \in \tilde{A} \text{ and } 0_1 \in \tilde{A}, \text{ but } 2_1 * (3_1 * (3_1 * 2_1^n)) = 2_1 \notin \tilde{A}. \quad \blacksquare$$

Corollary 3.2. *A fuzzy n -fold commutative ideal is a fuzzy ideal. But the converse does not hold in general.*

The following theorem gives a characterization of an n -fold commutative weak ideal.

Theorem 3.3. *Suppose that \tilde{A} is a weak ideal (namely A is a fuzzy ideal by Theorem 2.1), then the following conditions are equivalent:*

1. A is a fuzzy n -fold commutative ideal;
2. $\forall x_\lambda, y_\mu \in \tilde{X}$ such that $x_\lambda * y_\lambda \in \tilde{A}$, we have

$$x_{\min(\lambda, \mu)} * \left(y_\mu * \left(y_\mu * x_{\min(\lambda, \mu)}^n \right) \right) \in \tilde{A};$$

3. $\forall t \in (0, 1]$, the t -level subset $A^t = \{x \in X : A(x) \geq t\}$ is an n -fold commutative ideal when $A^t \neq \emptyset$;
4. $\forall x, y, z \in X$, $A(x * (y * (y * x^n))) \geq A(x * y)$;
5. \tilde{A} is an n -fold commutative weak ideal.

Proof.

1. \Rightarrow 2. Let $x_\lambda * y_\mu \in \tilde{A}$. Since A is a fuzzy n -fold commutative, we have

$$A(x * (y * (y * x^n))) \geq \min(A((x * y) * (x * y))),$$

$$A(x * y) \geq \min(A(0), A(x * y)) = A(x * y) \geq \min(\lambda, \mu).$$

Therefore,

$$(x * (y * (y * x^n)))_{\min(\lambda, \mu)} = x_{\min(\lambda, \mu)} * \left(y_\mu * \left(y_\mu * x_{\min(\lambda, \mu)}^n \right) \right) \in \tilde{A}.$$

2. \Rightarrow 3. – Obviously, $\forall t \in (0, 1]$, $0 \in A^t$.

– Let $(x * y) * z \in A^t$ and $z \in A^t$, then we have

$$((x * y) * z)_t = (x_t * y_t) * z_t \in \tilde{A} \text{ and } z_t \in \tilde{A}.$$

Since \tilde{A} is a weak ideal, we have $x_t * y_t = (x * y)_t \in \tilde{A}$. Using the hypothesis, we obtain

$$x_t * (y_t * (y_t * x_t^n)) = (x * (y * (y * x^n)))_t \in \tilde{A}, \text{ hence } x * (y * (y * x^n)) \in A^t.$$

By virtue of Lemma 3.2, we obtain that $A^t = \{x \in X : A(x) \geq t\}$ is an n -fold commutative ideal.

3. \Rightarrow 4. Let $x, y \in X$ and $t = A(x * y)$, then $x * y \in A^t$. Since A^t is an n -fold commutative ideal, we have

$$x * (y * (y * x^n)) \in A^t, \text{ hence } A(x * (y * (y * x^n))) \geq t = A(x * y).$$

4. \Rightarrow 5. – Let $\lambda \in \text{Im}(A)$. Obviously, $0_\lambda \in \tilde{A}$.

– Let $(x_\lambda * y_\mu) * z_\alpha \in \tilde{A}$ and $z_\alpha \in \tilde{A}$. Since \tilde{A} is a weak ideal, we obtain $(x * y)_{\min(\lambda, \mu, \alpha)} \in \tilde{A}$. According to the hypothesis, we obtain

$$A(x * (y * (y * x^n))) \geq A(x * y) \geq \min(\lambda, \mu, \alpha),$$

hence

$$\begin{aligned} & \left(x * (y * (y * x^n)) \right)_{\min(\lambda, \mu, \alpha)} \\ &= x_{\min(\lambda, \mu)} * \left(y_\mu * (y_\mu * x_{\min(\lambda, \alpha)}^n) \right) \in \tilde{A}. \end{aligned}$$

5. \Rightarrow 1. Follows from Theorem 3.2. ■

Theorem 3.4. *Let \tilde{A} and \tilde{B} be two weak ideals such that $\tilde{A} \subseteq \tilde{B}$, and $A(0) = B(0)$. If \tilde{A} is an n -fold commutative weak ideal, then \tilde{B} is also n -fold commutative weak ideal.*

Proof. To prove the theorem, we need the following result.

Lemma 3.3 [13]. *If I and J are two ideals of X such that $I \subseteq J$ with I n -fold commutative, then J is also n -fold commutative.*

Using this lemma, we can prove Theorem 3.4 as follows:

To prove that \tilde{B} is n -fold commutative, it suffices to show that $\forall t \in (0, 1]$, B^t is n -fold commutative ideal when $B^t \neq \emptyset$.

Since $A(0) = B(0)$, it is clear that $A^t \neq \emptyset$ when $B^t \neq \emptyset$.

$$\tilde{A} \subseteq \tilde{B} \implies A^t \subseteq B^t.$$

Since \tilde{A} is n -fold commutative, A^t is also n -fold commutative. According to Lemma 3.3, B^t is also n -fold commutative. So, \tilde{B} is also n -fold commutative. ■

Consequence 3.1. $\forall \lambda \in Im(A)$, if $\{0_\lambda\}$ is an n -fold commutative weak ideal, then \tilde{A} is also an n -fold commutative weak ideal.

Corollary 3.3. *Let A and B be two fuzzy ideals of X such that $A \leq B$ and $B(0) = A(0)$. If A is a fuzzy n -fold commutative ideal, then B is also a fuzzy n -fold commutative ideal.*

4. FUZZY n -FOLD WEAK COMMUTATIVE WEAK IDEALS

In this section, we define and give some characterizations of fuzzy n -fold weak commutative weak ideals in BCK-algebras.

Let us recall the following results.

Definition 4.1. A nonempty subset I of X is called an n -fold weak commutative ideal of X if it satisfies

1. $0 \in I$;
2. $\forall x, y, z \in X$, $(x * (x * y^n)) * z \in I$ and $z \in I \implies y * (y * x) \in I$.

Lemma 4.1 [13]. *An ideal I of a BCK-algebra X is an n -fold weak commutative ideal iff*

$$\forall x, y, z \in X, x * (x * y^n) \in I \implies y * (y * x) \in I.$$

Definition 4.2. A fuzzy subset A of X is called a fuzzy n -fold weak commutative ideal of X if it satisfies

1. $\forall x \in X$, $A(0) \geq A(x)$;
2. $\forall x, y, z \in X$, $A(y * (y * x)) \geq \min(A((x * (x * y^n)) * z), A(z))$.

Definition 4.3. \tilde{A} is a weak commutative weak ideal of \tilde{X} iff

1. $\forall \nu \in Im(A)$, $0_\nu \in \tilde{A}$;
2. $\forall x_\lambda, y_\mu, z_\alpha \in \tilde{X}$,
 $((x_\lambda * (x_\lambda * y_\mu)) * z_\alpha \in \tilde{A} \text{ and } z_\alpha \in \tilde{A}) \implies (y_\mu * (y_\mu * x_{\min(\lambda, \alpha)})) \in \tilde{A}$.

Definition 4.4. \tilde{A} is an n -fold weak commutative weak ideal of \tilde{X} iff

1. $\forall \nu \in Im(A), 0_\nu \in \tilde{A}$;
2. $\forall x_\lambda, y_\mu, z_\alpha \in \tilde{X}$,

$$((x_\lambda * (x_\lambda * y_\mu^n)) * z_\alpha \in \tilde{A} \text{ and } z_\alpha \in \tilde{A}) \Rightarrow (y_\mu * (y_\mu * x_{\min(\lambda, \alpha)})) \in \tilde{A}.$$

Example 4.1. Let $X = \{0, 1, 2, 3\}$ with $*$ defined by the following table:

*	0	1	2	3
0	0	0	0	0
1	1	0	0	0
2	2	2	0	0
3	3	3	3	0

By simple computations one can prove that $(X, *, 0)$ is a BCK-algebra. Let $t_1, t_2 \in (0, 1]$ and let us define a fuzzy subset $A : X \rightarrow [0, 1]$ by

$$t_1 = A(0) = A(1) = A(2) > A(3) = t_2.$$

It is easy to check that for any $n > 2$,

$$\begin{aligned} \tilde{A} = & \{0_\lambda : \lambda \in (0, t_1]\} \cup \{1_\lambda : \lambda \in (0, t_1]\} \\ & \cup \{2_\lambda : \lambda \in (0, t_1]\} \cup \{3_\lambda : \lambda \in (0, t_2]\} \end{aligned}$$

is an n -fold weak commutative weak ideal.

Remark 4.1. \tilde{A} is an 1-fold weak commutative weak ideal of a BCK-algebra X iff \tilde{A} is a weak commutative weak ideal.

Theorem 4.1. *If A is a fuzzy subset of X , then A is a fuzzy n -fold weak commutative ideal iff \tilde{A} is an n -fold weak commutative weak ideal.*

Proof.

\implies – Let $\lambda \in \text{Im}(A)$. Obviously $0_\lambda \in \tilde{A}$;

– Let $(x_\lambda * (x_\lambda * y_\mu^n)) * z_\alpha \in \tilde{A}$ and $z_\alpha \in \tilde{A}$, then

$$A((x * (x * y^n)) * z) \geq \min(\lambda, \mu, \alpha) \text{ and } A(z) \geq \alpha.$$

Since A is a fuzzy n -fold weak commutative ideal, we have

$$A(y * (y * x)) \geq \min(A((x * (x * y)) * z),$$

$$A(z)) \geq \min(\min(\lambda, \mu, \alpha), \alpha) = \min(\lambda, \mu, \alpha).$$

Therefore $(y * (y * x))_{\min(\lambda, \mu, \alpha)} = y_\mu * (y_\mu * x_{\min(\lambda, \alpha)}) \in \tilde{A}$.

\Leftarrow – Let $x \in X$, it is easy to prove that $A(0) \geq A(x)$;

– Let $x, y, z \in X$, $A((x * (x * y^n)) * z) = \beta$ and $A(z) = \alpha$.

Then,

$$((x * (x * y^n)) * z)_{\min(\beta, \alpha)} = (x_\beta * (x_\beta * y_\beta^n)) * z_\alpha \in \tilde{A} \text{ and } z_\alpha \in \tilde{A}.$$

Since \tilde{A} is n -fold weak commutative weak ideal, we have

$$y_\beta * (y_\beta * x_{\min(\beta, \alpha)}) = (y * (y * x))_{\min(\beta, \alpha)} \in \tilde{A}.$$

Hence, $A(y * (y * x)) \geq \min(\beta, \alpha) = \min(A((x * (x * y^n)) * z), A(z))$. ■

Proposition 4.1. *In an n -fold commutative BCK-algebra, the concepts of weak ideals, n -fold commutative weak ideals and n -fold weak commutative weak ideals are the same.*

Proof. The proof is straightforward. ■

Corollary 4.1. *In an n -fold commutative BCK-algebra, the concepts of fuzzy ideals, fuzzy n -fold commutative ideals and fuzzy n -fold weak commutative ideals are the same.*

Proposition 4.2. *An n -fold weak commutative weak ideal is a weak ideal.*

Proof. By setting $y_\mu = x_\lambda$ in Definition 4.4 and using the fact that $x*x = 0$ and $x*0 = x$, one obtains that

$$\forall x_\lambda, z_\alpha \in \tilde{X} \text{ such that } x_\lambda * z_\alpha \in \tilde{A} \text{ and } z_\alpha \in \tilde{A}, x_{\min(\lambda, \alpha)} \in \tilde{A}. \quad \blacksquare$$

Corollary 4.2. *A fuzzy n -fold weak commutative ideal is a fuzzy ideal.*

The following theorem summarizes a characterization of an n -fold weak commutative weak ideal.

Theorem 4.2. *Suppose that \tilde{A} is a weak ideal (namely A is a fuzzy ideal by Theorem 2.1), then the following conditions are equivalent:*

- 1) A is fuzzy n -fold weak commutative ideal;
- 2) $\forall x_\lambda, y_\mu \in \tilde{X}$ such that $x_\lambda * (x_\lambda * y_{\min(\lambda, \mu)}^n) \in \tilde{A}$, we have

$$y_\mu * (y_\mu * x_{\min(\lambda, \mu)}) \in \tilde{A};$$

- 3) $\forall t \in (0, 1]$, the t -level subset $A^t = \{x \in X : A(x) \geq t\}$ is an n -fold weak commutative ideal when $A^t \neq \emptyset$;
- 4) $\forall x, y \in X$, $A(y * (y * x)) \geq A(x * (x * y^n))$;
- 5) \tilde{A} is an n -fold weak commutative weak ideal.

Proof.

- 1) \Rightarrow 2) Let $x_\lambda * (x_\lambda * y_{\min(\lambda, \mu)}^n) \in \tilde{A}$. Since A is a fuzzy n -fold weak commutative ideal, we have

$$A(y * (y * x)) \geq \min(A((x * (x * y^n)) * 0),$$

$$A(0)) = A((x * (x * y^n))) \geq \min(\lambda, \mu).$$

Therefore, $(y * (y * x))_{\min(\lambda, \mu)} = y_\mu * (y_\mu * x_{\min(\lambda, \mu)}) \in \tilde{A}$.

2) \Rightarrow 3) – Obviously, $\forall t \in (0, 1]$, $0 \in A^t$.

– Let $x * (x * y^n) \in A^t$, we have

$$(x * (x * y^n))_t = x_t * (x_t * y_t^n) \in \tilde{A}.$$

By virtue of the hypothesis, one obtains $y_t * (y_t * x_t) \in \tilde{A}$, therefore $y * (y * x) \in A^t$. Using Lemma 4.1, we can conclude that $A^t = \{x \in X : A(x) \geq t\}$ is an n -fold weak commutative ideal.

3) \Rightarrow 4) Let $x, y \in X$ and $t = A(x * (x * y^n))$, then $x * (x * y^n) \in A^t$. Since A^t is an n -fold weak commutative ideal, we have

$$y * (y * x) \in A^t, \text{ therefore } A(y * (y * x)) \geq t = A(x * (x * y^n)).$$

4) \Rightarrow 5) – Let $\lambda \in \text{Im}(A)$, it is clear that $0_\lambda \in \tilde{A}$.

– Let $(x_\lambda * (x_\lambda * y_\mu^n)) * z_\alpha \in \tilde{A}$ and $z_\alpha \in \tilde{A}$. Since \tilde{A} is a weak ideal, $(x * (x * y^n))_{\min(\lambda, \mu, \alpha)} \in \tilde{A}$. Using the hypothesis, we obtain

$$A(y * (y * x)) \geq A(x * (x * y^n)) \geq \min(\lambda, \mu, \alpha).$$

From this, one can deduce that

$$(y * (y * x))_{\min(\lambda, \mu, \alpha)} = y_\mu * (y_\mu * x_{\min(\lambda, \alpha)}) \in \tilde{A}.$$

5) \Rightarrow 1) Follows from Theorem 4.1. ■

Theorem 4.3. *Theorem 3.4, its corollary (Corollary 3.3) and consequence (Consequence 3.1) are valid if “ n -fold commutative” is replaced by “ n -fold weak commutative”.*

Proof.

The proof is similar to that of Theorem 3.4 and is therefore omitted. ■

Appendix A

ALGORITHMS

Algorithm for BCK-algebras

Input(X : set, $*$: binary operation)
Output(“ X is a BCK-algebra or not”)
Begin
 If $X = \emptyset$ **then**
 go to (1.);
 EndIf
 If $0 \notin X$ **then**
 go to (1.);
 EndIf
 $Stop := false$;
 $i := 1$;
 While $i \leq |X|$ **and not**($Stop$) **do**
 If $x_i * x_i \neq 0$ **then**
 $Stop := true$;
 EndIf
 If $0 * x_i \neq 0$ **then**
 $Stop := true$;
 EndIf
 $j := 1$
 While $j \leq |X|$ **and not**($Stop$) **do**
 If $(x_i * (x_i * y_j)) * y_j \neq 0$ **then**
 $Stop := true$;
 EndIf
 If $(x_i * y_j = 0)$ **and** $(y_j * x_i = 0)$ **then**
 If $x_i \neq y_j$ **then**
 $Stop := true$;
 EndIf
 EndIf
 $k := 1$;
 While $k \leq |X|$ **and not**($Stop$) **do**
 If $((x_i * y_j) * (x_i * z_k)) * (z_k * y_j) \neq 0$ **then**
 $Stop := true$;
 EndIf
 EndWhile
 EndWhile
 EndWhile
 If $Stop$ **then**
 (1.) **Output**(“ X is not a BCK-algebra”)
 Else
 Output(“ X is a BCK-algebra”)
 EndIf
End

Algorithm for ideals of BCK-algebras

Input(X : BCK-algebra, I : subset of X);
Output(“ I is an ideal of X or not”);
Begin
 If $I = \emptyset$ **then**
 go to (1.);
 EndIf
 If $0 \notin I$ **then**
 go to (1.);
 EndIf
 $Stop := false$;
 $i := 1$;
 While $i \leq |X|$ **and not**($Stop$) **do**
 $j := 1$
 While $j \leq |X|$ **and not**($Stop$) **do**
 If $x_i * y_j \in I$ **and** $y_j \in I$ **then**
 If $x_i \notin I$ **then**
 $Stop := true$;
 EndIf
 EndIf
 EndWhile
 EndWhile
 If $Stop$ **then**
 Output(“ I is an ideal of X ”)
 Else
 (1.) **Output**(“ I is not an ideal of X ”)
 EndIf
End

Algorithm for n -fold commutative ideals

Input(X : BCK-algebra, I : subset of X , $n \in \mathbb{N}$);
Output(“ I is an n -fold commutative ideal of X or not”);
Begin
 If $I = \emptyset$ **then**
 go to (1.);
 EndIf
 If $0 \notin I$ **then**
 go to (1.);
 EndIf
 $Stop := false$;
 $i := 1$;
 While $i \leq |X|$ **and not**($Stop$) **do**
 $j := 1$
 While $j \leq |X|$ **and not**($Stop$) **do**
 $k := 1$
 While $k \leq |X|$ **and not**($Stop$) **do**
 If $(x_i * y_j) * z_k \in I$ **and** $z_k \in I$ **then**
 If $x_i * (y_j * (y_j * x_i^n)) \notin I$ **then**
 $Stop := true$;
 EndIf
 EndIf
 EndWhile
 EndWhile
 EndWhile
 If $Stop$ **then**
 Output(“ I is an n -fold commutative ideal of X ”)
 Else
 (1.)**Output**(“ I is not an n -fold commutative ideal of X ”)
 EndIf
End

Algorithm for n -fold weak commutative ideals

Input(X : BCK-algebra, I : subset of X , $n \in \mathbb{N}$);

Output(“ I is an n -fold weak commutative ideal of X or not”);

Begin
If $I = \emptyset$ **then**

go to (1.);

EndIf
If $0 \notin I$ **then**

go to (1.);

EndIf
 $Stop := false$;

 $i := 1$;

While $i \leq |X|$ **and not**($Stop$) **do**
 $j := 1$
While $j \leq |X|$ **and not**($Stop$) **do**
 $k := 1$
While $k \leq |X|$ **and not**($Stop$) **do**
If $(x_i * (x_i * y_j^n)) * z_k \in I$ **and** $z_k \in I$ **then**
If $y_j * (y_j * x_i) \notin I$ **then**
 $Stop := true$;

EndIf
EndIf
EndWhile
EndWhile
EndWhile
If $Stop$ **then**
Output(“ I is an n -fold weak commutative ideal of X ”)

Else

(1.) **Output**(“ I is not an n -fold weak commutative ideal of X ”)

EndIf
End

Algorithm for fuzzy subsets

Input($X : BCI\text{-algebra}, A : X \rightarrow [0, 1]$);
Output(“ A is a fuzzy subset of X or not”);
Begin
 $Stop := false$;
 $i := 1$;
 While $i \leq |X|$ **and not**($Stop$) **do**
 If ($A(x_i) < 0$) **or** ($A(x_i) > 1$) **then**
 $Stop := true$;
 EndIf
 EndWhile
 If $Stop$ **then**
 Output(“ A is a fuzzy subset of X ”)
 Else
 Output(“ A is not a fuzzy subset of X ”)
 EndIf
End

Algorithm for fuzzy n -fold commutative ideals

Input($X : BCK\text{-algebra}, *: \text{binary operation}, A : \text{fuzzy subset of } X$);
Output(“ A is a fuzzy n -fold commutative ideal of X or not”);
Begin
 $Stop := false$;
 $i := 1$;
 While $i \leq |X|$ **and not**($Stop$) **do**
 If $A(0) < A(x_i)$ **then**
 $Stop := true$;
 EndIf
 $j := 1$
 While $j \leq |X|$ **and not**($Stop$) **do**
 $k := 1$;
 While $k \leq |X|$ **and not**($Stop$) **do**
 If $A(x_i * (y_j * (y_j * x_i^n))) < \text{Min}(A(x_i * y_j) * z_k), A(z_k)$ **then**
 $Stop := true$;
 EndIf
 EndWhile
 EndWhile
 EndWhile
 If $Stop$ **then**
 Output(“ A is not a fuzzy n -fold commutative ideal of X ”)
 Else
 Output(“ A is a fuzzy n -fold commutative ideal of X ”)
 EndIf
End

Algorithm for fuzzy n -fold weak commutative ideals

Input(X : BCK-algebra, $*$: binary operation, A : fuzzy subset of X);
Output(“ A is a fuzzy n -fold weak commutative ideal of X or not”);
Begin
 $Stop := false$;
 $i := 1$;
 While $i \leq |X|$ **and not**($Stop$) **do**
 If $A(0) < A(x_i)$ **then**
 $Stop := true$;
 EndIf
 $j := 1$
 While $j \leq |X|$ **and not**($Stop$) **do**
 $k := 1$;
 While $k \leq |X|$ **and not**($Stop$) **do**
 If $A(y_j * (y_j * x_i)) < Min(A((x_i * (x_i * y_j^n) * z_k)), A(z_k))$ **then**
 $Stop := true$;
 EndIf
 EndWhile
 EndWhile
 EndWhile
 If $Stop$ **then**
 Output(“ A is not a fuzzy n -fold weak commutative ideal of X ”)
 Else
 Output(“ A is a fuzzy n -fold weak commutative ideal of X ”)
 EndIf
End

REFERENCES

- [1] J. Meng, *On ideals in BCK-algebras*, Math. Japonica **40** (1994), 143–154.
- [2] J. Meng and X. Guo, *On fuzzy ideals in BCK/BCI-algebras*, Fuzzy Sets and Systems **149** (2005), 509–525.
- [3] J. Meng, Y.B. Jun and H.S. Kim, *Fuzzy implicative ideals of BCK-algebras*, Fuzzy Sets and Systems **89** (1997), 243–248.
- [4] Y.B. Jun and E.H. Roh, *Fuzzy commutative ideals in BCK-algebras*, Fuzzy Sets and Systems **64** (1994), 401–405.
- [5] Y.B. Jun, *Characterization of Noetherian in BCK-algebras via fuzzy ideals*, Fuzzy Sets and Systems **108** (1999), 231–234.
- [6] Y.B. Jun, S.Z. Song and C. Lele, *Foldness of quasi-associative ideals in BCI-algebras*, Scientiae Mathematicae **6** (2002), 227–231.
- [7] Y.B. Jun and K.H. Kim, *Fuzzifications of foldness of quasi-associative ideals in BCI-algebras*, J. Appl. Math. Computing **11** (2003), 255–263.
- [8] Y.B. Jun and C. Lele, *Fuzzy point BCK/BCI-algebras*, International Journal of Pure and Applied Mathematics **1** (2002), 33–39.
- [9] Y.B. Jun, *On n -fold fuzzy implicative/commutative ideals of BCK-algebras*, IJMMS, (2001), 419–424.
- [10] C. Lele, C. Wu, P. Weke, T. Mamadou and G.E. Njock, *Fuzzy ideals and weak ideals in BCK-algebras*, Scientiae Mathematicae Japonicae **54** (2001), 323–336.
- [11] C. Hoo, *Some fuzzy concepts of BCI, BCK and MV-algebras*, Int. J. Approx. Reason. **18** (2002), 177–189.
- [12] L.Z. Liu and K.T. Li, *Fuzzy filters of BL-algebras*, Information Sciences (In press).
- [13] Y. Huang and Z. Chen, *On ideals in BCK-algebra*, Math. Japonica **50** (1999), 211–226.
- [14] K. Iseki and S. Tanaka, *An introduction to the theory of BCK-algebra*, Math. Japonica **23** (1978), 935–942.

- [15] O.G. Xi, *Fuzzy BCK-algebras*, *Math. Japonica* **36** (1991), 935–942.
- [16] L.A. Zadeh, *Fuzzy sets*, *Inform. and Control* **8** (1965), 338–353.

Received September 2005

Revised January 2006