

## A COMMON APPROACH TO DIRECTOIDS WITH AN ANTITONE INVOLUTION AND $D$ -QUASIRINGS\*

IVAN CHAJDA AND MIROSLAV KOLAŘÍK

*Department of Algebra and Geometry*  
*Palacký University Olomouc*  
*Tomkova 40, 779 00 Olomouc, Czech Republic*

**e-mail:** chajda@inf.upol.cz

**e-mail:** kolarik@inf.upol.cz

### Abstract

We introduce the so-called  $DN$ -algebra whose axiomatic system is a common axiomatization of directoids with an antitone involution and the so-called  $D$ -quasiring. It generalizes the concept of Newman algebras (introduced by H. Dobbertin) for a common axiomatization of Boolean algebras and Boolean rings.

**Keywords:** directoid, antitone involution,  $D$ -quasiring,  $DN$ -algebra,  $a$ -mutation.

**2000 Mathematics Subject Classification:** 06A12, 06A06, 06E20.

By a *Newman algebra* (see [1]) is meant a (generally non-associative) semi-ring  $\mathcal{A} = (A; +, \cdot, ', 0, 1)$  with neutral elements 0 and 1 and complementation operation  $'$  (i.e.  $x \cdot x' = 0$  and  $x + x' = 1$  for all  $x \in A$ ). These algebras were introduced by M.H.A. Newman in 1941 when studying the relationship between a non-associative modification of Boolean rings with unit and Boolean algebras. For the associative modification of a Newman algebra, so-called  $\mathbf{N}$ -algebra, the simple axiomatic system is given in [4].

---

\*This work is supported by the Research and Development Council of the Czech Government via the project MSM6198959214.

One of the axioms is as follows

$$(N) \quad x + y = ((1 + 1)' \cdot (x \cdot y))' \cdot (x' \cdot y)'$$

H. Dobbertin shows in [4] that if  $1 + 1 = 1$  in (N) then the corresponding  $N$ -algebra is a Boolean algebra and if  $1 + 1 = 0$  then it is a Boolean ring.

Since Boolean algebras and Boolean rings with unit are in a one-to-one correspondence, there was a question if an algebra similar to a Boolean ring can be constructed to obtain a one-to-one correspondence between ortholattices and these algebras. It was solved in [5] where the concept of *Boolean quasiring* is introduced. Hence ortholattices and Boolean quasirings are in the same correspondence as Boolean algebras and Boolean rings with unit. The natural question arised if also the concept of  $N$ -algebra can be generalized to serve as a common axiomatization of ortholattices and Boolean quasirings. This was answered in [3] where the concept of a **QN**-algebra is introduced by several simple axioms including the axiom (N). The main result of [3] is that if  $1 + 1 = 1$  then the  $QN$ -algebra is an ortholattice and if  $1 + 1 = 0$  then it is a Boolean quasiring. Hence, the analogy is complete.

The concept of an ortholattice can be generalized when the underlying semilattice is substituted by a so-called directoid (see e.g. [6]).

For the readers convenience, we repeat the definition: A *directoid* is an algebra  $\mathcal{D} = (D; \sqcap)$  of type (2) satisfying the identities

$$(D1) \quad x \sqcap x = x;$$

$$(D2) \quad (x \sqcap y) \sqcap x = x \sqcap y;$$

$$(D3) \quad y \sqcap (x \sqcap y) = x \sqcap y;$$

$$(D4) \quad x \sqcap ((x \sqcap y) \sqcap z) = (x \sqcap y) \sqcap z.$$

The *induced order* of a directoid  $\mathcal{D} = (D; \sqcap)$  is defined by  $x \leq y$  if and only if  $x \sqcap y = x$ . With respect to  $\leq$ , the couple  $(D; \leq)$  is a downward directed set where for every  $x, y \in D$  the element  $x \sqcap y$  is a common lower bound of  $x, y$ .

Also conversely, if  $(D; \leq)$  is a downward directed ordered set and for each  $x, y \in D$  we choose a common lower bound  $d$  of  $x, y$  arbitrarily with the constraint that  $x \leq y$  implies  $d = x$ , then, putting  $x \sqcap y = d$ , the resulting algebra  $(D; \sqcap)$  is a directoid.

Let  $(D; \sqcap)$  be a directoid with a least element 0 and a greatest element 1. A mapping from  $D$  to  $D$  assigning  $x'$  to  $x$  is called an *antitone involution* if  $x'' = x$  and if  $x \leq y$  implies  $y' \leq x'$  with respect to the induced order. A bounded directoid with an antitone involution will be denoted by  $\mathcal{D} = (D; \sqcap, ', 0, 1)$ . The term operation  $\sqcup$  on  $D$  defined by  $x \sqcup y = (x' \sqcap y)'$  will be called an *assigned operation*, see e.g. [2] for its properties and further results.

Hence, bounded directoids with an antitone involution are in fact a generalization of ortholattices since the underlying semilattice  $(D; \wedge)$  of an ortholattice  $(D; \wedge, \vee, ', 0, 1)$  is a directoid and, due to De Morgan laws,  $x \vee y = (x' \wedge y')'$  is an assigned operation. The question how the induced ring-like structure looks like was completely solved in [2]:

By a **D-quasiring** is meant an algebra  $\mathcal{R} = (R; +, \cdot, 0, 1)$  of type  $(2, 2, 0, 0)$  satisfying the identities

- (Q1)  $(x \cdot y) \cdot x = x \cdot y$ ;
- (Q2)  $y \cdot (x \cdot y) = x \cdot y$ ;
- (Q3)  $x \cdot ((x \cdot y) \cdot z) = (x \cdot y) \cdot z$ ;
- (Q4)  $x \cdot 0 = 0$ ;
- (Q5)  $x \cdot 1 = x$ ;
- (Q6)  $x + 0 = x$ ;
- (Q7)  $1 + ((1 + (x \cdot y)) \cdot (1 + y)) = y$ .

The following correspondence is shown in [2] (Theorems 4 and 5):

**Proposition.** *If  $\mathcal{R} = (R; +, \cdot, 0, 1)$  is a D-quasiring and*

$$x \sqcap y = x \cdot y, \quad x' = 1 + x \quad \text{and} \quad x \sqcup y = 1 + ((1 + x) \cdot (1 + y))$$

*then  $\mathcal{D}(R) = (R; \sqcap, ', 0, 1)$  is a bounded directoid with an antitone involution where  $\sqcup$  is the assigned operation.*

*If  $\mathcal{D} = (D; \sqcap, ', 0, 1)$  is a bounded directoid with an antitone involution and  $\sqcup$  its assigned operation then for*

$$x + y = (x \sqcup y) \sqcap (x \sqcap y)' \quad \text{and} \quad x \cdot y = x \sqcap y$$

*the algebra  $\mathcal{R}(D) = (D; +, \cdot, 0, 1)$  is a D-quasiring.*

Due to an analogy of the relationship between ortholattices and Boolean quasirings, we can search for the analogy of a  $QN$ -algebra. A suitable candidate can be as follows.

**Definition 1.** By a **DN-algebra** we mean an algebra  $\mathcal{A} = (A; +, \cdot, ', 0, 1)$  of type  $(2, 2, 1, 0, 0)$  satisfying the following identities

- (1)  $(x \cdot y) \cdot x = x \cdot y$ ;
- (2)  $y \cdot (x \cdot y) = x \cdot y$ ;
- (3)  $x \cdot ((x \cdot y) \cdot z) = (x \cdot y) \cdot z$ ;
- (4)  $x \cdot 0 = 0 = 0 \cdot x$ ;
- (5)  $x \cdot 1 = x = 1 \cdot x$ ;
- (6)  $x'' = x$ ;
- (7)  $(x' \cdot y')' \cdot x = x$ ;
- (N)  $x + y = ((1 + 1)' \cdot (x \cdot y))' \cdot (x' \cdot y)'$ .

Let us note that not only (N) but also the axioms (4) and (6) are common with a Newman algebra. Moreover, every  $N$ -algebra and every  $QN$ -algebra is a  $DN$ -algebra as well.

We are going to show that the aforementioned algebras can be derived from  $DN$ -algebras in the same way as described before.

**Theorem 1.** *Let  $\mathcal{A} = (A; +, \cdot, ', 0, 1)$  be a  $DN$ -algebra satisfying  $1 + 1 = 1$ . Then its reduct  $\mathcal{D}(\mathcal{A}) = (A; \cdot, ', 0, 1)$  is a bounded directoid with an antitone involution where the assigned operation is  $+$ .*

**Proof.** Of course, (1) is (D2), (2) is (D3) and (3) is (D4). Putting  $y = z = 1$  in (3) and using (5) we obtain (D1). Hence,  $(A; \cdot)$  is a directoid. Let  $\leq$  be its induced order. By (4) and (5) we conclude  $0 \leq x \leq 1$  for any  $x \in A$ . By (6) we see that the mapping  $x \mapsto x'$  is an involution. Suppose  $y' \leq x'$ . Then  $x' \cdot y' = y'$  and, by (6) and (7),  $y \cdot x = (y')' \cdot x = (x' \cdot y')' \cdot x = x$  thus  $x \leq y$ . Due to (6), also  $x \leq y$  implies  $y' \leq x'$  thus this involution is antitone. Hence also  $0' = 1$  and  $1' = 0$ . Assume  $1 + 1 = 1$ . We compute

$$x + y = (1' \cdot (x \cdot y))' \cdot (x' \cdot y')' = 0' \cdot (x' \cdot y')' = (x' \cdot y')'$$

whence  $+$  is the assigned operation. ■

We show that also the second conclusion is analogous.

**Theorem 2.** *Let  $\mathcal{A} = (A; +, \cdot, ', 0, 1)$  be a  $DN$ -algebra. If  $1 + 1 = 0$  then its reduct  $\mathcal{R}(\mathcal{A}) = (A; +, \cdot, 0, 1)$  is a  $D$ -quasiring.*

**Proof.** Of course, (1) is (Q1), (2) is (Q2), (3) is (Q3), (4) implies (Q4) and (5) implies (Q5). By (6) and (7),  $x \mapsto x'$  is an antitone involution as shown in the proof of Theorem 1 (where the induced order is that of the directoid  $(A; \cdot)$ ). Hence  $0' = 1$  and  $1' = 0$ .

Then by (N) we have

$$x + 0 = ((1 + 1)' \cdot (x \cdot 0))' \cdot (x' \cdot 0')' = 0' \cdot (x' \cdot 0')' = (x' \cdot 1)' = x'' = x$$

proving (Q6).

Further, by (N) we conclude  $1 + x = x'$  and, applying (6),  $1 + (1 + x) = x$ . Since  $(A; \cdot)$  is a directoid, by (D3) we have  $y \cdot (x \cdot y) = x \cdot y$ , i.e.  $x \cdot y \leq y$ . Hence  $y' \leq (x \cdot y)'$  and  $(x \cdot y)' \cdot y' = (1 + (x \cdot y)) \cdot (1 + y) = 1 + y$ . Applying the previous identity, we conclude

$$1 + ((1 + (x \cdot y)) \cdot (1 + y)) = 1 + (1 + y) = y$$

which is (Q7). Hence  $\mathcal{R}(\mathcal{A}) = (A; +, \cdot, 0, 1)$  is a  $D$ -quasiring. ■

**Corollary 1.**  *$QN$ -algebras are exactly the  $DN$ -algebras satisfying  $x \cdot (y \cdot z) = (y \cdot x) \cdot z$  and  $x \cdot x' = 0$ .*

**Proof.** It is evident that the identity  $x \cdot (y \cdot z) = (y \cdot x) \cdot z$  yields commutativity and associativity of the operation " $\cdot$ ". Moreover, taking  $x'$  and  $y'$  instead of  $x$  and  $y$  in (7) and using (6) we obtain  $(x \cdot y)' \cdot x' = x'$ , i.e.  $((x \cdot y)' \cdot x')' = x$ . Hence, if the given  $DN$ -algebra satisfies also  $x \cdot x' = 0$  then it is a  $QN$ -algebra. ■

**Corollary 2.** *Ortholattices are exactly the  $DN$ -algebras satisfying  $x \cdot x' = 0$ ,  $x \cdot (y \cdot z) = (y \cdot x) \cdot z$  and  $1 + 1 = 1$ .*

**Proof.** As shown by Corollary 1, the corresponding  $DN$ -algebra is a  $QN$ -algebra. Hence, if it satisfies also  $1 + 1 = 1$ , it is an ortholattice by Theorem 2.3. in [3]. ■

**Definition 2.** Let  $\mathcal{A} = (A; +, \cdot, ', 0, 1)$  be a *DN-algebra* and  $a \in A$ . Define  $x +_a y = (a' \cdot (x \cdot y))' \cdot (x' \cdot y)'$ . Then  $\mathcal{A}_a = (A; +_a, \cdot, ', 0, 1)$  will be called the *a-mutation of  $\mathcal{A}$* .

An analogous concept was defined for *N-algebras* in [4] and for *QN-algebras* in [3].

**Theorem 3.** Let  $\mathcal{A} = (A; +, \cdot, ', 0, 1)$  be a *DN-algebra* and  $a, b \in A$ . Then the following hold:

- (i)  $1 +_a 1 = a$ ;
- (ii)  $\mathcal{A}_a$  is a *DN-algebra*;
- (iii)  $\mathcal{A}_1$  is a bounded directoid with an antitone involution;
- (iv)  $\mathcal{A}_0$  is a *D-quasiring*;
- (v)  $\mathcal{A}_{1+1} = \mathcal{A}$ ;
- (vi)  $(\mathcal{A}_a)_b = \mathcal{A}_b$ ;
- (vii)  $\{\mathcal{A}_a; a \in A\}$  is the set of all *DN-algebras* with base set  $A$  having the same multiplication and the same unary operation as  $\mathcal{A}$ ;
- (viii)  $\mathcal{A}$  and  $\mathcal{A}_a$  admit the same congruences.

**Proof.**

- (i)  $1 +_a 1 = (a' \cdot (1 \cdot 1))' \cdot (1' \cdot 1) = (a')' \cdot (1') = a \cdot 1 = a$ .
- (ii) By (i),  $x +_a y = (a' \cdot (x \cdot y))' \cdot (x' \cdot y) = ((1 +_a 1)' \cdot (x \cdot y))' \cdot (x' \cdot y)$  for all  $x, y \in A$ .
- (iii) According to (ii),  $\mathcal{A}_1$  is a *DN-algebra* and according to (i),  $1 +_1 1 = 1$ . Hence  $\mathcal{A}_1$  is a bounded directoid with an antitone involution where the assigned operation is  $+_1$  (by Theorem 1).
- (iv) According to (ii),  $\mathcal{A}_0$  is a *DN-algebra* and according to (i),  $1 +_0 1 = 0$  thus  $\mathcal{A}_0$  is a *D-quasiring* (by Theorem 2).
- (v)  $x +_{1+1} y = ((1 + 1)' \cdot (x \cdot y))' \cdot (x' \cdot y) = x + y$  for all  $x, y \in A$ .
- (vi) Since  $\mathcal{A}$  is a *DN-algebra*, the same is true for  $\mathcal{A}_a = (A; +_a, \cdot, ', 0, 1)$  according to (ii), and  $x +_a y = (a' \cdot (x \cdot y))' \cdot (x' \cdot y)$  for all  $x, y \in A$ .

Since  $\mathcal{A}_a$  is a *DN*-algebra, the same is true for  $(\mathcal{A}_a)_b = (A; (+_a)_b, \cdot', 0, 1)$  according to (ii) and  $x(+_a)_b y = (b' \cdot (x \cdot y))' \cdot (x' \cdot y')' = x +_b y$  for all  $x, y \in A$ .

- (vii) Let  $\mathcal{S} = (A; \oplus, \cdot', 0, 1)$  be a *DN*-algebra. Then  $x \oplus y = ((1 \oplus 1)' \cdot (x \cdot y))' \cdot (x' \cdot y')' = x +_{1 \oplus 1} y$  for all  $x, y \in A$  and hence  $\mathcal{S} = \mathcal{A}_{1 \oplus 1}$ .
- (viii)  $\mathcal{A}$  and  $\mathcal{A}_a$  admit the same congruences as  $(A; \cdot, ')$  because  $+$  and  $+_a$  are polynomials of the algebra  $(A; \cdot, ')$ .

■

## REFERENCES

- [1] G. Birkhoff, *Lattice Theory*, (3<sup>rd</sup> edition), Colloq. Publ. 25, Proc. Amer. Math. Soc., Providence, R. I., 1967.
- [2] I. Chajda and M. Kolařík, *Directoids with an antitone involution*, Comment. Math. Univ. Carolinae (CMUC) **48** (2007), 555–567.
- [3] I. Chajda and H. Länger, *A common generalization of ortholattices and Boolean quasirings*, Demonstratio Math. **15** (2007), 769–774.
- [4] H. Dobbertin, *Note on associative Newman algebras*, Algebra Universalis **9** (1979), 396–397.
- [5] D. Dorninger, H. Länger and M. Mączyński, *The logic induced by a system of homomorphisms and its various algebraic characterizations*, Demonstratio Math. **30** (1997), 215–232.
- [6] J. Ježek and R. Quackenbush, *Directoids: algebraic models of up-directed sets*, Algebra Universalis **27** (1990), 49–69.

Received 18 May 2007

Revised 10 July 2007