

A COMMON APPROACH TO DIRECTOIDS WITH AN ANTITONE INVOLUTION AND D -QUASIRINGS*

IVAN CHAJDA AND MIROSLAV KOLAŘÍK

Department of Algebra and Geometry
Palacký University Olomouc
Tomkova 40, 779 00 Olomouc, Czech Republic

e-mail: chajda@inf.upol.cz

e-mail: kolarik@inf.upol.cz

Abstract

We introduce the so-called DN -algebra whose axiomatic system is a common axiomatization of directoids with an antitone involution and the so-called D -quasiring. It generalizes the concept of Newman algebras (introduced by H. Dobbertin) for a common axiomatization of Boolean algebras and Boolean rings.

Keywords: directoid, antitone involution, D -quasiring, DN -algebra, a -mutation.

2000 Mathematics Subject Classification: 06A12, 06A06, 06E20.

By a *Newman algebra* (see [1]) is meant a (generally non-associative) semi-ring $\mathcal{A} = (A; +, \cdot, ', 0, 1)$ with neutral elements 0 and 1 and complementation operation $'$ (i.e. $x \cdot x' = 0$ and $x + x' = 1$ for all $x \in A$). These algebras were introduced by M.H.A. Newman in 1941 when studying the relationship between a non-associative modification of Boolean rings with unit and Boolean algebras. For the associative modification of a Newman algebra, so-called \mathbf{N} -algebra, the simple axiomatic system is given in [4].

*This work is supported by the Research and Development Council of the Czech Government via the project MSM6198959214.

One of the axioms is as follows

$$(N) \quad x + y = ((1 + 1)' \cdot (x \cdot y))' \cdot (x' \cdot y)'$$

H. Dobbertin shows in [4] that if $1 + 1 = 1$ in (N) then the corresponding N -algebra is a Boolean algebra and if $1 + 1 = 0$ then it is a Boolean ring.

Since Boolean algebras and Boolean rings with unit are in a one-to-one correspondence, there was a question if an algebra similar to a Boolean ring can be constructed to obtain a one-to-one correspondence between ortholattices and these algebras. It was solved in [5] where the concept of *Boolean quasiring* is introduced. Hence ortholattices and Boolean quasirings are in the same correspondence as Boolean algebras and Boolean rings with unit. The natural question arised if also the concept of N -algebra can be generalized to serve as a common axiomatization of ortholattices and Boolean quasirings. This was answered in [3] where the concept of a **QN**-algebra is introduced by several simple axioms including the axiom (N). The main result of [3] is that if $1 + 1 = 1$ then the QN -algebra is an ortholattice and if $1 + 1 = 0$ then it is a Boolean quasiring. Hence, the analogy is complete.

The concept of an ortholattice can be generalized when the underlying semilattice is substituted by a so-called directoid (see e.g. [6]).

For the readers convenience, we repeat the definition: A *directoid* is an algebra $\mathcal{D} = (D; \sqcap)$ of type (2) satisfying the identities

$$(D1) \quad x \sqcap x = x;$$

$$(D2) \quad (x \sqcap y) \sqcap x = x \sqcap y;$$

$$(D3) \quad y \sqcap (x \sqcap y) = x \sqcap y;$$

$$(D4) \quad x \sqcap ((x \sqcap y) \sqcap z) = (x \sqcap y) \sqcap z.$$

The *induced order* of a directoid $\mathcal{D} = (D; \sqcap)$ is defined by $x \leq y$ if and only if $x \sqcap y = x$. With respect to \leq , the couple $(D; \leq)$ is a downward directed set where for every $x, y \in D$ the element $x \sqcap y$ is a common lower bound of x, y .

Also conversely, if $(D; \leq)$ is a downward directed ordered set and for each $x, y \in D$ we choose a common lower bound d of x, y arbitrarily with the constraint that $x \leq y$ implies $d = x$, then, putting $x \sqcap y = d$, the resulting algebra $(D; \sqcap)$ is a directoid.

Let $(D; \sqcap)$ be a directoid with a least element 0 and a greatest element 1. A mapping from D to D assigning x' to x is called an *antitone involution* if $x'' = x$ and if $x \leq y$ implies $y' \leq x'$ with respect to the induced order. A bounded directoid with an antitone involution will be denoted by $\mathcal{D} = (D; \sqcap, ', 0, 1)$. The term operation \sqcup on D defined by $x \sqcup y = (x' \sqcap y')'$ will be called an *assigned operation*, see e.g. [2] for its properties and further results.

Hence, bounded directoids with an antitone involution are in fact a generalization of ortholattices since the underlying semilattice $(D; \wedge)$ of an ortholattice $(D; \wedge, \vee, ', 0, 1)$ is a directoid and, due to De Morgan laws, $x \vee y = (x' \wedge y')'$ is an assigned operation. The question how the induced ring-like structure looks like was completely solved in [2]:

By a **D-quasiring** is meant an algebra $\mathcal{R} = (R; +, \cdot, 0, 1)$ of type $(2, 2, 0, 0)$ satisfying the identities

- (Q1) $(x \cdot y) \cdot x = x \cdot y$;
- (Q2) $y \cdot (x \cdot y) = x \cdot y$;
- (Q3) $x \cdot ((x \cdot y) \cdot z) = (x \cdot y) \cdot z$;
- (Q4) $x \cdot 0 = 0$;
- (Q5) $x \cdot 1 = x$;
- (Q6) $x + 0 = x$;
- (Q7) $1 + ((1 + (x \cdot y)) \cdot (1 + y)) = y$.

The following correspondence is shown in [2] (Theorems 4 and 5):

Proposition. *If $\mathcal{R} = (R; +, \cdot, 0, 1)$ is a D-quasiring and*

$$x \sqcap y = x \cdot y, \quad x' = 1 + x \quad \text{and} \quad x \sqcup y = 1 + ((1 + x) \cdot (1 + y))$$

then $\mathcal{D}(R) = (R; \sqcap, ', 0, 1)$ is a bounded directoid with an antitone involution where \sqcup is the assigned operation.

If $\mathcal{D} = (D; \sqcap, ', 0, 1)$ is a bounded directoid with an antitone involution and \sqcup its assigned operation then for

$$x + y = (x \sqcup y) \sqcap (x \sqcap y)' \quad \text{and} \quad x \cdot y = x \sqcap y$$

the algebra $\mathcal{R}(D) = (D; +, \cdot, 0, 1)$ is a D-quasiring.

Due to an analogy of the relationship between ortholattices and Boolean quasirings, we can search for the analogy of a QN -algebra. A suitable candidate can be as follows.

Definition 1. By a **DN-algebra** we mean an algebra $\mathcal{A} = (A; +, \cdot, ', 0, 1)$ of type $(2, 2, 1, 0, 0)$ satisfying the following identities

- (1) $(x \cdot y) \cdot x = x \cdot y$;
- (2) $y \cdot (x \cdot y) = x \cdot y$;
- (3) $x \cdot ((x \cdot y) \cdot z) = (x \cdot y) \cdot z$;
- (4) $x \cdot 0 = 0 = 0 \cdot x$;
- (5) $x \cdot 1 = x = 1 \cdot x$;
- (6) $x'' = x$;
- (7) $(x' \cdot y')' \cdot x = x$;
- (N) $x + y = ((1 + 1)' \cdot (x \cdot y))' \cdot (x' \cdot y)'$.

Let us note that not only (N) but also the axioms (4) and (6) are common with a Newman algebra. Moreover, every N -algebra and every QN -algebra is a DN -algebra as well.

We are going to show that the aforementioned algebras can be derived from DN -algebras in the same way as described before.

Theorem 1. *Let $\mathcal{A} = (A; +, \cdot, ', 0, 1)$ be a DN -algebra satisfying $1 + 1 = 1$. Then its reduct $\mathcal{D}(\mathcal{A}) = (A; \cdot, ', 0, 1)$ is a bounded directoid with an antitone involution where the assigned operation is $+$.*

Proof. Of course, (1) is (D2), (2) is (D3) and (3) is (D4). Putting $y = z = 1$ in (3) and using (5) we obtain (D1). Hence, $(A; \cdot)$ is a directoid. Let \leq be its induced order. By (4) and (5) we conclude $0 \leq x \leq 1$ for any $x \in A$. By (6) we see that the mapping $x \mapsto x'$ is an involution. Suppose $y' \leq x'$. Then $x' \cdot y' = y'$ and, by (6) and (7), $y \cdot x = (y')' \cdot x = (x' \cdot y')' \cdot x = x$ thus $x \leq y$. Due to (6), also $x \leq y$ implies $y' \leq x'$ thus this involution is antitone. Hence also $0' = 1$ and $1' = 0$. Assume $1 + 1 = 1$. We compute

$$x + y = (1' \cdot (x \cdot y))' \cdot (x' \cdot y')' = 0' \cdot (x' \cdot y')' = (x' \cdot y')'$$

whence $+$ is the assigned operation. ■

We show that also the second conclusion is analogous.

Theorem 2. *Let $\mathcal{A} = (A; +, \cdot, ', 0, 1)$ be a DN -algebra. If $1 + 1 = 0$ then its reduct $\mathcal{R}(\mathcal{A}) = (A; +, \cdot, 0, 1)$ is a D -quasiring.*

Proof. Of course, (1) is (Q1), (2) is (Q2), (3) is (Q3), (4) implies (Q4) and (5) implies (Q5). By (6) and (7), $x \mapsto x'$ is an antitone involution as shown in the proof of Theorem 1 (where the induced order is that of the directoid $(A; \cdot)$). Hence $0' = 1$ and $1' = 0$.

Then by (N) we have

$$x + 0 = ((1 + 1)' \cdot (x \cdot 0))' \cdot (x' \cdot 0')' = 0' \cdot (x' \cdot 0')' = (x' \cdot 1)' = x'' = x$$

proving (Q6).

Further, by (N) we conclude $1 + x = x'$ and, applying (6), $1 + (1 + x) = x$. Since $(A; \cdot)$ is a directoid, by (D3) we have $y \cdot (x \cdot y) = x \cdot y$, i.e. $x \cdot y \leq y$. Hence $y' \leq (x \cdot y)'$ and $(x \cdot y)' \cdot y' = (1 + (x \cdot y)) \cdot (1 + y) = 1 + y$. Applying the previous identity, we conclude

$$1 + ((1 + (x \cdot y)) \cdot (1 + y)) = 1 + (1 + y) = y$$

which is (Q7). Hence $\mathcal{R}(\mathcal{A}) = (A; +, \cdot, 0, 1)$ is a D -quasiring. ■

Corollary 1. *QN -algebras are exactly the DN -algebras satisfying $x \cdot (y \cdot z) = (y \cdot x) \cdot z$ and $x \cdot x' = 0$.*

Proof. It is evident that the identity $x \cdot (y \cdot z) = (y \cdot x) \cdot z$ yields commutativity and associativity of the operation " \cdot ". Moreover, taking x' and y' instead of x and y in (7) and using (6) we obtain $(x \cdot y)' \cdot x' = x'$, i.e. $((x \cdot y)' \cdot x')' = x$. Hence, if the given DN -algebra satisfies also $x \cdot x' = 0$ then it is a QN -algebra. ■

Corollary 2. *Ortholattices are exactly the DN -algebras satisfying $x \cdot x' = 0$, $x \cdot (y \cdot z) = (y \cdot x) \cdot z$ and $1 + 1 = 1$.*

Proof. As shown by Corollary 1, the corresponding DN -algebra is a QN -algebra. Hence, if it satisfies also $1 + 1 = 1$, it is an ortholattice by Theorem 2.3. in [3]. ■

Definition 2. Let $\mathcal{A} = (A; +, \cdot, ', 0, 1)$ be a *DN-algebra* and $a \in A$. Define $x +_a y = (a' \cdot (x \cdot y))' \cdot (x' \cdot y)'$. Then $\mathcal{A}_a = (A; +_a, \cdot, ', 0, 1)$ will be called the *a-mutation of \mathcal{A}* .

An analogous concept was defined for *N-algebras* in [4] and for *QN-algebras* in [3].

Theorem 3. Let $\mathcal{A} = (A; +, \cdot, ', 0, 1)$ be a *DN-algebra* and $a, b \in A$. Then the following hold:

- (i) $1 +_a 1 = a$;
- (ii) \mathcal{A}_a is a *DN-algebra*;
- (iii) \mathcal{A}_1 is a bounded directoid with an antitone involution;
- (iv) \mathcal{A}_0 is a *D-quasiring*;
- (v) $\mathcal{A}_{1+1} = \mathcal{A}$;
- (vi) $(\mathcal{A}_a)_b = \mathcal{A}_b$;
- (vii) $\{\mathcal{A}_a; a \in A\}$ is the set of all *DN-algebras* with base set A having the same multiplication and the same unary operation as \mathcal{A} ;
- (viii) \mathcal{A} and \mathcal{A}_a admit the same congruences.

Proof.

- (i) $1 +_a 1 = (a' \cdot (1 \cdot 1))' \cdot (1' \cdot 1) = (a')' \cdot (1')' = a \cdot 1 = a$.
- (ii) By (i), $x +_a y = (a' \cdot (x \cdot y))' \cdot (x' \cdot y) = ((1 +_a 1)' \cdot (x \cdot y))' \cdot (x' \cdot y)$ for all $x, y \in A$.
- (iii) According to (ii), \mathcal{A}_1 is a *DN-algebra* and according to (i), $1 +_1 1 = 1$. Hence \mathcal{A}_1 is a bounded directoid with an antitone involution where the assigned operation is $+_1$ (by Theorem 1).
- (iv) According to (ii), \mathcal{A}_0 is a *DN-algebra* and according to (i), $1 +_0 1 = 0$ thus \mathcal{A}_0 is a *D-quasiring* (by Theorem 2).
- (v) $x +_{1+1} y = ((1 + 1)' \cdot (x \cdot y))' \cdot (x' \cdot y) = x + y$ for all $x, y \in A$.
- (vi) Since \mathcal{A} is a *DN-algebra*, the same is true for $\mathcal{A}_a = (A; +_a, \cdot, ', 0, 1)$ according to (ii), and $x +_a y = (a' \cdot (x \cdot y))' \cdot (x' \cdot y)$ for all $x, y \in A$.

Since \mathcal{A}_a is a *DN*-algebra, the same is true for $(\mathcal{A}_a)_b = (A; (+_a)_b, \cdot', 0, 1)$ according to (ii) and $x(+_a)_b y = (b' \cdot (x \cdot y))' \cdot (x' \cdot y')' = x +_b y$ for all $x, y \in A$.

- (vii) Let $\mathcal{S} = (A; \oplus, \cdot', 0, 1)$ be a *DN*-algebra. Then $x \oplus y = ((1 \oplus 1)' \cdot (x \cdot y))' \cdot (x' \cdot y')' = x +_{1 \oplus 1} y$ for all $x, y \in A$ and hence $\mathcal{S} = \mathcal{A}_{1 \oplus 1}$.
- (viii) \mathcal{A} and \mathcal{A}_a admit the same congruences as $(A; \cdot, ')$ because $+$ and $+_a$ are polynomials of the algebra $(A; \cdot, ')$. ■

REFERENCES

- [1] G. Birkhoff, *Lattice Theory*, (3rd edition), Colloq. Publ. 25, Proc. Amer. Math. Soc., Providence, R. I., 1967.
- [2] I. Chajda and M. Kolařík, *Directoids with an antitone involution*, Comment. Math. Univ. Carolinae (CMUC) **48** (2007), 555–567.
- [3] I. Chajda and H. Länger, *A common generalization of ortholattices and Boolean quasirings*, Demonstratio Math. **15** (2007), 769–774.
- [4] H. Dobbertin, *Note on associative Newman algebras*, Algebra Universalis **9** (1979), 396–397.
- [5] D. Dorninger, H. Länger and M. Mączyński, *The logic induced by a system of homomorphisms and its various algebraic characterizations*, Demonstratio Math. **30** (1997), 215–232.
- [6] J. Ježek and R. Quackenbush, *Directoids: algebraic models of up-directed sets*, Algebra Universalis **27** (1990), 49–69.

Received 18 May 2007

Revised 10 July 2007