

LEAPING CONVERGENTS OF TASOEV CONTINUED FRACTIONS

TAKAO KOMATSU*

Graduate School of Science and Technology
Hirosaki University, Hirosaki, 036-8561, Japan

e-mail: komatsu@cc.hirosaki-u.ac.jp

Abstract

Denote the n -th convergent of the continued fraction by $p_n/q_n = [a_0; a_1, \dots, a_n]$. We give some explicit forms of leaping convergents of Tasoev continued fractions. For instance, $[0; ua, ua^2, ua^3, \dots]$ is one of the typical types of Tasoev continued fractions. Leaping convergents are of the form p_{rn+i}/q_{rn+i} ($n = 0, 1, 2, \dots$) for fixed integers $r \geq 2$ and $0 \leq i \leq r - 1$.

Keywords: leaping convergents, Tasoev continued fractions.

2010 Mathematics Subject Classification: 05A19, 11A55, 11J70.

1. HURWITZ AND TASOEV CONTINUED FRACTIONS

Let $\alpha = [a_0; a_1, a_2, \dots]$ denote the regular (or simple) continued fraction expansion of a real number α , where

$$\begin{aligned} \alpha &= a_0 + 1/\alpha_1, & a_0 &= \lfloor \alpha \rfloor, \\ \alpha_n &= a_n + 1/\alpha_{n+1}, & a_n &= \lfloor \alpha_n \rfloor \quad (n \geq 1). \end{aligned}$$

*This research was supported in part by the Grant-in-Aid for Scientific research (C) (No. 22540005), the Japan Society for the Promotion of Science.

Hurwitz continued fraction expansions, quasi-periodic simple continued fractions, have the form

$$(1) \quad [a_0; a_1, \dots, a_n, \overline{Q_1(k), \dots, Q_p(k)}]_{k=1}^\infty \\ = [a_0; a_1, \dots, a_n, Q_1(1), \dots, Q_p(1), Q_1(2), \dots, Q_p(2), Q_1(3), \dots],$$

where a_0 is an integer, a_1, \dots, a_n are positive integers, Q_1, \dots, Q_p are polynomials with rational coefficients which take positive integral values for $k = 1, 2, \dots$ and at least one of the polynomials is not constant. Well-known examples are

$$e = [2; 1, 2, 1, 1, 4, 1, 1, 6, 1, \dots] = [2; \overline{1, 2k, 1}]_{k=1}^\infty, \\ \tanh 1 = \frac{e^2 - 1}{e^2 + 1} = [0; 1, 3, 5, 7, \dots] = [0; \overline{2k - 1}]_{k=1}^\infty, \\ \tan 1 = [1; 1, 1, 3, 1, 5, 1, \dots] = [1; \overline{2k - 1, 1}]_{k=1}^\infty.$$

It seems that every known example belongs to one of the three types, e -type, \tanh -type and \tan -type. No concrete example where the degree of any polynomial exceeds 1 has been given.

Recently, the author [5] found more general forms of Hurwitz continued fractions belonging to \tanh -type and \tan -type. Namely,

$$[0; ua, v(a + b), u(a + 2b), v(a + 3b), u(a + 4b), v(a + 5b), \dots] \\ (2) \quad = \frac{\sum_{n=0}^\infty (n!)^{-1} u^{-n-1} (vb)^{-n} \prod_{i=1}^n (a + bi)^{-1}}{\sum_{n=0}^\infty (n!)^{-1} (uvb)^{-n} \prod_{i=1}^{n-1} (a + bi)^{-1}}$$

and

$$[0; ua - 1, 1, v(a + b) - 2, 1, u(a + 2b) - 2, 1, v(a + 3b) - 2, 1, \dots] \\ (3) \quad = \frac{\sum_{n=0}^\infty (-1)^n (n!)^{-1} u^{-n-1} (vb)^{-n} \prod_{i=1}^n (a + bi)^{-1}}{\sum_{n=0}^\infty (-1)^n (n!)^{-1} (uvb)^{-n} \prod_{i=1}^{n-1} (a + bi)^{-1}},$$

respectively. In [9], the author constituted more general forms of Hurwitz continued fractions of e -type, namely, the quasi-periodic continued fractions with period 3 whose partial quotients include at least one 1;

$$(4) \quad \frac{[0; \overline{u(a+bk) - 1, 1, v - 1}]_{k=1}^{\infty}}{= \frac{\sum_{n=0}^{\infty} u^{-2n-1} v^{-2n} b^{-n} (n!)^{-1} \prod_{i=1}^{n+1} (a+bi)^{-1}}{\sum_{n=0}^{\infty} b^{-n} (n!)^{-1} ((uv)^{-2n} \prod_{i=1}^n (a+bi)^{-1} - (uv)^{-2n-1} \prod_{i=1}^{n+1} (a+bi)^{-1})}}$$

and

$$(5) \quad \frac{[0; \overline{v - 1, 1, u(a+bk) - 1}]_{k=1}^{\infty}}{= \frac{\sum_{n=0}^{\infty} b^{-n} (n!)^{-1} (u^{-2n} v^{-2n-1} \prod_{i=1}^n (a+bi)^{-1} + u^{-2n-1} v^{-2n-2} \prod_{i=1}^{n+1} (a+bi)^{-1})}{\sum_{n=0}^{\infty} (uv)^{-2n} b^{-n} (n!)^{-1} \prod_{i=1}^n (a+bi)^{-1}}}$$

Tasoev continued fractions ([4, 5, 8, 9, 10, 18]) are also systematic but have hardly been known before. They are also quasi-periodic but at least one of $Q_j(k)$'s in (1) includes exponentials in k instead of polynomials. In [5], the author found some more general Tasoev continued fractions. Namely,

$$(6) \quad [0; \overline{ua^k}]_{k=1}^{\infty} = \frac{\sum_{n=0}^{\infty} u^{-2n-1} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}{\sum_{n=0}^{\infty} u^{-2n} a^{-n^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}},$$

$$(7) \quad [0; \overline{ua - 1, 1, ua^{k+1} - 2}]_{k=1}^{\infty} = \frac{\sum_{n=0}^{\infty} (-1)^n u^{-2n-1} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}{\sum_{n=0}^{\infty} (-1)^n u^{-2n} a^{-n^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}},$$

$$(8) \quad [0; \overline{ua^k, va^k}]_{k=1}^{\infty} = \frac{\sum_{n=0}^{\infty} u^{-n-1} v^{-n} a^{-(n+1)(n+2)/2} \prod_{i=1}^n (a^i - 1)^{-1}}{\sum_{n=0}^{\infty} u^{-n} v^{-n} a^{-n(n+1)/2} \prod_{i=1}^n (a^i - 1)^{-1}}$$

and

$$(9) \quad \begin{aligned} & [0; ua - 1, 1, va - 2, 1, ua^{k+1} - 2, 1, va^{k+1} - 2]_{k=1}^{\infty} \\ &= \frac{\sum_{n=0}^{\infty} (-1)^n u^{-n-1} v^{-n} a^{-(n+1)(n+2)/2} \prod_{i=1}^n (a^i - 1)^{-1}}{\sum_{n=0}^{\infty} (-1)^n u^{-n} v^{-n} a^{-n(n+1)/2} \prod_{i=1}^n (a^i - 1)^{-1}}. \end{aligned}$$

We can safely say that Tasoev continued fractions are geometric and Hurwitz continued fractions are arithmetic ([8]). The Tasoev continued fractions corresponding to e -type Hurwitz continued fractions were also derived in [9];

$$(10) \quad \begin{aligned} & [0; ua^k - 1, 1, v - 1]_{k=1}^{\infty} \\ &= \frac{\sum_{n=0}^{\infty} u^{-2n-1} v^{-2n} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}{\sum_{n=0}^{\infty} ((uv)^{-2n} a^{-n^2} - (uv)^{-2n-1} a^{-(n+1)^2}) \prod_{i=1}^n (a^{2i} - 1)^{-1}} \end{aligned}$$

and

$$(11) \quad \begin{aligned} & [0; v - 1, 1, ua^k - 1]_{k=1}^{\infty} \\ &= \frac{\sum_{n=0}^{\infty} (u^{-2n} v^{-2n-1} a^{-n^2} + u^{-2n-1} v^{-2n-2} a^{-(n+1)^2}) \prod_{i=1}^n (a^{2i} - 1)^{-1}}{\sum_{n=0}^{\infty} (uv)^{-2n} a^{-n^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}. \end{aligned}$$

The different types of Tasoev continued fractions with period 3 shown in [8] are

$$(12) \quad \begin{aligned} & [0; ua^{2k-1} - 1, 1, va^{2k} - 1]_{k=1}^{\infty} \\ &= \frac{\sum_{n=0}^{\infty} u^{-n-1} v^{-n} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - (-1)^i)^{-1}}{\sum_{n=0}^{\infty} (-1)^n u^{-n} v^{-n} a^{-n^2} \prod_{i=1}^n (a^{2i} - (-1)^i)^{-1}}, \end{aligned}$$

$$\begin{aligned}
 & [0; \overline{ua, va^{2k} - 1, 1, ua^{2k+1} - 1}]_{k=1}^{\infty} \\
 (13) \quad &= \frac{\sum_{n=0}^{\infty} (-1)^n u^{-n-1} v^{-n} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - (-1)^i)^{-1}}{\sum_{n=0}^{\infty} u^{-n} v^{-n} a^{-n^2} \prod_{i=1}^n (a^{2i} - (-1)^i)^{-1}},
 \end{aligned}$$

$$\begin{aligned}
 & [0; \overline{ua^k - 1, 1, va^k - 1}]_{k=1}^{\infty} \\
 (14) \quad &= \frac{\sum_{n=0}^{\infty} u^{-n-1} v^{-n} a^{-(n+1)(n+2)/2} \prod_{i=1}^n (a^i - (-1)^i)^{-1}}{\sum_{n=0}^{\infty} (-1)^n u^{-n} v^{-n} a^{-n(n+1)/2} \prod_{i=1}^n (a^i - (-1)^i)^{-1}}
 \end{aligned}$$

and

$$\begin{aligned}
 & [0; \overline{ua, va^k - 1, 1, ua^{k+1} - 1}]_{k=1}^{\infty} \\
 (15) \quad &= \frac{\sum_{n=0}^{\infty} (-1)^n u^{-n-1} v^{-n} a^{-(n+1)(n+2)/2} \prod_{i=1}^n (a^i - (-1)^i)^{-1}}{\sum_{n=0}^{\infty} u^{-n} v^{-n} a^{-n(n+1)/2} \prod_{i=1}^n (a^i - (-1)^i)^{-1}}.
 \end{aligned}$$

2. LEAPING CONVERGENTS

Leaping convergents are every r -th convergent of the convergents $p_n/q_n = [a_0; a_1, a_2, \dots, a_n]$. Namely, Leaping convergents are of the form p_{rn+i}/q_{rn+i} ($n = 0, 1, 2, \dots$) for fixed integers $r \geq 2$ and $0 \leq i \leq r - 1$. Some explicit forms of the leaping convergents of Hurwitz continued fractions have been known. For example, if p_n/q_n and p_n^*/q_n^* are the n -th convergent of the continued fraction $e^{1/s} = [1; (2k-1)s, 1, 1]_{k=1}^{\infty}$ and $e = [2; \overline{1, 2k, 1}]_{k=1}^{\infty}$, respectively, then

$$\begin{aligned}
 p_{3n} &= p_{3n-2}^* = \sum_{k=0}^n \frac{(2n-k)!}{k!(n-k)!} s^{n-k}, \\
 q_{3n} &= q_{3n-2}^* = \sum_{k=0}^n (-1)^k \frac{(2n-k)!}{k!(n-k)!} s^{n-k},
 \end{aligned}$$

$$\begin{aligned}
p_{3n-1} &= p_{3n-3}^* = n \sum_{k=0}^n \frac{(2n-k-1)!}{k!(n-k)!} s^{n-k}, \\
q_{3n-1} &= q_{3n-3}^* = \sum_{k=0}^{n-1} (-1)^k \frac{(2n-k-1)!}{k!(n-k-1)!} s^{n-k}, \\
p_{3n-2} &= p_{3n-4}^* = \sum_{k=0}^{n-1} \frac{(2n-k-1)!}{k!(n-k-1)!} s^{n-k}, \\
q_{3n-2} &= q_{3n-4}^* = n \sum_{k=0}^n (-1)^k \frac{(2n-k-1)!}{k!(n-k)!} s^{n-k}.
\end{aligned}$$

(See e.g. [11, 12, 13, 14]). The study of leaping convergents of $e^{1/s}$ has been initiated by Elsner in the case of $s = 1$ ([1]) and by the author in the case of $s \geq 2$ ([6, 7]).

Such explicit forms are useful to lead varieties of applications. In fact, explicit forms of Hurwitz continued fractions have already yielded several interesting applications. In [2, 3] we give Diophantine approximations of values of hypergeometric functions formed from Diophantine equations. In [16, 17] we give several Diophantine approximations of \tanh , \tan , and some linear forms of e in terms of integrals. In [15] we give some new non-regular continued fraction expansions under the aspect of N continued fractions. These results are obtained by using some explicit forms of Hurwitz continued fractions.

In this paper, we shall give some explicit forms of leaping convergents of Tasoev continued fractions.

3. MAIN RESULTS

Let p_n/q_n be the n -th convergent of the Tasoev continued fraction

$$[0; \overline{ua^{2k-1}, va^{2k}}]_{k=1}^{\infty} = \frac{\sum_{n=0}^{\infty} u^{-n-1} v^{-n} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}{\sum_{n=0}^{\infty} u^{-n} v^{-n} a^{-n^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}.$$

Theorem 1. For $n \geq 0$,

$$p_n = \sum_{\nu=1}^{\lfloor \frac{n}{2} \rfloor} u^{\lfloor \frac{n}{2} \rfloor - \nu} v^{\lfloor \frac{n+1}{2} \rfloor - \nu} a^{\frac{n(n+1)}{2} - 2(\nu-1)n - \nu^2} \prod_{i=1}^{\nu-1} \frac{a^{2n} - a^{2(\nu+i-1)}}{a^{2i} - 1},$$

$$q_n = \sum_{\nu=1}^{\lfloor \frac{n}{2} \rfloor + 1} u^{\lfloor \frac{n}{2} \rfloor - \nu + 1} v^{\lfloor \frac{n+1}{2} \rfloor - \nu} a^{\frac{n(n+1)}{2} - 2(\nu-1)n - (\nu-1)^2} \prod_{i=1}^{\nu-1} \frac{a^{2n} - a^{2(\nu+i-2)}}{a^{2i} - 1}.$$

Remark 1. If $u = v$, then these identities become those of (6). If u is replaced by ua and a is replaced by $a^{1/2}$, then these become those of (8).

Proof. We shall prove the identities by induction. For simplicity, put

$$P(n) = \sum_{\nu=1}^{\lfloor \frac{n}{2} \rfloor} u^{\lfloor \frac{n}{2} \rfloor - \nu} v^{\lfloor \frac{n+1}{2} \rfloor - \nu} a^{\frac{n(n+1)}{2} - 2(\nu-1)n - \nu^2} \prod_{i=1}^{\nu-1} \frac{a^{2n} - a^{2(\nu+i-1)}}{a^{2i} - 1},$$

$$Q(n) = \sum_{\nu=1}^{\lfloor \frac{n}{2} \rfloor + 1} u^{\lfloor \frac{n}{2} \rfloor - \nu + 1} v^{\lfloor \frac{n+1}{2} \rfloor - \nu} a^{\frac{n(n+1)}{2} - 2(\nu-1)n - (\nu-1)^2} \prod_{i=1}^{\nu-1} \frac{a^{2n} - a^{2(\nu+i-2)}}{a^{2i} - 1}.$$

It is easy to see that $p_0 = 0 = P(0)$, $p_1 = 1 = P(1)$, $q(0) = 1 = Q(0)$ and $q(1) = ua = Q(1)$. Assume that $p_{2n-1} = P(2n-1)$ and $p_{2n-2} = P(2n-2)$. Then

$$\begin{aligned} p_{2n} &= va^{2n}p_{2n-1} + p_{2n-2} \\ &= va^{2n} \sum_{\nu=1}^n u^{n-\nu} v^{n-\nu} a^{n(2n-1) - 2(\nu-1)(2n-1) - \nu^2} \prod_{i=1}^{\nu-1} \frac{a^{4n-2} - a^{2(\nu+i-1)}}{a^{2i} - 1} \\ &\quad + \sum_{\nu=1}^{n-1} u^{n-\nu-1} v^{n-\nu} a^{(n-1)(2n-1) - 2(\nu-1)(2n-2) - \nu^2} \prod_{i=1}^{\nu-1} \frac{a^{4n-4} - a^{2(\nu+i-1)}}{a^{2i} - 1} \\ &= \sum_{\nu=1}^n u^{n-\nu} v^{n-\nu+1} a^{n(2n+1) - 4(\nu-1)n - \nu^2} a^{2(\nu-1)} \prod_{i=1}^{\nu-1} \frac{a^{4n-2} - a^{2(\nu+i-1)}}{a^{2i} - 1} \\ &\quad + \sum_{\nu=2}^n u^{n-\nu} v^{n-\nu+1} a^{(n-1)(2n-1) - 2(\nu-2)(2n-2) - (\nu-1)^2} \prod_{i=1}^{\nu-2} \frac{a^{4n-4} - a^{2(\nu+i-2)}}{a^{2i} - 1} \end{aligned}$$

$$\begin{aligned}
&= \sum_{\nu=1}^n u^{n-\nu} v^{n-\nu+1} a^{n(2n+1)-4(\nu-1)n-\nu^2} \left((a^{4n} - a^{2(2\nu-1)}) + a^{2\nu} (a^{2(\nu-1)} - 1) \right) \\
&\quad \times \frac{(a^{4n} - a^{2(\nu+1)})(a^{4n} - a^{2(\nu+2)}) \dots (a^{4n} - a^{2(2\nu-2)})}{(a^2 - 1)(a^4 - 1) \dots (a^{2(\nu-1)} - 1)} \\
&= \sum_{\nu=1}^n u^{n-\nu} v^{n-\nu+1} a^{n(2n+1)-4(\nu-1)n-\nu^2} \prod_{i=1}^{\nu-1} \frac{a^{4n} - a^{2(\nu+i-1)}}{a^{2i} - 1} \\
&= P(2n).
\end{aligned}$$

Next, assume that $p_{2n} = P(2n)$ and $p_{2n-1} = P(2n-1)$. Then

$$\begin{aligned}
p_{2n+1} &= ua^{2n+1}p_{2n} + p_{2n-1} \\
&= ua^{2n+1} \sum_{\nu=1}^n u^{n-\nu} v^{n-\nu+1} a^{n(2n+1)-4(\nu-1)n-\nu^2} \prod_{i=1}^{\nu-1} \frac{a^{4n} - a^{2(\nu+i-1)}}{a^{2i} - 1} \\
&\quad + \sum_{\nu=1}^n u^{n-\nu} v^{n-\nu} a^{n(2n-1)-2(\nu-1)(2n-1)-\nu^2} \prod_{i=1}^{\nu-1} \frac{a^{4n-2} - a^{2(\nu+i-1)}}{a^{2i} - 1} \\
&= \sum_{\nu=1}^n (uv)^{n-\nu+1} a^{(n+1)(2n+1)-2(\nu-1)(2n+1)-\nu^2} a^{2(\nu-1)} \prod_{i=1}^{\nu-1} \frac{a^{4n} - a^{2(\nu+i-1)}}{a^{2i} - 1} \\
&\quad + \sum_{\nu=2}^{n+1} (uv)^{n-\nu+1} a^{n(2n-1)-2(\nu-2)(2n-1)-(\nu-1)^2} \prod_{i=1}^{\nu-2} \frac{a^{2(2n-1)} - a^{2(\nu+i-2)}}{a^{2i} - 1} \\
&= \sum_{\nu=1}^{n+1} (uv)^{n-\nu+1} a^{(n+1)(2n+1)-2(\nu-1)(2n+1)-\nu^2} \\
&\quad \times \left((a^{4n+2} - a^{2(2\nu-1)}) + a^{2\nu} (a^{2(\nu-1)} - 1) \right) \frac{\prod_{i=1}^{\nu-2} (a^{4n+2} - a^{2(\nu+i)})}{\prod_{i=1}^{\nu-1} (a^{2i} - 1)} \\
&= \sum_{\nu=1}^{n+1} (uv)^{n-\nu+1} a^{(n+1)(2n+1)-2(\nu-1)(2n+1)-\nu^2} \prod_{i=1}^{\nu-1} \frac{a^{2(2n+1)} - a^{2(\nu+i-1)}}{a^{2i} - 1} \\
&= P(2n+1).
\end{aligned}$$

In similar manners, we can prove that if $q_{2n-1} = Q(2n-1)$ and $q_{2n-2} = Q(2n-2)$ then $q_{2n} = Q(2n)$, and if $q_{2n} = Q(2n)$ and $q_{2n-1} = Q(2n-1)$ then $q_{2n+1} = Q(2n+1)$. \blacksquare

We can rewrite the formulas as follows.

Corollary 1. For $n \geq 0$,

$$\begin{aligned} p_{2n} &= \sum_{\nu=0}^{n-1} u^\nu v^{\nu+1} a^{(\nu+2)(2\nu+1)} \prod_{i=1}^{n-\nu-1} \frac{a^{2(2\nu+i+1)} - 1}{a^{2i} - 1}, \\ p_{2n+1} &= \sum_{\nu=0}^n (uv)^\nu a^{\nu(2\nu+3)} \prod_{i=1}^{n-\nu} \frac{a^{2(2\nu+i)} - 1}{a^{2i} - 1}, \\ q_{2n} &= \sum_{\nu=0}^n (uv)^\nu a^{\nu(2\nu+1)} \prod_{i=1}^{n-\nu} \frac{a^{2(2\nu+i)} - 1}{a^{2i} - 1}, \\ q_{2n+1} &= \sum_{\nu=0}^n u^{\nu+1} v^\nu a^{(\nu+1)(2\nu+1)} \prod_{i=1}^{n-\nu} \frac{a^{2(2\nu+i+1)} - 1}{a^{2i} - 1}. \end{aligned}$$

Let p_n/q_n be the n -th convergent of the Tasoev continued fraction

$$\begin{aligned} &[0; ua - 1, 1, va^{2k} - 2, 1, ua^{2k+1} - 2]_{k=1}^\infty \\ &= \frac{\sum_{n=0}^\infty (-1)^n u^{-n-1} v^{-n} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}{\sum_{n=0}^\infty (-1)^n u^{-n} v^{-n} a^{-n^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}. \end{aligned}$$

For simplicity, put

$$\begin{aligned} P^*(n) &= \\ &= \sum_{\nu=1}^{\lfloor \frac{n}{2} \rfloor} (-1)^{\nu-1} u^{\lfloor \frac{n}{2} \rfloor - \nu} v^{\lfloor \frac{n+1}{2} \rfloor - \nu} a^{\frac{n(n+1)}{2} - 2(\nu-1)n - \nu^2} \prod_{i=1}^{\nu-1} \frac{a^{2n} - a^{2(\nu+i-1)}}{a^{2i} - 1}, \end{aligned}$$

$$\begin{aligned} Q^*(n) &= \\ &= \sum_{\nu=1}^{\lfloor \frac{n}{2} \rfloor + 1} (-1)^{\nu-1} u^{\lfloor \frac{n}{2} \rfloor - \nu + 1} v^{\lfloor \frac{n+1}{2} \rfloor - \nu} a^{\frac{n(n+1)}{2} - 2(\nu-1)n - (\nu-1)^2} \prod_{i=1}^{\nu-1} \frac{a^{2n} - a^{2(\nu+i-2)}}{a^{2i} - 1}. \end{aligned}$$

Theorem 2. For $n \geq 0$ we have

$$\begin{aligned} p_{2n} &= P^*(n), & p_{2n+1} &= P^*(n+1) - P^*(n), \\ q_{2n} &= Q^*(n), & q_{2n+1} &= Q^*(n+1) - Q^*(n). \end{aligned}$$

Remark 2. If $u = v$, then these identities become those of (7). If u is replaced by ua and a is replaced by $a^{1/2}$, then these become those of (9).

Proof. The proof is similar to the previous one and omitted. ■

We can rewrite the formulas as follows.

Corollary 2. For $n \geq 0$,

$$\begin{aligned} P^*(2n) &= \sum_{\nu=0}^{n-1} (-1)^{n-\nu+1} u^\nu v^{\nu+1} a^{(\nu+2)(2\nu+1)} \prod_{i=1}^{n-\nu-1} \frac{a^{2(2\nu+i+1)} - 1}{a^{2i} - 1}, \\ P^*(2n+1) &= \sum_{\nu=0}^n (-1)^{n-\nu} (uv)^\nu a^{\nu(2\nu+3)} \prod_{i=1}^{n-\nu} \frac{a^{2(2\nu+i)} - 1}{a^{2i} - 1}, \\ Q^*(2n) &= \sum_{\nu=0}^n (-1)^{n-\nu} (uv)^\nu a^{\nu(2\nu+1)} \prod_{i=1}^{n-\nu} \frac{a^{2(2\nu+i)} - 1}{a^{2i} - 1}, \\ Q^*(2n+1) &= \sum_{\nu=0}^n (-1)^{n-\nu} u^{\nu+1} v^\nu a^{(\nu+1)(2\nu+1)} \prod_{i=1}^{n-\nu} \frac{a^{2(2\nu+i+1)} - 1}{a^{2i} - 1}. \end{aligned}$$

4. EXPLICIT FORMS OF e -TYPE TASOEV CONTINUED FRACTIONS

In this section, we shall show some explicit forms of the leaping convergents of e -type Tasoev continued fractions. Proofs are similarly done by induction and omitted.

Let p_n/q_n be the n -th convergent of the Tasoev continued fraction (10):

$$\begin{aligned} & [0; \overline{ua^k - 1, 1, v - 1}]_{k=1}^\infty \\ &= \frac{\sum_{n=0}^\infty u^{-2n-1} v^{-2n} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}{\sum_{n=0}^\infty ((uv)^{-2n} a^{-n^2} - (uv)^{-2n-1} a^{-(n+1)^2}) \prod_{i=1}^n (a^{2i} - 1)^{-1}}. \end{aligned}$$

Theorem 3. For $n \geq 0$ we have

$$\begin{aligned}
p_{3n} &= \sum_{\nu=1}^{\lfloor \frac{n}{2} \rfloor} u^{n-2\nu+1} v^{n-2\nu+2} a^{\frac{n(n+1)}{2}-2(\nu-1)n-\nu^2} \prod_{i=1}^{\nu-1} \frac{a^{2n} - a^{2(\nu+i-1)}}{a^{2i} - 1}, \\
p_{3n+2} &= \sum_{\nu=1}^{n+1} (uv)^{n-\nu+1} a^{\frac{(n-\nu+2)(n-\nu+3)}{2}-1} \prod_{i=1}^{\lceil \nu/2 \rceil - 1} \frac{a^{2(n-\nu+i+1)} - 1}{a^{2i} - 1} \\
&= \sum_{\nu=1}^{n+1} (uv)^{n-\nu+1} a^{\frac{n(n+1)}{2} - (\nu-2)n - \lfloor \frac{\nu+1}{2} \rfloor^2 + \frac{1-(-1)^\nu}{2} \lfloor \frac{\nu-1}{2} \rfloor} \prod_{i=1}^{\lfloor \frac{\nu-1}{2} \rfloor} \frac{a^{2n} - a^{2(\lfloor \frac{\nu}{2} \rfloor + i - 1)}}{a^{2i} - 1}, \\
q_{3n} &= \sum_{\kappa=0}^{\lfloor \frac{n}{2} \rfloor} (uv)^{n-2\kappa} a^{\frac{n(n+1)}{2} - 2\kappa n - \kappa^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i-1)}}{a^{2i} - 1} \\
&\quad - \sum_{\kappa=0}^{\lfloor \frac{n-1}{2} \rfloor} (uv)^{n-2\kappa-1} a^{\frac{n(n+1)}{2} - 2\kappa n - (\kappa+1)^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i)}}{a^{2i} - 1} \\
&= \sum_{\nu=1}^{n+1} (-1)^{\nu-1} (uv)^{n-\nu+1} a^{\frac{n(n+1)}{2} - 2\lfloor \frac{\nu-1}{2} \rfloor n - \lfloor \frac{\nu}{2} \rfloor^2} \prod_{i=1}^{\lfloor \frac{\nu-1}{2} \rfloor} \frac{a^{2n} - a^{2(\lfloor \frac{\nu}{2} \rfloor + i - 1)}}{a^{2i} - 1}, \\
q_{3n+2} &= \sum_{\kappa=0}^{\lfloor \frac{n}{2} \rfloor} u^{n-2\kappa+1} v^{n-2\kappa} a^{\frac{(n+1)(n+2)}{2} - 2\kappa n - \kappa^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i-2)}}{a^{2i} - 1} \\
&\quad + \sum_{\kappa=0}^{\lfloor \frac{n-1}{2} \rfloor} u^{n-2\kappa} v^{n-2\kappa-1} a^{\frac{n(n+1)}{2} - 2\kappa n - \kappa^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i-1)}}{a^{2i} - 1} \\
&\quad - \sum_{\kappa=0}^{\lfloor \frac{n-1}{2} \rfloor} u^{n-2\kappa} v^{n-2\kappa-1} a^{\frac{(n+1)(n+2)}{2} - 2\kappa n - (\kappa+1)^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i-1)}}{a^{2i} - 1} \\
&\quad - \sum_{\kappa=0}^{\lfloor \frac{n-2}{2} \rfloor} u^{n-2\kappa-1} v^{n-2\kappa-2} a^{\frac{n(n+1)}{2} - 2\kappa n - (\kappa+1)^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i)}}{a^{2i} - 1}.
\end{aligned}$$

Remark 3. The forms of p_{3n+1} and q_{3n+1} are not simpler, and are obtained by using the recurrence relations: $p_{3n+1} = p_{3n+2} - p_{3n}$ and $q_{3n+1} = q_{3n+2} - q_{3n}$.

Let p_n/q_n be the n -th convergent of the Tasoev continued fraction (11):

$$\begin{aligned} & [0; \overline{v-1, 1, ua^k - 1}]_{k=1}^\infty \\ &= \frac{\sum_{n=0}^\infty (u^{-2n}v^{-2n-1}a^{-n^2} + u^{-2n-1}v^{-2n-2}a^{-(n+1)^2}) \prod_{i=1}^n (a^{2i} - 1)^{-1}}{\sum_{n=0}^\infty (uv)^{-2n}a^{-n^2} \prod_{i=1}^n (a^{2i} - 1)^{-1}}. \end{aligned}$$

Theorem 4. For $n \geq 0$ we have

$$\begin{aligned} p_{3n} &= \sum_{\kappa=0}^{\lfloor \frac{n-1}{2} \rfloor} u^{n-2\kappa}v^{n-2\kappa-1}a^{\frac{n(n+1)}{2}-2\kappa n-\kappa^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i-1)}}{a^{2i} - 1} \\ &+ \sum_{\kappa=0}^{\lfloor \frac{n-2}{2} \rfloor} u^{n-2\kappa-1}v^{n-2\kappa-2}a^{\frac{n(n+1)}{2}-2\kappa n-(\kappa+1)^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i)}}{a^{2i} - 1} \\ &- \sum_{\kappa=0}^{\lfloor \frac{n-2}{2} \rfloor} u^{n-2\kappa-1}v^{n-2\kappa-2}a^{\frac{n(n-1)}{2}-2\kappa n-\kappa^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i)}}{a^{2i} - 1} \\ &- \sum_{\kappa=0}^{\lfloor \frac{n-3}{2} \rfloor} u^{n-2\kappa-2}v^{n-2\kappa-3}a^{\frac{n(n-1)}{2}-2\kappa n-(\kappa+1)^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i+1)}}{a^{2i} - 1}, \\ p_{3n+2} &= \sum_{\kappa=0}^{\lfloor \frac{n}{2} \rfloor} (uv)^{n-2\kappa}a^{\frac{n(n+1)}{2}-2\kappa n-\kappa^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i-1)}}{a^{2i} - 1} \\ &+ \sum_{\kappa=0}^{\lfloor \frac{n-1}{2} \rfloor} (uv)^{n-2\kappa-1}a^{\frac{n(n+1)}{2}-2\kappa n-(\kappa+1)^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i)}}{a^{2i} - 1} \\ &= \sum_{\nu=1}^{n+1} (uv)^{n-\nu+1}a^{\frac{n(n+1)}{2}-2\lfloor \frac{\nu-1}{2} \rfloor n - \lfloor \frac{\nu}{2} \rfloor^2} \prod_{i=1}^{\lfloor \frac{\nu-1}{2} \rfloor} \frac{a^{2n} - a^{2(\lfloor \frac{\nu}{2} \rfloor + i - 1)}}{a^{2i} - 1}, \\ q_{3n} &= \sum_{\kappa=0}^{\lfloor \frac{n}{2} \rfloor} (uv)^{n-2\kappa}a^{\frac{n(n+1)}{2}-2\kappa n-\kappa^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i-1)}}{a^{2i} - 1} \end{aligned}$$

$$\begin{aligned}
& - \sum_{\kappa=0}^{\lfloor \frac{n-1}{2} \rfloor} (uv)^{n-2\kappa-1} a^{\frac{n(n-1)}{2}-2\kappa n-\kappa^2} \prod_{i=1}^{\kappa} \frac{a^{2n} - a^{2(\kappa+i)}}{a^{2i} - 1} \\
& = \sum_{\nu=1}^{n+1} (-1)^{\nu-1} (uv)^{n-\nu+1} a^{\frac{n(n+1)}{2}-(\nu-1)n-\lfloor \frac{\nu-1}{2} \rfloor^2} \prod_{i=1}^{\lfloor \frac{\nu-1}{2} \rfloor} \frac{a^{2n} - a^{2(\lfloor \frac{\nu}{2} \rfloor+i-1)}}{a^{2i} - 1}, \\
q_{3n+2} & = \sum_{\nu=1}^{\lfloor \frac{n+1}{2} \rfloor} u^{n-2\nu+2} v^{n-2\nu+3} a^{\frac{n(n+5)}{2}-2\nu n-(\nu-1)^2} \prod_{i=1}^{\nu-1} \frac{a^{2n} - a^{2(\nu+i-2)}}{a^{2i} - 1}.
\end{aligned}$$

Remark 4. The forms of p_{3n+1} and q_{3n+1} are not simpler, and are obtained by using the recurrence relations: $p_{3n+1} = p_{3n+2} - p_{3n}$ and $q_{3n+1} = q_{3n+2} - q_{3n}$.

Let p_n/q_n be the n -th convergent of the Tasoev continued fraction (12):

$$[0; \overline{ua^{2k-1} - 1, 1, va^{2k} - 1}]_{k=1}^{\infty} = \frac{\sum_{n=0}^{\infty} u^{-n-1} v^{-n} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - (-1)^i)^{-1}}{\sum_{n=0}^{\infty} (-1)^n u^{-n} v^{-n} a^{-n^2} \prod_{i=1}^n (a^{2i} - (-1)^i)^{-1}}.$$

Theorem 5. For $n \geq 0$ we have

$$\begin{aligned}
p_{3n} & = \\
& = \sum_{\nu=1}^n u^{n-\nu} v^{n-\nu+1} a^{n(2n+1)-4(\nu-1)n-\nu^2} \prod_{i=1}^{\nu-1} \frac{a^{4n} - (-1)^{\nu+i-1} a^{2(\nu+i-1)}}{a^{2i} - (-1)^i}, \\
p_{3n+2} & = \\
& = \sum_{\nu=1}^{n+1} (uv)^{n-\nu+1} a^{(2n+1)(n+1)-2(\nu-1)(2n+1)-\nu^2} \prod_{i=1}^{\nu-1} \frac{a^{4n+2} - (-1)^{\nu+i} a^{2(\nu+i-1)}}{a^{2i} - (-1)^i}, \\
q_{3n} & = \\
& = \sum_{\nu=1}^{n+1} (-1)^{\nu-1} (uv)^{n-\nu+1} a^{n(2n+1)-4(\nu-1)n-(\nu-1)^2} \prod_{i=1}^{\nu-1} \frac{a^{4n} - (-1)^{\nu+i} a^{2(\nu+i-2)}}{a^{2i} - (-1)^i},
\end{aligned}$$

$$q_{3n+2} = \sum_{\nu=1}^{n+1} (-1)^{\nu-1} u^{n-\nu+2} v^{n-\nu+1} a^{(2n+1)(n+1)-2(\nu-1)(2n+1)-(\nu-1)^2} \\ \times \prod_{i=1}^{\nu-1} \frac{a^{4n+2} - (-1)^{\nu+i-1} a^{2(\nu+i-2)}}{a^{2i} - (-1)^i}.$$

Remark 5. The forms of p_{3n+1} and q_{3n+1} are not simpler, and are obtained by using the recurrence relations: $p_{3n+1} = p_{3n+2} - p_{3n}$ and $q_{3n+1} = q_{3n+2} - q_{3n}$.

If u is replaced by ua and a is replaced by $a^{1/2}$, then these become those of (14).

Let p_n/q_n be the n -th convergent of the Tasoev continued fraction (13):

$$[0; ua, \overline{va^{2k} - 1, 1, ua^{2k+1} - 1}]_{k=1}^{\infty} \\ = \frac{\sum_{n=0}^{\infty} (-1)^n u^{-n-1} v^{-n} a^{-(n+1)^2} \prod_{i=1}^n (a^{2i} - (-1)^i)^{-1}}{\sum_{n=0}^{\infty} u^{-n} v^{-n} a^{-n^2} \prod_{i=1}^n (a^{2i} - (-1)^i)^{-1}}.$$

Theorem 6. For $n \geq 0$ we have

$$p_{3n} = \sum_{\nu=1}^n (-1)^{\nu-1} u^{n-\nu} v^{n-\nu+1} a^{n(2n+1)-4(\nu-1)n-\nu^2} \\ \times \prod_{i=1}^{\nu-1} \frac{a^{4n} - (-1)^{\nu+i-1} a^{2(\nu+i-1)}}{a^{2i} - (-1)^i}, \\ p_{3n+1} = \sum_{\nu=1}^{n+1} (-1)^{\nu-1} (uv)^{n-\nu+1} a^{(2n+1)(n+1)-2(\nu-1)(2n+1)-\nu^2} \\ \times \prod_{i=1}^{\nu-1} \frac{a^{4n+2} - (-1)^{\nu+i} a^{2(\nu+i-1)}}{a^{2i} - (-1)^i}, \\ q_{3n} = \sum_{\nu=1}^{n+1} (uv)^{n-\nu+1} a^{n(2n+1)-4(\nu-1)n-(\nu-1)^2} \\ \times \prod_{i=1}^{\nu-1} \frac{a^{4n} - (-1)^{\nu+i} a^{2(\nu+i-2)}}{a^{2i} - (-1)^i},$$

$$q_{3n+1} = \sum_{\nu=1}^{n+1} u^{n-\nu+2} v^{n-\nu+1} a^{(2n+1)(n+1)-2(\nu-1)(2n+1)-(\nu-1)^2} \\ \times \prod_{i=1}^{\nu-1} \frac{a^{4n+2} - (-1)^{\nu+i-1} a^{2(\nu+i-2)}}{a^{2i} - (-1)^i}.$$

Remark 6. The forms of p_{3n+2} and q_{3n+2} are not simpler, and are obtained by using the recurrence relations: $p_{3n+2} = p_{3n+3} - p_{3n+1}$ and $q_{3n+2} = q_{3n+3} - q_{3n+1}$.

If u is replaced by ua and a is replaced by $a^{1/2}$, then these become those of (15).

REFERENCES

- [1] C. Elsner, *On arithmetic properties of the convergents of Euler's number*, Colloq. Math. **79** (1999), 133–145.
- [2] C. Elsner, T. Komatsu and I. Shiokawa, *Approximation of values of hypergeometric functions by restricted rationals*, J. Théor. Nombres Bordeaux **19** (2007), 393–404. doi:10.5802/jtnb.593
- [3] C. Elsner, T. Komatsu and I. Shiokawa, *On convergents formed from Diophantine equations*, Glasnik Mat. **44** (2009), 267–284. doi:10.3336/gm.44.2.02
- [4] T. Komatsu, *On Tasoev's continued fractions*, Math. Proc. Cambridge Philos. Soc. **134** (2003), 1–12. doi:10.1017/S0305004102006266
- [5] T. Komatsu, *On Hurwitzian and Tasoev's continued fractions*, Acta Arith. **107** (2003), 161–177. doi:10.4064/aa107-2-4
- [6] T. Komatsu, *Recurrence relations of the leaping convergents*, JP J. Algebra Number Theory Appl. **3** (2003), 447–459.
- [7] T. Komatsu, *Arithmetical properties of the leaping convergents of $e^{1/s}$* , Tokyo J. Math. **27** (2004), 1–12. doi:10.3836/tjm/1244208469
- [8] T. Komatsu, *Tasoev's continued fractions and Rogers-Ramanujan continued fractions*, J. Number Theory **109** (2004), 27–40. doi:10.1016/j.jnt.2004.06.001
- [9] T. Komatsu, *Hurwitz and Tasoev continued fractions*, Monatsh. Math. **145** (2005), 47–60. doi:10.1007/s00605-004-0281-0
- [10] T. Komatsu, *An algorithm of infinite sums representations and Tasoev continued fractions*, Math. Comp. **74** (2005), 2081–2094. doi:10.1090/S0025-5718-05-01752-7

- [11] T. Komatsu, *Some combinatorial properties of the leaping convergents*, in: Combinatorial Number Theory, Proceedings of the Integers Conference 2005 in Celebration of the 70th Birthday of Ronald Graham, Carrollton, Georgia, USA, October 27–30, 2005, eds. by B.M. Landman, M.B. Nathanson, J. Nešetřil, R.J. Nowakowski and C. Pomerance, Walter de Gruyter, 2007, pp. 315–325.
- [12] T. Komatsu, *Some combinatorial properties of the leaping convergents, II*, Applications of Fibonacci Numbers, Proceedings of 12th International Conference on Fibonacci Numbers and their Applications, Congr. Numer. **200** (2010), 187–196.
- [13] T. Komatsu, *Hurwitz continued fractions with confluent hypergeometric functions*, Czech. Math. J. **57** (2007), 919–932. doi:10.1007/s10587-007-0085-1
- [14] T. Komatsu, *Leaping convergents of Hurwitz continued fractions*, in: Diophantine Analysis and related fields (DARF 2007/2008), AIP Conf. Proc. 976, pp. 130–143. Amer. Inst. Phys., Melville, NY, 2008.
- [15] T. Komatsu, *Shrinking the period length of quasi-periodic continued fractions*, J. Number Theory **129** (2009), 358–366. doi:10.1016/j.jnt.2008.08.004
- [16] T. Komatsu, *A diophantine approximation of $e^{1/s}$ in terms of integrals*, Tokyo J. Math. **32** (2009), 159–176. doi:10.3836/tjm/1249648415
- [17] T. Komatsu, *Diophantine approximations of \tanh , \tan , and linear forms of e in terms of integrals*, Rev. Roum. Math. Pures Appl. **54** (2009), 223–242.
- [18] B.G. Tasoiev, *Rational approximations to certain numbers* (Russian), Mat. Zametki **67** (2000), 931–937; English transl. in Math. Notes **67** (2000), 786–791.

Received 3 March 2011

Revised 12 May 2011