

SOME CHARACTERIZATIONS OF 2-PRIMAL IDEALS OF A Γ -SEMIRING

SUHRID DHARA

Department of Pure Mathematics
University of Calcutta
35, Ballygunge Circular Road
Kolkata – 700019, India

e-mail: suhridd@yahoo.com

AND

TAPAN KUMAR DUTTA

Department of Pure Mathematics
University of Calcutta
35, Ballygunge Circular Road
Kolkata – 700019, India

e-mail: duttatapankumar@yahoo.co.in

Abstract

This paper is a continuation of our previous paper entitled “On 2-primal Γ -semirings”. In this paper we have introduced the notion of 2-primal ideal in Γ -semiring and studied it.

Keywords: Γ -semiring, nilpotent element, 2-primal Γ -semiring, 2-primal ideal, IFP (insertion of factor property), completely prime ideal, completely semiprime ideal.

2010 Mathematics Subject Classification: 16Y60, 16Y99.

1. INTRODUCTION

The notion of Γ -ring was introduced by N. Nobuswa [6] in 1964. Later W.E. Barnes [19] weakened the defining conditions of a Γ -ring. The notion of Γ -semiring was introduced by M.M.K. Rao in [4, 5]. Now-a-days there has been a remarkable growth of the theory of Γ -ring as well as of Γ -semiring.

Birkenmeier-Heatherly-Lee [3] introduced the notion of 2-primal ring in 1993. A ring R with identity is called 2-primal if $\mathcal{P}(R) = \mathcal{N}(R)$, where $\mathcal{P}(R)$ denotes the intersection of all prime ideals of R and $\mathcal{N}(R)$ denotes the set of all nilpotent elements of R . An ideal I of R is called 2-primal if $\mathcal{P}(R/I) = \mathcal{N}(R/I)$. Birkenmeier-Heatherly-Lee obtained some characterizations of 2-primal ideal in ring. They proved that an ideal I is 2-primal if and only if $\mathcal{P}(I)$ is a completely semiprime ideal of R .

In this paper we introduce the notion of 2-primal ideal in a Γ -semiring. We obtain some characterizations of 2-primal ideal in a Γ -semiring. Also we introduce the notion of $N_I(P)$ and N_I^P etc. in Γ -semiring and using them we obtain some characterizations of 2-primal ideals.

2. PRELIMINARIES

We first give the definition of a Γ -semiring.

Definition (See [12]). Let S and Γ be two additive commutative semigroups. Then S is called a Γ -semiring if there exists a mapping $S \times \Gamma \times S \rightarrow S$ (the image to be denoted by $a\alpha b$, for $a, b \in S$ and $\alpha \in \Gamma$) satisfying the following conditions:

- (i) $a\alpha(b + c) = a\alpha b + a\alpha c$
- (ii) $(a + b)\alpha c = a\alpha c + b\alpha c$
- (iii) $a(\alpha + \beta)c = a\alpha c + a\beta c$
- (iv) $a\alpha(b\beta c) = (a\alpha b)\beta c$ for all $a, b, c \in S$ and for all $\alpha, \beta \in \Gamma$.

Let S be a Γ -semiring. If there exists an element $0 \in S$ such that $0 + x = x$ and $0\alpha x = x\alpha 0 = 0$ for all $x \in S$ and for all $\alpha \in \Gamma$ then '0' is called the zero element or simply the zero of the Γ -semiring S . In this case we say that S is a Γ -semiring with zero.

Throughout this paper we assume that a Γ -semiring always contains a zero element and S^* denotes the set of all nonzero elements of S i.e., $S^* = S \setminus \{0\}$.

Definition (See [12]). Let S be a Γ -semiring and F be the free additive commutative semigroup generated by $S \times \Gamma$. Then the relation ρ on F defined by $\sum_{i=1}^m (x_i, \alpha_i)\rho \sum_{j=1}^n (y_j, \beta_j)$ if and only if $\sum_{i=1}^m x_i\alpha_i s = \sum_{j=1}^n y_j\beta_j s$ for all $s \in S$ ($m, n \in \mathbb{Z}^+$, the set of all positive integers), is a congruence on F . We denote the congruence class containing $\sum_{i=1}^m (x_i, \alpha_i)$ by $\sum_{i=1}^m [x_i, \alpha_i]$. Then F/ρ is an additive commutative semigroup. Now F/ρ forms a semiring with the multiplication defined by $(\sum_{i=1}^m [x_i, \alpha_i])(\sum_{j=1}^n [y_j, \beta_j]) = \sum_{i,j} [x_i\alpha_i y_j, \beta_j]$. We denote this semiring by L and call it the left operator semiring of the Γ -semiring S .

Dually, we define the right operator semiring R of the Γ -semiring S where $R = \{\sum_{i=1}^m [\alpha_i, x_i] : \alpha_i \in \Gamma, x_i \in S, i = 1, 2, \dots, m; m \in \mathbb{Z}^+\}$ and the multiplication on R is defined as $(\sum_{i=1}^m [\alpha_i, x_i])(\sum_{j=1}^n [\beta_j, y_j]) = \sum_{i,j} [\alpha_i, x_i \beta_j y_j]$.

Let S be a Γ -semiring and L be the left operator semiring and R be the right operator semiring of S . If there exists an element $\sum_{i=1}^m [e_i, \delta_i] \in L$ (respectively $\sum_{j=1}^n [\nu_j, f_j] \in R$) such that $\sum_{i=1}^m e_i \delta_i a = a$ (respectively $\sum_{j=1}^n a \nu_j f_j = a$) for all $a \in S$ then S is said to have the *left unity* $\sum_{i=1}^m [e_i, \delta_i]$ (respectively the *right unity* $\sum_{j=1}^n [\nu_j, f_j]$).

Definition (See [7]). If R is a commutative semiring and $R - \{0\}$ is a multiplicative group then R is called a Γ -semifield.

Definition (See [16]). Let A be a nonempty subset of a Γ -semiring S . The right annihilator of A with respect to $\Phi \subseteq \Gamma$ in S , denoted by $r(A, \Phi)$, is defined by $r(A, \Phi) = \{s \in S : A\Phi s = \{0\}\}$.

In particular, if $\Phi = \Gamma$ we denote $r(A, \Phi)$ by $ann_R(A)$. Again if $A = \{a\}$, then we denote $ann_R(A)$ by $ann_R(a)$.

Analogously we can define left annihilator $l(\Phi, A)$ and for $\Phi = \Gamma$ it is denoted by $ann_L(A)$.

Proposition 1 (See [16]). *The right annihilator $r(A, \Phi)$ of A with respect to Φ in a Γ -semiring S is a right ideal of S .*

Remark 2. Similar result holds for left annihilator.

For other preliminaries we refer to [17].

Throughout this paper we assume that a Γ -semiring S always contain a unity whose every ideal is a k -ideal.

3. 2-PRIMAL IDEALS

We begin with the following examples of Γ -semiring in which every ideal is a k -ideal.

Example 3. Let M be a Γ -ring with unity. Then M is a Γ -semiring with unity and every ideal of M is a k -ideal.

Example 4. Let R be a Γ -ring with unity, $S = \{r\omega : r \in \mathbb{R}_0^+\}$ and $\Gamma_1 = \{r\omega^2 : r \in \mathbb{R}_0^+\}$, where ω be a cube root of unity and \mathbb{R}_0^+ is the set of all non negative real numbers. Then S is a Γ_1 -semiring with unity with usual addition and multiplication. Also $R \times S$ is a $\Gamma \times \Gamma_1$ -semiring with unity which is not a $\Gamma \times \Gamma_1$ -ring but every ideal of $R \times S$ is a k -ideal.

Example 5. Let L be a bounded distributive lattice with maximal element 1. Then L is a Γ -semiring with unity, where $\Gamma = L$. Now L is not a Γ -ring. Also every ideal of L is a k -ideal.

Now we recall the following definitions:

Definition (See [13]). An element a of a Γ -semiring S is said to be *nilpotent* if for any $\gamma \in \Gamma$ there exists a positive integer $n = n(\gamma, a)$ such that $(a\gamma)^{n-1}a = 0$ and an element a of a Γ -semiring S is said to be *strongly nilpotent* if there exists a positive integer n such that $(a\Gamma)^{n-1}a = 0$.

Definition (See [17]). A Γ -semiring S is said to be a *2-primal Γ -semiring* if $\mathcal{P}(S) = \mathcal{N}(S)$, where $\mathcal{P}(S)$ denotes the intersection of all prime ideals of the Γ -semiring S i.e., the prime radical of S and $\mathcal{N}(S)$ denotes the set of all nilpotent elements of S .

Definition (See [17]). A one sided ideal I of a Γ -semiring S is said to have the *insertion of factors property* or simply IFP if for any $a, b \in S$, $a\Gamma b \subseteq I$ implies $a\Gamma S\Gamma b \subseteq I$.

Definition (See [17]). For a prime ideal P of a Γ -semiring S , we define
 $N(P) = \{x \in S : x\Gamma S\Gamma y \subseteq \mathcal{P}(S) \text{ for some } y \in S \setminus P\}$,
 $N_P = \{x \in S : x\Gamma y \subseteq \mathcal{P}(S) \text{ for some } y \in S \setminus P\}$,
 $\overline{N}_P = \{x \in S : (x\Gamma)^{n-1}x \subseteq N_P, \text{ for some positive integer } n\}$.

Definition. Let S be a Γ -semiring and I be an ideal of S . Then I is said to be a *2-primal ideal* of S if S/I is a 2-primal Γ -semiring i.e. if $\mathcal{P}(S/I) = \mathcal{N}(S/I)$, where $\mathcal{P}(S/I)$ denotes the intersection of all prime ideals of the factor Γ -semiring S/I and $\mathcal{N}(S/I)$ denotes the set of all nilpotent elements of S/I .

Definition. Let I be any ideal of a Γ -semiring S and P be a prime ideal of S . Then we define

$N_I(P) = \{x \in S : x\Gamma S\Gamma y \subseteq \mathcal{P}(I) \text{ for some } y \in S \setminus P\}$,
 $N_I^P = \{x \in S : x\Gamma y \subseteq \mathcal{P}(I) \text{ for some } y \in S \setminus P\}$,
 $\overline{N}_I(P) = \{x \in S : (x\Gamma)^{n-1}x \subseteq N_I(P), \text{ for some positive integer } n\}$,
 $\overline{N}_I^P = \{x \in S : (x\Gamma)^{n-1}x \subseteq N_I^P, \text{ for some positive integer } n\}$.

Example 6. Let F be a semifield. Consider the sets:

$$S = \left\{ \left(\begin{array}{cc} d_1 & d_2 \\ 0 & d_3 \\ 0 & 0 \end{array} \right) : d_1, d_2, d_3 \in F \right\}, \Gamma = \left\{ \left(\begin{array}{ccc} d_4 & d_5 & d_6 \\ 0 & d_7 & d_8 \end{array} \right) : d_4, d_5, d_6, d_7, d_8 \in F \right\}$$

and $I = \left\{ \begin{pmatrix} 0 & d \\ 0 & 0 \\ 0 & 0 \end{pmatrix} : d \in F \right\}$. Then S is a 2-primal Γ -semiring with respect to the usual matrix addition and usual matrix multiplication and I is a 2-primal ideal of S .

Proposition 7. *Let S be a Γ -semiring and I be an ideal of S . Then for any prime ideal P we have, $N(P) \subseteq N_I(P)$, $N_P \subseteq N_I^P$, $I \subseteq N_I(P) \subseteq \overline{N_I(P)}$ and $I \subseteq N_I^P \subseteq \overline{N_I^P}$.*

Definition (See [17]). A Γ -semiring S is said to satisfy (SI) if for each $a \in S$, $ann_R(a)$ is an ideal of S .

Definition (See [17]). A Γ -semiring S is said to be *SN Γ -semiring* if $\mathcal{N}(S) = \mathcal{N}_\Gamma(S)$, where $\mathcal{N}_\Gamma(S)$ is the set of all strongly nilpotent elements of S .

Definition (See [17]). A Γ -semiring S is said to be *right symmetric* if for $a, b, c \in S$, $a\Gamma b\Gamma c = 0$ implies $a\Gamma c\Gamma b = 0$. An ideal I of a Γ -semiring S is said to be *right symmetric* if $a\Gamma b\Gamma c \subseteq I$ implies $a\Gamma c\Gamma b \subseteq I$ for $a, b, c \in S$.

Analogously we can define *left symmetric Γ -semiring* and *left symmetric ideal*.

Proposition 8. *Let S be a Γ -semiring and I be an ideal of S . Then $\mathcal{P}(S/I) = \mathcal{P}(I)/I$.*

Proof. Let $s/I \in \mathcal{P}(S/I)$

$\Leftrightarrow s/I \in Q/I$ for all prime ideals Q of S containing I

$\Leftrightarrow s \in Q$ for all prime ideals Q of S containing I , as Q is a k -ideal

$\Leftrightarrow s \in \mathcal{P}(I)$

$\Leftrightarrow s/I \in \mathcal{P}(I)/I$.

Therefore, $\mathcal{P}(S/I) = \mathcal{P}(I)/I$. ■

Proposition 9. *Let S be an SN Γ -semiring and I be an ideal of S . If $(x\Gamma)^{n-1}x \subseteq I \Rightarrow x \in \mathcal{P}(I)$, then I is 2-primal.*

Proof. For any Γ -semiring S and any ideal I of S we have $\mathcal{P}(S/I) \subseteq \mathcal{N}(S/I)$ (Cf. Ref. Proposition 3.10 [17]). On the other hand let, $x/I \in \mathcal{N}(S/I)$. Since S is an SN Γ -semiring, S/I is an SN Γ -semiring. Then there exists a positive ineger say n such that $((x/I)\Gamma)^{n-1}x/I = 0/I$ which implies that $(x\Gamma)^{n-1}x \subseteq I$. By hypothesis $x \in \mathcal{P}(I)$. This implies that $x/I \in \mathcal{P}(I)/I$. Then by Proposition 8, $x/I \in \mathcal{P}(S/I)$. Thus $\mathcal{N}(S/I) \subseteq \mathcal{P}(S/I)$. Therefore, $\mathcal{P}(S/I) = \mathcal{N}(S/I)$. Hence I is 2-primal. ■

Proposition 10. *Let S be an SN Γ -semiring and I be an ideal of S . Then the following statements are equivalent:*

- (1) I is a 2-primal ideal of S .
- (2) $\mathcal{P}(I)$ is completely semiprime ideal of S .
- (3) $\mathcal{P}(I)$ is a left and right symmetric ideal of S .
- (4) $\mathcal{P}(I)$ has the IFP.

Proof. (1) implies (2). Let I be a 2-primal ideal of S . Then S/I is a 2-primal Γ -semiring. So $\mathcal{P}(S/I)$ is completely semiprime (Cf. Ref. Theorem 3.25 [17]). Now by Proposition 8, we have $\mathcal{P}(S/I) = \mathcal{P}(I)/I$. Thus $\mathcal{P}(I)/I$ is completely semiprime, so $\mathcal{P}(I)$ is completely semiprime.

(2) implies (3). Let $a\Gamma b\Gamma c \subseteq \mathcal{P}(I)$, where $a, b, c \in S$. Now $(c\Gamma a\Gamma b)\Gamma(c\Gamma a\Gamma b) = c\Gamma(a\Gamma b\Gamma c)\Gamma a\Gamma b \subseteq \mathcal{P}(I)$. Since $\mathcal{P}(I)$ is completely semiprime, $c\Gamma a\Gamma b \subseteq \mathcal{P}(I)$. Now $(a\Gamma b\Gamma a\Gamma c)\Gamma(a\Gamma b\Gamma a\Gamma c) = a\Gamma b\Gamma a\Gamma(c\Gamma a\Gamma b)\Gamma a\Gamma c \subseteq \mathcal{P}(I)$ as $\mathcal{P}(I)$ is an ideal of S . This implies that $a\Gamma b\Gamma a\Gamma c \subseteq \mathcal{P}(I)$. Again by similar argument $(b\Gamma a\Gamma c\Gamma b\Gamma a)\Gamma(b\Gamma a\Gamma c\Gamma b\Gamma a) = b\Gamma a\Gamma c\Gamma b\Gamma(a\Gamma b\Gamma a\Gamma c)\Gamma b\Gamma a \subseteq \mathcal{P}(I) \Rightarrow b\Gamma a\Gamma c\Gamma b\Gamma a \subseteq \mathcal{P}(I) \Rightarrow (a\Gamma c\Gamma b)\Gamma(a\Gamma c\Gamma b)\Gamma(a\Gamma c\Gamma b)\Gamma(a\Gamma c\Gamma b) = a\Gamma c\Gamma(b\Gamma a\Gamma c\Gamma b\Gamma a)\Gamma c\Gamma b\Gamma a\Gamma c\Gamma b \subseteq \mathcal{P}(I) \Rightarrow a\Gamma c\Gamma b \subseteq \mathcal{P}(I)$ as $\mathcal{P}(I)$ is completely semiprime. Hence $\mathcal{P}(I)$ is a right symmetric ideal of S . Also $(b\Gamma a\Gamma c)\Gamma(b\Gamma a\Gamma c) = b\Gamma(a\Gamma c\Gamma b)\Gamma a\Gamma c \subseteq \mathcal{P}(I) \Rightarrow b\Gamma a\Gamma c \subseteq \mathcal{P}(I)$. Hence $\mathcal{P}(I)$ is a left symmetric ideal of S . Therefore, $\mathcal{P}(I)$ is a left and a right symmetric ideal of S .

(3) implies (4). Let $x\Gamma y \subseteq \mathcal{P}(I)$, where $x, y \in S$. Suppose $s \in S$, then $s\Gamma x\Gamma y \subseteq \mathcal{P}(I)$. As $\mathcal{P}(I)$ is left symmetric, $x\Gamma s\Gamma y \subseteq \mathcal{P}(I)$. Therefore $x\Gamma s\Gamma y \subseteq \mathcal{P}(I)$. Hence $\mathcal{P}(I)$ has the IFP.

(4) implies (1). For any Γ -semiring S and any ideal I of S we have $\mathcal{P}(S/I) \subseteq \mathcal{N}(S/I)$. On the other hand let, $x/I \in \mathcal{N}(S/I)$. Since S is an SN Γ -semiring, S/I is an SN Γ -semiring. Then $((x/I)\Gamma)^{n-1}x/I = 0/I$ implies that $(x\Gamma)^{n-1}x \subseteq I$. Now we claim that $x \in \mathcal{P}(I)$. Suppose $x \notin \mathcal{P}(I)$, then there exists a prime ideal P of S containing I such that $x \notin P$, i.e. $x \in S - P$. Since P is a prime ideal of S , $S - P$ is an m-system. Then there exist $s_1 \in S, \alpha_1, \beta_1 \in \Gamma$ such that $x\alpha_1 s_1 \beta_1 x \in S \setminus P$. Again since $x\alpha_1 s_1 \beta_1 x, x \in S \setminus P$, applying m-system property $x\alpha_1 s_1 \beta_1 x\alpha_2 s_2 \beta_2 x \in S \setminus P$, for some $\alpha_2, \beta_2 \in \Gamma$ and $s_2 \in S$. Applying m-system property after finite step, we have $x\alpha_1 s_1 \beta_1 x\alpha_2 s_2 \beta_2 x \dots \alpha_{n-1} s_{n-1} \beta_{n-1} x \in S \setminus P$ for some $s_i \in S, \alpha_i, \beta_i \in \Gamma$, where $i = 1, 2, \dots, (n-1)$. Since $(x\Gamma)^{n-1}x \subseteq \mathcal{P}(I)$ and $\mathcal{P}(I)$ has the IFP, $x\alpha_1 s_1 \beta_1 x\alpha_2 s_2 \beta_2 x \dots \alpha_{n-1} s_{n-1} \beta_{n-1} x \in \mathcal{P}(I)$ i.e., $x\alpha_1 s_1 \beta_1 x\alpha_2 s_2 \beta_2 x \dots \alpha_{n-1} s_{n-1} \beta_{n-1} x \in P$, a contradiction. Therefore $x \in \mathcal{P}(I)$. Hence $x/I \in \mathcal{P}(I)/I = \mathcal{P}(S/I)$ by Proposition 8. So $\mathcal{P}(S/I) = \mathcal{N}(S/I)$ i.e., S/I is a 2-primal Γ -semiring. Hence I is a 2-primal ideal of S . ■

Proposition 11. Let S be a Γ -semiring and I be an ideal of S . If S/I satisfies (SI) then $x\Gamma y \subseteq I$ implies that $x\Gamma s\Gamma y \subseteq I$ for all $x, y \in S$ i.e., I has the IFP.

Proof. Let S be a Γ -semiring and I be an ideal of S such that S/I satisfies (SI). Let $x\Gamma y \subseteq I$. Then $(x/I)\Gamma(y/I) = 0/I$. So $y/I \in \text{ann}_R(x/I)$. Since S/I satisfies (SI), $(S/I)\Gamma\text{ann}_R(x/I) \subseteq \text{ann}_R(x/I)$ i.e., $(S/I)\Gamma(y/I) \subseteq \text{ann}_R(x/I)$ i.e., $(x/I)\Gamma(S/I)\Gamma(y/I) = 0/I$ i.e., $x\Gamma S\Gamma y \subseteq I$. This completes the proof. ■

Proposition 12. *Let S be a Γ -semiring and I be an ideal of S . If $S/\mathcal{P}(I)$ has no nonzero nilpotent elements, then $S/\mathcal{P}(I)$ satisfy (SI).*

Proof. Let S be a Γ -semiring and I be an ideal of S such that $S/\mathcal{P}(I)$ has no nonzero nilpotent elements. Then $S/\mathcal{P}(I)$ is a 2-primal Γ -semiring (Cf. Ref. Proposition 3.11, [17]). Hence $\mathcal{P}(I)$ is a 2-primal ideal of S . Now by Proposition 10(4), $\mathcal{P}(\mathcal{P}(I))$ has the IFP. Now $\mathcal{P}(\mathcal{P}(I)) = \mathcal{P}(I)$ (Cf. Ref. [10]). So $\mathcal{P}(I)$ has the IFP. Now we show that for any $a/\mathcal{P}(I) \in S/\mathcal{P}(I)$, $\text{ann}_R(a/\mathcal{P}(I))$ is an ideal of $S/\mathcal{P}(I)$. Let $b/\mathcal{P}(I) \in \text{ann}_R(a/\mathcal{P}(I))$. Then $(a/\mathcal{P}(I))\Gamma(b/\mathcal{P}(I)) = 0/\mathcal{P}(I)$. Then $a\Gamma b \subseteq \mathcal{P}(I)$, where $a, b \in S$. Since $\mathcal{P}(I)$ has the IFP, $a\Gamma S\Gamma b \subseteq \mathcal{P}(I)$, which implies that $(a/\mathcal{P}(I))\Gamma(S/\mathcal{P}(I))\Gamma(b/\mathcal{P}(I)) = 0/\mathcal{P}(I)$. Hence $(S/\mathcal{P}(I))\Gamma(b/\mathcal{P}(I)) \subseteq \text{ann}_R(a/\mathcal{P}(I))$. So $\text{ann}_R(a/\mathcal{P}(I))$ is a left ideal of $S/\mathcal{P}(I)$. Again we know $\text{ann}_R(a/\mathcal{P}(I))$ is a right ideal of $S/\mathcal{P}(I)$. Consequently $\text{ann}_R(a/\mathcal{P}(I))$ is an ideal of $S/\mathcal{P}(I)$. Therefore, $S/\mathcal{P}(I)$ satisfy (SI). ■

Proposition 13. *Let S be a Γ -semiring with unity. Then*

- (i) $I \subseteq P$ if and only if $N_I(P) \subseteq P$ for any ideal I and any prime ideal P of S .
- (ii) $N_I(P) \subseteq N_I^P$ for any prime ideal P and ideal I of S .
- (iii) If $I = P$ then $N_I(P) = P$ for any ideal I and any prime ideal P of S .
- (iv) If $P = Q$ if and only if $N_Q(P) = P$ for any prime ideals P and Q of S .

Proof. (i) Suppose $I \subseteq P$, then $\mathcal{P}(I) \subseteq P$. So, for any element $x \in N_I(P)$, there exists $b \in S - P$ such that $x\Gamma S\Gamma b \subseteq \mathcal{P}(I) \subseteq P$. Since P is a prime ideal of S and $b \in S - P$, we have $x \in P$. Therefore, $N_I(P) \subseteq P$. Conversely, let $N_I(P) \subseteq P$. Let $x \in I$. Now for any $y \in S - P$, $x\Gamma S\Gamma y \subseteq I$. Again $I \subseteq \mathcal{P}(I)$, so we have $x\Gamma S\Gamma y \subseteq \mathcal{P}(I)$. Hence $x \in N_I(P) \subseteq P$. This completes the proof.

(ii) Let $x \in N_I(P)$. Then there exists $b \in S - P$ such that $x\Gamma S\Gamma b \subseteq \mathcal{P}(I)$. Since P is a prime ideal of S , there exists $s \in S$ and $\alpha, \beta \in \Gamma$ such that $b\alpha s\beta b \in S - P$. Thus we have $x\Gamma b\alpha s\beta b \subseteq x\Gamma S\Gamma b \subseteq \mathcal{P}(I)$. Now since $b\alpha s\beta b \in S - P$, $x \in N_I^P$. Therefore, $N_I(P) \subseteq N_I^P$.

(iii) Let $P = I$ and $x \in I$. Since $I \subseteq \mathcal{P}(I)$ and I is an ideal, $x\Gamma S\Gamma S \subseteq I \subseteq \mathcal{P}(I)$. So for any $y \in S - P$, $x\Gamma S\Gamma y \subseteq \mathcal{P}(I)$. Hence $x \in N_I(P)$. Therefore, $P \subseteq N_I(P)$. Now by (i) $N_I(P) \subseteq P$. This completes the proof.

(iv) Suppose that $P = Q$, then by (iii), $N_Q(P) = P$. On the other hand, let $N_Q(P) = P$. Then $Q \subseteq N_Q(P) = P$ i.e., $Q \subseteq P$. Let $x \in P$. Then $x \in N_Q(P)$.

Then there exists $b \in S - P$ such that $x\Gamma S\Gamma b \subseteq \mathcal{P}(Q) \subseteq Q$ as Q is prime. Since $Q \subseteq P$, $b \in S - P \subseteq S - Q$. Hence $x \in Q$ as Q is prime. Therefore, $P = Q$. ■

Lemma 14. *Let S be a Γ -semiring and ‘ a ’ be a nonzero strongly nilpotent element of S . Then there exists a nonzero element b in S such that $b\Gamma b = 0$.*

Proof. Let ‘ a ’ be a nonzero strongly nilpotent element of S . Let n be the smallest positive integer such that $(a\Gamma)^{n-1}a = 0$.

Case 1. Suppose that n is odd say $n = 2k + 1$, where $1 \leq k < n$. Then we have $(a\Gamma)^{2k}a = 0$ which implies $(a\Gamma)^{2k}a\Gamma a = 0$. So

$$\begin{aligned} & \underbrace{(a\Gamma a\Gamma a \dots a\Gamma a)}_{\text{‘a’ appears (k+1) times}} \Gamma \underbrace{(a\Gamma a\Gamma a \dots a\Gamma a)}_{\text{‘a’ appears (k+1) times}} = 0. \\ \Rightarrow & \underbrace{(a\gamma a\gamma a \dots a\gamma a)}_{\text{‘a’ appears (k+1) times}} \Gamma \underbrace{(a\gamma a\gamma a \dots a\gamma a)}_{\text{‘a’ appears (k+1) times}} = 0 \text{ for all } \gamma \in \Gamma. \\ \text{Let } b = & \underbrace{a\gamma a\gamma a \dots a\gamma a}_{\text{‘a’ appears (k+1)-times}} \text{ for some nonzero } \gamma \in \Gamma. \text{ Then } b \neq 0 \text{ and } b\Gamma b = 0. \end{aligned}$$

Case 2. Suppose that n is even say $n = 2k$, where $1 \leq k < n$. Then we have $(a\Gamma)^{2k-1}a = 0$ which implies

$$\begin{aligned} & \underbrace{(a\Gamma a\Gamma a \dots a\Gamma a)}_{\text{‘a’ appears k-times}} \Gamma \underbrace{(a\Gamma a\Gamma a \dots a\Gamma a)}_{\text{‘a’ appears k-times}} = 0. \\ \Rightarrow & \underbrace{(a\gamma a\gamma a \dots a\gamma a)}_{\text{‘a’ appears k-times}} \Gamma \underbrace{(a\gamma a\gamma a \dots a\gamma a)}_{\text{‘a’ appears k-times}} = 0 \text{ for all } \gamma \in \Gamma. \\ \text{Let } b = & \underbrace{a\gamma a\gamma a \dots a\gamma a}_{\text{‘a’ appears k-times}} \text{ for some nonzero } \gamma \in \Gamma. \text{ Then } b \neq 0 \text{ and } b\Gamma b = 0. \quad \blacksquare \end{aligned}$$

Note 15. $\text{Spec}(S)$ denotes the set of all prime ideals of S .

Theorem 16. *Let S be an SN Γ -semiring with unity and I be an ideal of S . Then the following are equivalent:*

- (i) I is a 2-primal ideal of S ,
- (ii) $\mathcal{P}(I)$ has the IFP,
- (iii) $N_I(P)$ has the IFP for each $P \in \text{Spec}(S)$,
- (iv) $N_I(P) = \overline{N_I^P}$ for each $P \in \text{Spec}(S)$,
- (v) $N_I(P) = N_I^P$ for each $P \in \text{Spec}(S)$,
- (vi) $N_I^P \subseteq P$ for each prime ideal P which contains I ,
- (vii) $N_{P/\mathcal{P}(I)} \subseteq P/\mathcal{P}(I)$ for each prime ideal P which contains I ,

- (viii) $\overline{N_J^Q} \subseteq N_I(P)$ for any ideal $J \subseteq I$ and prime ideals P, Q of S such that $P \subseteq Q$,
- (ix) $N_J^Q \subseteq N_I(P)$ for any ideal $J \subseteq I$ and prime ideals P, Q of S such that $P \subseteq Q$,
- (x) $\overline{N_J^Q} \subseteq P$ for any ideal J and prime ideals P, Q of S such that $J \subseteq I \subseteq P \subseteq Q$,
- (xi) $N_J^Q \subseteq P$ for any ideal J and prime ideals P, Q of S such that $J \subseteq I \subseteq P \subseteq Q$,
- (xii) $N_{Q/P(I)} \subseteq P/P(I)$ for each prime ideal P, Q of S , such that $I \subseteq P \subseteq Q$.

Proof. (i) implies (ii). Let I be a 2-primal ideal of S . Then S/I is a 2-primal Γ -semiring. Let $x/P(I) \in \mathcal{N}(S/P(I))$. Since S is a SN Γ -semiring with unity, $S/P(I)$ is a SN Γ -semiring with unity. Then there exists a positive integer n such that $(x/P(I)\Gamma)^{n-1}(x/P(I)) = \mathcal{P}(I)$, i.e., $((x\Gamma)^{n-1}x)/P(I) = \mathcal{P}(I)$, i.e., $(x\Gamma)^{n-1}x \subseteq \mathcal{P}(I)$. Since I is a 2-primal ideal of S , then by Proposition 10, $\mathcal{P}(I)$ is a completely semiprime ideal of S . Hence $x \in \mathcal{P}(I)$, i.e., $x/P(I) = 0/P(I)$. Hence $S/P(I)$ has no strongly nilpotent elements. Then by Proposition 12, $S/P(I)$ satisfies (SI). Hence by Proposition 11, $\mathcal{P}(I)$ has the IFP.

(ii) implies (iii). Let $P \in \text{Spec}(S)$ and $x\Gamma y \subseteq N_I(P)$. Then $x\Gamma y\Gamma S\Gamma b \subseteq \mathcal{P}(I)$ for any $b \in S - P$. Now by (ii), $\mathcal{P}(I)$ has the IFP, so $x\Gamma S\Gamma y\Gamma S\Gamma b \subseteq \mathcal{P}(I)$ i.e. $(x\Gamma S\Gamma y)\Gamma S\Gamma b \subseteq \mathcal{P}(I)$, where $b \in S - P$. Therefore, $x\Gamma S\Gamma y \subseteq N_I(P)$ for each $P \in \text{Spec}(S)$. Thus $N_I(P)$ has the IFP for each $P \in \text{Spec}(S)$.

(iii) implies (i). Let $(a\Gamma)^{n-1}a \in I$, for some positive integer n . Claim: $a \in \mathcal{P}(I)$. Suppose $a \notin \mathcal{P}(I)$. Then there exists a prime ideal P which contains I , such that $a \notin P$ i.e., $a \in S \setminus P$. Since P is prime, $S \setminus P$ is an m-system. Then there exist $s_1 \in S, \alpha_1, \beta_1 \in \Gamma$ such that $a\alpha_1 s_1 \beta_1 a \in S \setminus P$. Again since $a\alpha_1 s_1 \beta_1 a, a \in S \setminus P$, applying m-system property $a\alpha_1 s_1 \beta_1 a\alpha_2 s_2 \beta_2 a \in S \setminus P$, for some $\alpha_2, \beta_2 \in \Gamma$ and $s_2 \in S$. Applying m-system property after finite step, we have $a\alpha_1 s_1 \beta_1 a\alpha_2 s_2 \beta_2 a \dots \alpha_{n-1} s_{n-1} \beta_{n-1} a \in S \setminus P$ for some $s_i \in S, \alpha_i, \beta_i \in \Gamma$, where $i = 1, 2, \dots, (n-1)$. Since $(a\Gamma)^{n-1}a \in I \subseteq N_I(P)$ and $N_I(P)$ has the IFP, $a\alpha_1 s_1 \beta_1 a\alpha_2 s_2 \beta_2 a \dots \alpha_{n-1} s_{n-1} \beta_{n-1} a \in N_I(P)$. Again by Proposition 13 (i), $N_I(P) \subseteq P$, then $a\alpha_1 s_1 \beta_1 a\alpha_2 s_2 \beta_2 a \dots \alpha_{n-1} s_{n-1} \beta_{n-1} a \in P$, a contradiction. Hence $a \in \mathcal{P}(I)$. Now by Proposition 9, I is a 2-primal ideal of S .

(i) implies (iv). Let $a \in \overline{N_I^P}$ for each $P \in \text{Spec}(S)$. Then $(a\Gamma)^{n-1}a \subseteq N_I^P$, for some positive integer n . Hence there exists $b \in S - P$ such that $(a\Gamma)^{n-1}a\Gamma b \subseteq \mathcal{P}(I)$ i.e., $(a\Gamma)^n b \subseteq \mathcal{P}(I)$. Since I is a 2-primal ideal of S , by Proposition 10(3), $\mathcal{P}(I)$ is a left and a right symmetric ideal of S . Suppose $n = 1$, $a\Gamma b \subseteq \mathcal{P}(I)$. Let $n = 2$, $a\Gamma a\Gamma b \subseteq \mathcal{P}(I) \Rightarrow a\Gamma b\Gamma a \subseteq \mathcal{P}(I)$ (as $\mathcal{P}(I)$ is a right symmetric

ideal of $S \Rightarrow a\Gamma b\Gamma a\Gamma b \subseteq \mathcal{P}(I)$ (as $\mathcal{P}(I)$ is an ideal of S). Now by Proposition 10(2), $\mathcal{P}(I)$ is a completely semiprime ideal of S , then we have $a\Gamma b \subseteq \mathcal{P}(I)$. Let $n = 3$. Then $a\Gamma a\Gamma a\Gamma b \subseteq \mathcal{P}(I) \Rightarrow b\Gamma a\Gamma a\Gamma a\Gamma b \subseteq \mathcal{P}(I)$ (as $\mathcal{P}(I)$ is an ideal of $S \Rightarrow a\Gamma b\Gamma a\Gamma a\Gamma b \subseteq \mathcal{P}(I)$ (as $\mathcal{P}(I)$ is a left symmetric ideal of $S \Rightarrow a\Gamma b\Gamma a\Gamma b\Gamma a \subseteq \mathcal{P}(I)$ (as $\mathcal{P}(I)$ is a right symmetric ideal of $S \Rightarrow a\Gamma b\Gamma a\Gamma b\Gamma a\Gamma b\Gamma a\Gamma b \subseteq \mathcal{P}(I)$ (as $\mathcal{P}(I)$ is an ideal of S). Hence by Proposition 10(2), $\mathcal{P}(I)$ is a completely semiprime ideal of S , then we have $a\Gamma b \subseteq \mathcal{P}(I)$. Continuing this process for $n \geq 2$, $(a\Gamma)^n b \subseteq \mathcal{P}(I) \Rightarrow \underbrace{(a\Gamma b)\Gamma(a\Gamma b)\Gamma(a\Gamma b)\Gamma \dots \Gamma(a\Gamma b)}_{(n\text{-times})} \subseteq \mathcal{P}(I)$. If n is even, then $a\Gamma b \subseteq \mathcal{P}(I)$ (by

Proposition 10(2), $\mathcal{P}(I)$ is a completely semiprime ideal of S). If n is odd, then multiplying by $a\Gamma b$ and applying Proposition 10(2) we have $a\Gamma b \subseteq \mathcal{P}(I)$. Now by Proposition 10(4), we have $a\Gamma S\Gamma b \subseteq \mathcal{P}(I)$, where $b \in \overline{S - P}$. Hence $a \in N_I(P)$. Again by Proposition 13, we have, $N_I(P) \subseteq N_I^P \subseteq \overline{N_I^P}$. Therefore, $N_I(P) = \overline{N_I^P}$.

(iv) implies (v). Since $N_I(P) \subseteq N_I^P \subseteq \overline{N_I^P}$, by (iv) we have $N_I(P) = N_I^P$.

(v) implies (vi). Let P be a prime ideal of S which contains I . Then by Proposition 13 (i), we have $N_I(P) \subseteq P$. Now by (v) we have, $N_I(P) = N_I^P$. Hence $N_I^P \subseteq P$.

(vi) implies (vii). Let $a/\mathcal{P}(I) \in N_{P/\mathcal{P}(I)}$. Then there exists $b/\mathcal{P}(I) \in S/\mathcal{P}(I) - P/\mathcal{P}(I)$ such that $(a/\mathcal{P}(I))\Gamma(b/\mathcal{P}(I)) \subseteq \mathcal{P}(S/\mathcal{P}(I)) = \mathcal{P}(\mathcal{P}(I))/\mathcal{P}(I)$ (by Proposition 8). This implies that $a\Gamma b \subseteq \mathcal{P}(I)$ as $\mathcal{P}(\mathcal{P}(I)) = \mathcal{P}(I)$, where $b \in S - P$. So $a \in N_I^P$. Hence by (vi), $a \in P$. This implies that $a/\mathcal{P}(I) \in P/\mathcal{P}(I)$. Therefore, $N_{P/\mathcal{P}(I)} \subseteq P/\mathcal{P}(I)$.

(vii) implies (i). First we shall show that $S/\mathcal{P}(I)$ is reduced. Suppose, $S/\mathcal{P}(I)$ is not reduced. Then there exists a nonzero nilpotent element say $a/\mathcal{P}(I) \in S/\mathcal{P}(I)$. Since S is an SN Γ -semiring, $S/\mathcal{P}(I)$ is an SN Γ -semiring. Then $a/\mathcal{P}(I) \in S/\mathcal{P}(I)$ is a strongly nilpotent element. Hence by Lemma 14, there exists a nonzero element say $b/\mathcal{P}(I) \in S/\mathcal{P}(I)$ such that $(b/\mathcal{P}(I))\Gamma(b/\mathcal{P}(I)) = \mathcal{P}(I)/\mathcal{P}(I)$ i.e., $(b/\mathcal{P}(I))\Gamma(b/\mathcal{P}(I)) = \mathcal{P}(S/\mathcal{P}(I))$ (by Proposition 8). Since $b/\mathcal{P}(I)$ is a nonzero element of $S/\mathcal{P}(I)$, $b \notin \mathcal{P}(I)$. So $b/\mathcal{P}(I) \in S/\mathcal{P}(I) - P/\mathcal{P}(I)$ for some prime ideal P of S containing I . Hence $b/\mathcal{P}(I) \in N_{P/\mathcal{P}(I)}$. Now by (vii) $N_{P/\mathcal{P}(I)} \subseteq P/\mathcal{P}(I)$, so $b/\mathcal{P}(I) \in P/\mathcal{P}(I)$ which is a contradiction. Therefore, $S/\mathcal{P}(I)$ has no nonzero strongly nilpotent elements. Thus $S/\mathcal{P}(I)$ is reduced. Then using Proposition 12, we have $S/\mathcal{P}(I)$ satisfy (SI). Then by Proposition 11, $\mathcal{P}(I)$ has the IFP. Then by Proposition 10, we have I is a 2-primal ideal of S .

(i) implies (viii). Let $x \in \overline{N_J^Q}$. Then there exists a positive integer n such that $(x\Gamma)^{n-1}x \subseteq N_J^Q$. Then $(x\Gamma)^{n-1}x\Gamma y \subseteq \mathcal{P}(J)$ for some $y \in S - Q$. Since $P \subseteq Q$ and $J \subseteq I$, $(x\Gamma)^{n-1}x\Gamma y \subseteq \mathcal{P}(I)$ for some $y \in S - P$. Since I is 2-primal, by Proposition

10, $\mathcal{P}(I)$ is a left and right symmetric ideal and completely semiprime ideal of S . Proceeding as in the proof of (i) implies (iv), we get $x\Gamma y \subseteq \mathcal{P}(I)$, where $y \in S - P$. Again by same proposition $\mathcal{P}(I)$ has the IFP, then $x\Gamma y \subseteq \mathcal{P}(I) \Rightarrow x\Gamma S\Gamma y \subseteq \mathcal{P}(I)$. Hence $x \in N_I(P)$. Therefore, $\overline{N_J^Q} \subseteq N_I(P)$.

(viii) implies (ix). It is obvious as $N_J^Q \subseteq \overline{N_J^Q}$.

(ix) implies (xi). Let $J \subseteq I \subseteq P \subseteq Q$. Then by (ix), $N_J^Q \subseteq N_I(P)$. Again since $I \subseteq P$, by Proposition 13(i), $N_I(P) \subseteq P$. Hence $N_J^Q \subseteq P$ for any ideal J and prime ideals P, Q of S such that $J \subseteq I \subseteq P \subseteq Q$.

(xi) implies (vi). On assuming $I = J$ and $P = Q$ in (xi), we get $N_I^P \subseteq P$.

(vi) implies (xii). Let $a/\mathcal{P}(I) \in N_{Q/\mathcal{P}(I)}$. Then there exists $b/\mathcal{P}(I) \in S/\mathcal{P}(I) - Q/\mathcal{P}(I)$ such that $(a/\mathcal{P}(I))\Gamma(b/\mathcal{P}(I)) \subseteq \mathcal{P}(S/\mathcal{P}(I)) = \mathcal{P}(I)/\mathcal{P}(I)$ (by Proposition 8). This implies that $a\Gamma b \subseteq \mathcal{P}(I)$, where $b \in S - Q$. This implies that $a\Gamma b \subseteq \mathcal{P}(I)$, where $b \in S - P$ as $P \subseteq Q$. Hence $a \in N_I^P$. Since $I \subseteq P$, by (vi) $N_I^P \subseteq P$, so $a \in P$. This implies that $a/\mathcal{P}(I) \in P/\mathcal{P}(I)$. Therefore, $N_{Q/\mathcal{P}(I)} \subseteq P/\mathcal{P}(I)$.

(xii) implies (vii). On assuming $Q = P$ in (xii), we get $N_{P/\mathcal{P}(I)} \subseteq P/\mathcal{P}(I)$.

(viii) implies (x). Let I, J be two any ideals of S and P, Q be two prime ideals of S such that $J \subseteq I \subseteq P \subseteq Q$. Now by (viii), we have $\overline{N_J^Q} \subseteq N_I(P)$. Again since $I \subseteq P$, by Proposition 13(i), $N_I(P) \subseteq P$. Therefore, $\overline{N_J^Q} \subseteq P$.

(x) implies (xi). Let I, J be two any ideals of S and P, Q be two prime ideals of S such that $J \subseteq I \subseteq P \subseteq Q$. Then by (x), $\overline{N_J^Q} \subseteq P$. Again by Proposition 7, we have $N_J^Q \subseteq \overline{N_J^Q}$. Therefore, $N_J^Q \subseteq P$. ■

Corollary 17. *Let S be an SN Γ -semiring with unity and I be a 2-primal ideal of S . Then for any prime ideal P of S , $I \subseteq P$ if and only if $N_I^P \subseteq P$.*

Proof. Let $I \subseteq P$. Then by Theorem 16 (vi), we have $N_I^P \subseteq P$. Conversely let, $N_I^P \subseteq P$. Let $x \in I$. Then for any $y \in S - P$ we have $x\Gamma y \subseteq I \Rightarrow x\Gamma y \subseteq \mathcal{P}(I)$ as $I \subseteq \mathcal{P}(I)$, where $y \in S - P$. Hence $x \in N_I^P$ and so $x \in P$. Thus $I \subseteq P$. ■

Corollary 18. *Let S be an SN Γ -semiring with unity. Then $I = P$ if and only if $N_I^P = P$ for any completely prime ideal I and prime ideal P of S .*

Proof. Let I be any completely prime ideal and P be any prime ideal of S such that $I = P$. Now we have, $I \subseteq \mathcal{P}(I) \subseteq P$. Since $I = P$, $I = \mathcal{P}(I)$. Hence $\mathcal{P}(I)$ is a completely prime ideal and hence a completely semiprime ideal of S . Then by Proposition 10, I is a 2-primal ideal of S . By Corollary 17, $N_I^P \subseteq P$. Hence $P = I \subseteq N_I^P \subseteq P$. Therefore, $N_I^P = P$. Conversely let, $N_I^P = P$ for any completely prime ideal I and prime ideal P of S . Then by Corollary 17, $I \subseteq P$.

Let $x \in P$. Then $x \in N_I^P$. So $x\Gamma y \subseteq \mathcal{P}(I)$ for some $y \in S - P$. Since I is a completely prime ideal of S , I is a prime ideal of S . Hence $\mathcal{P}(I) = I$. So $x\Gamma y \subseteq I$. Since $I \subseteq P$ and $y \in S - P$, $y \in S - I$. Again I being a completely prime ideal, then $x \in I$ as $y \in S - P$. Hence $I = P$. ■

Corollary 19. *Let S be an SN Γ -semiring with unity. Then the following are equivalent:*

- (i) S is a 2-primal Γ -semiring,
- (ii) $\mathcal{P}(S)$ has the IFP,
- (iii) $N(P)$ has the IFP for each $P \in \text{Spec}(S)$,
- (iv) $N(P) = \overline{N_P}$ for each $P \in \text{Spec}(S)$,
- (v) $N(P) = N_P$ for each $P \in \text{Spec}(S)$,
- (vi) $N_P \subseteq P$ for each $P \in \text{Spec}(S)$,
- (vii) $N_{P/\mathcal{P}(S)} \subseteq P/\mathcal{P}(S)$ for each $P \in \text{Spec}(S)$,
- (viii) $\overline{N_Q} \subseteq N(P)$ for any prime ideals P, Q of S such that $P \subseteq Q$,
- (ix) $N_Q \subseteq N(P)$ for any prime ideals P, Q of S such that $P \subseteq Q$,
- (x) $\overline{N_Q} \subseteq P$ for any prime ideals P, Q of S such that $P \subseteq Q$,
- (xi) $N_Q \subseteq P$ for any prime ideals P, Q of S such that $P \subseteq Q$,
- (xii) $N_{Q/\mathcal{P}(S)} \subseteq P/\mathcal{P}(S)$ for each prime ideals P, Q , such that $P \subseteq Q$.

Proof. The proof follows from Theorem 16. ■

Acknowledgement

The authors are thankful to the learned referee for his kind suggestions.

REFERENCES

- [1] C. Selvaraj and S. Petchimuthu, *Characterization of 2-Primal Γ -Rings*, Southeast Asian Bull. Math. **34** (2010) 1083–1094.
- [2] C. Selvaraj and S. Petchimuthu, *Characterization of 2-primal ideals*, Far East J. Math. Sci. **28** (2008) 249–256.
- [3] G.F. Birkenmeier, H.E. Heatherly and E.K. Lee, *Completely Prime Ideals and Associated Radicals*, in: Proc. Biennial Ohio State-Denison Conference 1992, S.K. Jain and S.T. Rizvi (Ed(s)), (World Scientific, New Jersey, 1993) 102–129.
- [4] M.M.K. Rao, *Γ -semiring-I*, Southeast Asian Bull. Math. **19** (1995) 49–54.
- [5] M.M.K. Rao, *Γ -semiring-II*, Southeast Asian Bull. Math. **21** (1997) 281–287.
- [6] N. Nobusawa, *On a generalization of the ring theory*, Osaka J. Math. **1** (1964) 81–89.

- [7] P.J. Allen and W.R. Windham, *Operator semigroup with applications to semiring*, Publicationes Mathematicae **20** (1973) 161–175.
- [8] S.K. Sardar, *On Jacobson radical of a Γ -semiring*, Far East J. Math. Sci. **35** (2005) 1–9.
- [9] S.K. Sardar and U. Dasgupta, *On primitive Γ -semiring*, Far East J. Math. Sci. **34** (2004) 1–12.
- [10] T.K. Dutta and S.K. Sardar, *On prime ideals and prime radical of a Γ -semirings*, ANALELE ȘTIINȚIFICE ALE UNIVERSITĂȚII “AL.I.CUZA” IAȘI, Matematică **Tomul XLVI.s.I.a, f.2** (2000) 319–329.
- [11] T.K. Dutta and S.K. Sardar, *Semiprime ideals and irreducible ideals of Γ -semirings*, Novi Sad J. Math. **30** (2000) 97–108.
- [12] T.K. Dutta and S.K. Sardar, *On the operator semirings of a Γ -semiring*, Southeast Asian Bull. Math., Springer-Verlag **26** (2002) 203–213.
- [13] T.K. Dutta and S.K. Sardar, *On Levitzki radical of a Γ -semiring*, Bull. Calcutta Math. Soc. **95** (2003) 113–120.
- [14] T.K. Dutta and S. Dhara, *On uniformly strongly prime Γ -semirings*, Southeast Asian Bull. Math. **30** (2006) 39–48.
- [15] T.K. Dutta and S. Dhara, *On uniformly strongly prime Γ -semirings (II)*, Discuss. Math. General Algebra and Appl. **26** (2006) 219–231.
- [16] T.K. Dutta and S. Dhara, *On strongly prime Γ -semirings*, ANALELE ȘTIINȚIFICE ALE UNIVERSITĂȚII “AL.I.CUZA” DIN IAȘI(S. N.) Matematică **Tomul LV, f.1** (2009) 213–224.
- [17] T.K. Dutta and S. Dhara, *On 2-primal Γ -semirings*, Southeast Asian Bull. Math. **37** (2013) 699–714.
- [18] T.K. Dutta, K.P. Shum and S. Dhara, *On NI Γ -semirings*, Int. J. Pure and Appl. Math. **84** (2013) 279–298. doi:10.12732/ijpam.v84i3.14
- [19] W.E. Barnes, *On the Γ -rings of Nobusawa*, Pacific J. Math. **18** (1966) 411–422. doi:10.2140/pjm.1966.18.411

Received 10 September 2013

Revised 27 October 2013