

4 **SOME ALGEBRAIC CHARACTERISATIONS OF**
5 **GENERALISED MIDDLE BOL LOOPS**

6 BENARD OSOBA¹

7 *Bells University of Technology*
8 *Ota, Ogun State, Nigeria*

9 **e-mail:** benardomth@gmail.com, b_osoba@bellsuniversity.edu.ng

10 ABDULKAREEM ABDULAFEEZ OLALEKAN

11 *Department of Mathematics*
12 *Lagos State University Ojo*
13 *Lagos State, 102101 Nigeria*

14 **e-mail:** abdulafeez.abdulkareem@lasu.edu.ng

15 YAKUB TUNDE OYEBO

16 *Department of Mathematics*
17 *Lagos State University Ojo*
18 *Lagos State, 102101 Nigeria*

19 **e-mail:** oyeboyt@yahoo.com, yakub.oyebo@lasu.edu.ng

20 AND

21 ANTHONY OYEM

22 *Department of Mathematics*
23 *University of Lagos, Akoka, Lagos*

24 **e-mail:** tonyoyem@yahoo.com

25 **Abstract**

26 In this article, some algebraic characterisations of generalised middle
27 Bol loop (GMBL) using its parastrophes and holomorph were studied. In
28 particular, it was shown that if the generalised map α is bijective such
29 $\alpha : e \rightarrow e$, then the (12)– parastrophe of GMBL is a GMBL. The conditions
30 for (13)– and (123)–parastrophes of a GMBL to be GMBL of exponent
31 two were unveiled. We further established that a commutative (13)– and

¹All correspondence to be addressed to this author.

(123)–parastrophes of GMBL has an inverse properties. (23)– parastrophe of Q was shown to be super α –elastic property if it has a middle symmetric while (132)–parastrophe of Q satisfies left α –symmetric. It is further shown that a commutative (13)– and (123)– parastrophes of Q are generalised Moufang loops of exponent two. Also, commutative (132)– and (23)– parastrophes of Q are shown to be Steiner loops. A necessary and sufficient condition for holomorph of generalised middle Bol loop to be GMBL was presented. The holomorph of a commutative loop was shown to be a commutative generalised middle Bol loop if and only if the loop is a GMBL.

Keywords: loop, parastrophe, Holomorph, Generalised middle Bol loop.

2020 Mathematics Subject Classification: Primary 20N05; Secondary 08A05.

1 INTRODUCTION

1.1 Quasigroups and Loops

Let Q be a non -empty set. Define a binary operation " \cdot " on Q . If $x \cdot y \in Q$ for all $x, y \in Q$, then the pair (Q, \cdot) is called a groupoid or magma. If the equations $a \cdot x = b$ and $y \cdot a = b$ have unique solutions $x, y \in Q$ for all $a, b \in Q$ then (Q, \cdot) is called a quasigroup. Let (Q, \cdot) be a quasigroup and let there exist a unique element $e \in Q$ called the identity element such that for all $x \in Q, x \cdot e = e \cdot x = x$, then (Q, \cdot) is called a loop. At times, we shall write xy instead of $x \cdot y$ and stipulate that " \cdot " has lower priority than juxtaposition among factors to be multiplied. Let (Q, \cdot) be a groupoid and a be a fixed element in Q , then the left and right translations L_a and R_a of a are respectively defined by $xL_a = a \cdot x$ and $xR_a = x \cdot a$ for all $x \in Q$. It can now be seen that a groupoid

(Q, \cdot) is a quasigroup if its left and right translation mappings are permutations. Since the left and right translation mappings of a quasigroup are bijective, then the inverse mappings L_a^{-1} and R_a^{-1} exist.

Let

$$a \setminus b = bL_a^{-1} = aM_b \quad \text{and} \quad a/b = aR_b^{-1} = bM_a^{-1}$$

and note that

$$a \setminus b = c \iff a \cdot c = b \quad \text{and} \quad a/b = c \iff c \cdot b = a.$$

Thus, for any quasigroup (Q, \cdot) , we have two new binary operations; right division ($/$) and left division (\setminus). M_a is the middle translation for any fixed $a \in Q$. Consequently, (Q, \setminus) and $(Q, /)$ are also quasigroups. Using the operations (\setminus) and ($/$), the definition of a loop can be restated as follows.

65 **Definition 1.1.** A loop $(Q, \cdot, /, \backslash, e)$ is a set Q together with three binary oper-
 66 ations (\cdot) , $(/)$, (\backslash) and one nullary operation e such that

- 67 (i) $a \cdot (a \backslash b) = b$, $(b/a) \cdot a = b$ for all $a, b \in Q$,
 68 (ii) $a \backslash a = b/b$ or $e \cdot a = a \cdot e = a$ for all $a, b \in Q$.

69 We also stipulate that $(/)$ and (\backslash) have higher priority than (\cdot) among factors
 70 to be multiplied. For instance, $a \cdot b/c$ and $a \cdot b \backslash c$ stand for $a(b/c)$ and $a(b \backslash c)$
 71 respectively.

72 In a loop (Q, \cdot) with identity element e , the *left inverse element* of $x \in Q$ is
 73 the element $xJ_\lambda = x^\lambda \in Q$ such that

$$x^\lambda \cdot x = e$$

74 while the *right inverse element* of $x \in G$ is the element $xJ_\rho = x^\rho \in G$ such that

$$x \cdot x^\rho = e.$$

75 It is well known that every quasigroup (Q, \cdot) belongs to a set of six quasigroups,
 76 called adjugates by (Fisher, Yates [5] 1934), conjugates by (Stein , 1957) and
 77 parastrophes by (Belousov [4], 1967)

78 A binary groupoid (Q, A) with a binary operation “ A ” such that in the
 79 equality $A(x_1, x_2) = x_3$ knowledge of any 2 elements of x_1, x_2, x_3 uniquely spec-
 80 ifies remaining one is called a binary quasigroup. It follows that any quasi-
 81 group (Q, A) , associate $(3! - 1)$ quasigroups called parastrophes of quasigroup
 82 (Q, A) ; $A(x_1, x_2) = x_3 \iff A^{(12)}(x_2, x_1) = x_3 \iff A^{(13)}(x_3, x_2) = x_1 \iff$
 83 $A^{(23)}(x_1, x_2) = x_3 \iff A^{(123)}(x_2, x_3) = x_1 \iff A^{(132)}(x_3, x_1) = x_2$. [see (Shcherba-
 84 cov [32], 2008)]. For more on quasigroups and loops, check [31, 33].

85 1.2 Middle Bol Loop and its Generalisation

86 **Definition 1.2.** A loop (Q, \cdot) is called a middle Bol loop if

$$(x/y)(z \backslash x) = (x/(zy))x \text{ or } (x/y)(z \backslash x) = x((zy) \backslash x) \quad (1)$$

87 for all $x, y \in Q$.

88 Middle Bol loop were first studied in the work of V. D. Belousov [4], where he
 89 gave identity (1) characterizing loops that satisfy the universal anti-automorphic
 90 inverse property. After this beautiful characterisation by Belousov and the lay-
 91 ing of foundations for a classical study of this structure, Gwaramija in [6] gave
 92 isostrophic connection between right(left) with middle Bol loop.

93 Grecu [16] showed that the right multiplication group of a middle Bol loop
 94 coincides with the left multiplication group of the corresponding right Bol loop.

95 After that, middle Bol loops resurfaced in literature in 1994 and 1996 when Syrbu
 96 [20, 21] considered them in relation to the universality of the elasticity law. In
 97 2003, Kuznetsov [15], while studying gyrogroups (a special class of Bol loops)
 98 established some algebraic properties of middle Bol loop and designed a method
 99 of constructing a middle Bol loop from a gyrogroup.

100 In 2010, Syrbu [22] studied the connections between structure and properties
 101 of middle Bol loops and of the corresponding left Bol loops. It was noted that two
 102 middle Bol loops are isomorphic if and only if the corresponding left (right) Bol
 103 loops are isomorphic, and a general form of the autotopisms of middle Bol loops
 104 was deduced. Relations between different sets of elements, such as nucleus, left
 105 (right, middle) nuclei, the set of Moufang elements, the center of a middle Bol
 106 loop and left Bol loop were established. In 2012, Grecu and Syrbu [17] proved
 107 that two middle Bol loops are isotopic if and only if the corresponding right (left)
 108 Bol loops are isotopic.

109 In 2012, Drapal and Shcherbacov [18] rediscovered the middle Bol identities in
 110 a new way. In 2013, Syrbu and Grecu [19] established a necessary and sufficient
 111 condition for the quotient loop of a middle Bol loop and of its corresponding
 112 right Bol loop to be isomorphic. In 2014, Grecu and Syrbu [24] established that
 113 the commutant (centrum) of a middle Bol loop is an AIP-subloop and gave a
 114 necessary and sufficient condition when the commutant is an invariant under
 115 the existing isotropy between middle Bol loop and the corresponding right Bol
 116 loop and the same authors presented a study of loops with invariant flexibility
 117 law under the isotropy of loop [23].

118 In 2017, Jaiyéḡlá et al. [8] presented the holomorphic structure of middle Bol
 119 loop and showed that the holomorph of a commutative loop is a commutative
 120 middle Bol loop if and only if the loop is a middle Bol loop and its automorphism
 121 group is abelian. Adeniran et al. [1], Jaiyéḡlá and Popoola [13] studied gener-
 122 alised Bol loops. It was revealed in [10] that isotopy-isomorphy is a necessary
 123 and sufficient condition for any distinct quasigroups to be parastophic invariance
 124 relative to the associative law.

125 (Osoba et al. [25] and [26]) investigate further the multiplication group of
 126 middle Bol loop in relation to left Bol loop and the relationship of multiplication
 127 groups and isotropic quasigroups respectively while Jaiyéḡlá [11, 12] studied
 128 second Smarandache Bol loops. The Smarandache nuclei of second Smarandache
 129 Bol loops was further studied by Osoba [27].

130 (Jaiyéḡlá et al. [7], 2015) in furtherance to their exploit obtained new alge-
 131 braic identities of middle Bol loop, where necessary and sufficient conditions for
 132 a bi-variate mapping of a middle Bol loop to have RIP, LIP, RAIP, LAIP and
 133 flexible property were presented. Additional algebraic properties of middle Bol
 134 loop were announced in (Jaiyéḡlá et al. [9], 2021).

135 The new algebraic connections between right and middle Bol loops and their

136 cores were unveiled by (Osoba and Jaiyéplá (2022),[28]). More results on the
 137 algebraic properties of middle Bol loops using its parastrophes was presented by
 138 (Oyebo and Osoba, [30]). The paper revealed some of the algebraic properties the
 139 parastrophic structures of middle Bol loop shared with its underline structure.
 140 The connections between middle Bol loop and right Bol loop with their crypto-
 141 automorphism features were unveiled in [29] by Oyebo et.al. In [14], Bryant-
 142 Schneider group of middle Bol loop with some of the isostrophy-group invariance
 143 results was linked. It was further shown that some subgroups of the Bryant-
 144 Schneider group of a middle Bol loop were isomorphic to the automorphism and
 145 pseudo-aumorphism groups of its corresponding right (left) Bol loop.

146 A generalised middle Bol loop characterised by

$$(x/y)(z^\alpha \backslash x^\alpha) = x(z^\alpha y \backslash x^\alpha) \quad (2)$$

147 was first introduced in [2], as a consequence of a generalised Moufang loop with
 148 universal α -elastic property where the map $\alpha : Q \mapsto Q$ is a homomorphism.
 149 Thus, if $\alpha : x \mapsto x$, then identity of generalised middle Bol loop reduces to
 150 the identity of middle Bol loop. The authors in [3], presented the basic algebraic
 151 properties of generalised middle Bol loop, where it revealed the necessary and suf-
 152 ficient conditions for the identity to satisfies left(right) inverse and α -alternative
 153 property was also presented.

154 Furtherance to earlier studies, this paper investigates some structural char-
 155 acterisation of generalised middle Bol loop using its parastrophes and holomorph.
 156 The second section provides preliminaries for necessary background of the study.
 157 Section 3 contains the main results where the parastrophic characterisation of
 158 generalised middle Bol loop is presented. It is shown that a (12)-parastrophe of
 159 a generalised middle Bol is also a generalised middle Bol loop and further estab-
 160 lished the conditions for (13)- and (123)-parastrophes of Q to be GMBL. We
 161 further investigate the algebraic properties of the parastrophes to obtain some of
 162 the related properties and identities they share with the underline structure. In-
 163 terestingly, some new identities are found. In the fourth section, the holomorphic
 164 characterisations of generalised middle Bol loop is studied and the necessary and
 165 sufficient condition is found.

166 **2 PRELIMINARIES**

167 **Definition 2.1.** A loop $(Q, \cdot, /, \backslash)$ is called a generalised middle Bol loop if is
 168 satisfies the identity

$$(x/y)(z^\alpha \backslash x^\alpha) = (x/(z^\alpha y))x^\alpha \quad (3)$$

169 **Definition 2.2.** For any non-empty set Q , the set of all permutations on Q forms
 170 a group $SYM(Q)$ called the symmetric group of Q . Let (Q, \cdot) be a loop and let

171 $A, B, C \in SYM(Q)$. If

$$xA \cdot yB = (x \cdot y)C \quad \forall x, y \in Q$$

172 then the triple (A, B, C) is called an autotopism (ATP) and such triples form a
173 group $AUT(Q, \cdot)$ called the autotopism group of (Q, \cdot) . Also, suppose that

$$xA \cdot yB = (y \cdot x)C \quad \forall x, y \in Q$$

174 then the triple (A, B, C) is called anti-autotopism (AATP). If $A = B = C$, then
175 A is called an automorphism of (Q, \cdot) which form a group $AUM(Q, \cdot)$ called the
176 automorphism group of (Q, \cdot) .

177 **Definition 2.3.** A groupoid (quasigroup) (Q, \cdot) is said to have the

- 178 1. left inverse property (LIP) if there exists a mapping $J_\lambda : x \mapsto x^\lambda$ such that
179 $x^\lambda \cdot xy = y$ for all $x, y \in Q$.
- 180 2. right inverse property (RIP) if there exists a mapping $J_\rho : x \mapsto x^\rho$ such
181 that $yx \cdot x^\rho = y$ for all $x, y \in Q$.
- 182 3. inverse property (IP) if it has both the LIP and RIP. for all $x, y \in Q$.
- 183 4. flexibility or elasticity if $xy \cdot x = x \cdot yx$ holds for all $x, y \in Q$.
- 184 5. α -elastic if $xy \cdot x^\alpha = x \cdot yx^\alpha$ holds for all $x, y \in Q$.
- 185 6. super α -elastic if $(x \cdot y^\alpha) \cdot x^\alpha = x \cdot (y^\alpha \cdot x^\alpha)$ holds for all $x, y \in Q$.
- 186 7. cross inverse property (CIP) if there exist mapping $J_\lambda : x \mapsto x^\lambda$ or $J_\rho :$
187 $x \mapsto x^\rho$ such that $xy \cdot x^\rho = y$ or $x \cdot yx^\rho = y$ or $x^\lambda \cdot yx = y$ or $x^\lambda y \cdot x = y$ for
188 all $x, y \in Q$.

189 **Definition 2.4.** A loop (Q, \cdot) is said to be

- 190 1. commutative loop if $R_x = L_x$ and a commutative square loop if $R_x^2 = L_x^2$
191 for all $x, y \in Q$
- 192 2. an automorphic inverse property loop (AIPL) if $(xy)^{-1} = x^{-1}y^{-1}$ for all
193 $x, y \in Q$
- 194 3. an anti- automorphic inverse property loop (AAIPL) if $(xy)^{-1} = y^{-1}x^{-1}$
195 for all $x, y \in Q$.

196 **Definition 2.5.** [31] Moufang loops are loops satisfying the identities $(xy \cdot z)y =$
197 $x(y \cdot zy), yz \cdot xy = y(zx \cdot y)$ and $(yz \cdot y)x = y(z \cdot yx)$

198 **Definition 2.6.** A groupoid (quasigroup) (Q, \cdot) is

- 199 1. right symmetric if $yx \cdot x = y$ for all $x, y \in Q$
- 200 2. left symmetric if $x \cdot xy = y$ for all $x, y \in Q$
- 201 3. middle symmetric if $x \cdot yx = y$ or $xy \cdot x = y$ for all $x, y \in Q$
- 202 4. idempotent if $x \cdot x = x$ for all $x \in Q$
- 203 5. right α -symmetric if $y^\alpha x \cdot x = y^\alpha$ for all $x, y \in Q$
- 204 6. left α -symmetric if $x \cdot xy^\alpha = y^\alpha$ for all $x, y \in Q$
- 205 7. middle α -symmetric if $x \cdot y^\alpha x = y^\alpha$ or $xy^\alpha \cdot x = y$ for all $x, y \in Q$
- 206 8. super middle α -symmetric if $x \cdot (y^\alpha \cdot x^\alpha) = y^\alpha$ or $(x \cdot y^\alpha) \cdot x^\alpha = y^\alpha$ for all
- 207 $x, y \in Q$

208 **Definition 2.7.** A quasigroup (Q, \cdot) is totally symmetric if any relation $xy = z$
 209 implies any other such relation can be obtained by permuting x, y and z .

210 **Definition 2.8.** [31] If a totally symmetric quasigroup (Q, \cdot) is a loop, then it is
 211 called Steiner loop.

212 **Theorem 2.1.** [31] A quasigroup (Q, \cdot) is totally symmetric if and only if it is
 213 commutative ($xy = yx$) for all $x, y \in Q$) and is right or left symmetric

214 **Theorem 2.2.** [31] A loop (Q, \cdot) is totally symmetric if and only if (Q, \cdot) is an
 215 IP loop of exponent 2.

216 **Corollary 2.1.** [31] Every T.S. quasigroup is a commutative I.M. quasigroup.

217 **Definition 2.9.** Let (Q, \cdot) be a loop. The pair $(H, \circ) = H(Q, \cdot)$ given by

$$H = A(Q) \times Q, \text{ where } A(Q) \leq AUT(Q, \cdot) \text{ such that } (\phi, x) \circ (\psi, y) = (\phi\psi, x\psi \cdot y)$$

218 for all $(\phi, x), (\psi, y) \in H$ is called the $A(H)$ - Holomorph of (Q, \cdot)

219 **Lemma 2.1.** [8] Let $(L, \cdot, /, \backslash)$ be a loop with holomorph $G(L, \cdot)$. Then, $G(L, \cdot)$
 220 is a commutative if and only if $A(L, \cdot)$ is an abelian group and $(\psi, \phi^{-1}, I_e) \in$
 221 $AATP(L, \cdot)$ for all $\phi, \psi \in A(L)$

222 **Definition 2.10.** [32] Let (Q, \cdot) be quasigroup with e_l and e_r identity elemente.
 223 (Q, \cdot) is called:

- 224 1. a left loop if $e_l \cdot x = x \ \forall x \in Q$
- 225 2. a right loop if $x \cdot e_r = x \ \forall x \in Q$
- 226 3. a loop if $e_l \cdot x = x \cdot e_r = x \ \forall x \in Q$

227 A quasigroup (Q, \cdot) , for which $e_l = e_r$ is called a loop. In more general note
 228 $e_l = e_r = e$

229

3 MAIN RESULTS

230 **3.1 Some algebraic connections between identities (2) and (3)**

231 Here, we uncovered some characterisations of the two identities of GBML:
 232 (2) and (3), and further established that they are equivalent.

233 **Lemma 3.1.** Let (Q, \cdot) be a loop. Let x, y, z be arbitrary elements in Q .

234 1. If (Q, \cdot) obeys identity (2) such that $\alpha : e \mapsto e$, then

235 (a) $(x/y) \cdot x^\alpha = x \cdot (y \setminus x^\alpha)$. 237 (c) $y^\lambda = y^\rho$.

236 (b) $y^\lambda \cdot (z^\alpha)^\rho = (z^\alpha \cdot y)^\rho$.

238 2. If (Q, \cdot) obeys identity (3) such that $\alpha : e \mapsto e$, then

239 (a) $x \cdot (z^\alpha \setminus x^\alpha) = (x/z^\alpha) \cdot x^\alpha$. 241 (c) $(z^\alpha)^\lambda = (z^\alpha)^\rho$.

240 (b) $y^\lambda \cdot (z^\alpha)^\rho = (z^\alpha \cdot y)^\lambda$.

242 3. If (Q, \cdot) obeys identity (2) such that α is bijective and $\alpha : e \mapsto e$, then

243 (a) $(x/y) \cdot x^\alpha = x \cdot (y \setminus x^\alpha)$. 245 (c) $y^\lambda = y^\rho$.

244 (b) $y^\lambda \cdot z^\rho = (z \cdot y)^\rho$.

246 4. If (Q, \cdot) obeys identity (3) such that α is bijective and $\alpha : e \mapsto e$, then

247 (a) $x \cdot (z \setminus x^\alpha) = (x/z) \cdot x^\alpha$. 249 (c) $z^\lambda = z^\rho$.

248 (b) $y^\lambda \cdot z^\rho = (z \cdot y)^\lambda$.

250 5. Let $\alpha : e \mapsto e$. Then, (Q, \cdot) obeys identity (2) if and only if (Q, \cdot) obeys
 251 identity (3) and $(x/y) \cdot x^\alpha = x \cdot (y \setminus x^\alpha)$.

252 6. Let α be bijective such that $\alpha : e \mapsto e$. Then, (Q, \cdot) obeys identity (2) if
 253 and only if (Q, \cdot) obeys identity (3).

254 **Proof.** 1. Assume that (Q, \cdot) obeys the identity (2) such that $\alpha : e \mapsto e$.

255 (a) Put $z = e$ in (2) to get $(x/y) \cdot (e^\alpha \setminus x^\alpha) = x \cdot ((e^\alpha \cdot y) \setminus x^\alpha)$ which gives
 256 $(x/y) \cdot x^\alpha = x \cdot (y \setminus x^\alpha)$.

257 (b) In (2), put $x = e$ to get $(e/y) \cdot (z^\alpha \setminus e^\alpha) = e \cdot ((z^\alpha \cdot y) \setminus e^\alpha)$ to get
 258 $y^\lambda \cdot (z^\alpha)^\rho = (z^\alpha \cdot y)^\rho$.

259 (c) In (b), put $z = e$ to get $y^\lambda = y^\rho$.

- 290 2. (132)-parastrophe of right loop is not a loop
 291 3. (132)-parastrophe of loop is a loop if and only if $|x| = 2$ for all $x \in Q$

292 **Proof.** (a) " $\circ_{(12)}$ " denotes the operation of (12)-parastrophe of Q . If (Q, \cdot)
 293 is a left loop, then $e_l \cdot x = x$ this implies that (12)-parastrophe of Q
 294 is $x \circ_{(12)} e_r = x$ for all $x \in Q$. (Q, \cdot) is right loop if $x \circ_{(12)} e_r = x \Rightarrow$
 295 (12)-parastrophe of Q is $e_l \circ_{(12)} x = x$ for all $x \in Q$. Therefore, (12)-
 296 parastrophe of Q is a loop.

297 (b) (13)-parastrophe of a left loop is given as $x \circ_{(13)} x = e_l$. This is only possible
 298 iff $|x| = 2$ for all $x \in Q$. Conversely, suppose that (13)-parastrophe of a left
 299 loop is of exponent 2, this implies that $x^\lambda = x$, then $x^\lambda \cdot x = e_l$ Also, if (Q, \cdot)
 300 is right loop, then (13)-parastrophe of Q is also loop, that is $x \circ_{(13)} e_r = x$.
 301 Thus, $x^\lambda = x^\rho = x$ Therefore, (13)-parastrophe of Q is a loop if and only
 302 if $|x| = 2$. Similar results are obtained for (c), (d) and (e).
 303 ■

304 **Theorem 3.1.** Let $(Q, \cdot, /, \backslash)$ be a generalised middle Bol loop. Then, (12)-parastrophe
 305 of Q is also a generalised middle Bol loop

306 **Proof.** Let

$$a \cdot b = x(z^\alpha y \backslash x^\alpha) \quad (4)$$

307 in equation (2) where $a = x/y \Rightarrow x = ay \xRightarrow{\text{by (12)-permutation}} y \circ_{(12)} a = x \Rightarrow a =$

308 $y \backslash^{(12)} x$. And $b = z^\alpha \backslash x^\alpha \Rightarrow z^\alpha b = x^\alpha \xRightarrow{\text{take (12)-permutation}} b \circ_{(12)} z^\alpha = x^\alpha \Rightarrow b =$

309 $x^\alpha /^{(12)} z^\alpha$.

310 Substitute for a and b into equation (4), give

$$(y \backslash^{(12)} x) \cdot (x^\alpha /^{(12)} z^\alpha) = x(z^\alpha y \backslash x^\alpha) \quad (5)$$

311 Applying (12)-permutation on equation (5), to get

$$(x^\alpha /^{(12)} z^\alpha) \circ_{(12)} (y \backslash^{(12)} x) = ((y \circ_{(12)} z^\alpha) \backslash x^\alpha) \circ_{(12)} x \quad (6)$$

312 Let

$$(y \circ_{(12)} z^\alpha) \backslash x^\alpha = c \Rightarrow (y \circ_{(12)} z^\alpha) \cdot c = x^\alpha \xRightarrow{\text{take (12)-permutation}} c \circ_{(12)} (y \circ_{(12)} z^\alpha) = x^\alpha \Rightarrow$$

$$c = x^\alpha /^{(12)} (y \circ_{(12)} z^\alpha)$$

313 Put c into equation (6) and make the substitution $x \leftrightarrow x^\alpha$, $z^\alpha \leftrightarrow y$, one obtains

$$(x /^{(12)} y) \circ_{(12)} (z^\alpha \backslash^{(12)} x^\alpha) = (x /^{(12)} (z^\alpha \circ_{(12)} y)) \circ_{(12)} x^\alpha$$

314 ■

315 **Lemma 3.3.** Let $(Q, \cdot, /, \backslash)$ be a generalised middle Bol loop. Then, the (13)–parastrophe
 316 of Q is given by

$$(x \circ_{(13)} y) /^{(13)} (x^\alpha \backslash^{(13)} z^\alpha) = x /^{(13)} [x^\alpha \backslash^{(13)} (z^\alpha /^{(13)} y)] \quad (7)$$

317 **Proof.** Let

$$a \cdot b = x(z^\alpha y \backslash x^\alpha) \quad (8)$$

318 in equation (2), where

$$a = x/y \Rightarrow x = ay \quad \underbrace{\Rightarrow}_{\text{taking (13)-permutation}} \quad a = x \circ_{(13)} y \quad (9)$$

319 and

$$b = z^\alpha \backslash x^\alpha \Rightarrow z^\alpha b = x^\alpha \quad \underbrace{\Rightarrow}_{\text{take (13)-permutation}} \quad z^\alpha = x^\alpha \circ_{(13)} b \Rightarrow x^\alpha \backslash^{(13)} z^\alpha = b \quad (10)$$

320 Let $c = z^\alpha y$ in identity (2), this implies that $z^\alpha = \underbrace{c \circ_{(13)} y}_{(13)\text{-permutation}} \Rightarrow c = z^\alpha /^{(13)} y$.

321 Also, let $d = c \backslash x^\alpha \Rightarrow c \cdot d = x^\alpha \quad \underbrace{\Rightarrow}_{\text{by taking (13)-permutation}} \quad x^\alpha \circ_{(13)} d = c \Rightarrow d =$

322 $x^\alpha \backslash^{(13)} c$. Then, substituting c into d , we have

$$d = x^\alpha \backslash^{(13)} (z^\alpha /^{(13)} y) \quad (11)$$

323 Let $s = x \cdot d \Rightarrow x = s \circ_{(13)} d \Rightarrow s = x /^{(13)} d \quad \underbrace{\Rightarrow}_{\text{substitute } d \text{ into } s}$

$$s = x /^{(13)} [x^\alpha \backslash^{(13)} (z^\alpha /^{(13)} y)] \quad (12)$$

324 Now, according to identity (2), we have $a \cdot b = s \Rightarrow \underbrace{s \circ_{(13)} b}_{(13)\text{-permutation}} = a \Rightarrow$

325 $a /^{(13)} b = s$. Substituting (9), (10) and (12) into the last equality, we have

$$(x \circ_{(13)} y) /^{(13)} (x^\alpha \backslash^{(13)} z^\alpha) = x /^{(13)} [x^\alpha \backslash^{(13)} (z^\alpha /^{(13)} y)]$$

326 which is the (13)–parastrophe of Q as required. ■

327 **Theorem 3.2.** Let $(Q, \cdot, /, \backslash)$ be a generalised middle Bol loop. Then, the fol-
 328 lowing hold in (13)–parastrophe of Q

329 1. $(L_x, L_{x^\alpha}^{-1}, L_{x^\alpha}^{-1} M_x^{-1}) \in AATP(Q, /^{(13)})$

330 2. $t^\lambda \circ_{(13)} (t \circ_{(13)} y) = y$ that is left inverse property for all $t \in Q$

- 331 3. $L_x R_{(x^\alpha)^\rho}^{-1} = \lambda J L_{x^\alpha}^{-1} M_x^{-1}$
 332 4. $L_x M_x = L_{x^\alpha}^{-1} M_{x^\alpha}^{-1}$
 333 5. $x /^{(13)}(x^\alpha)^\rho = (x \cdot_{(13)} y) /^{(13)}(x \setminus^{(13)} y)$ for all $x, y \in Q$
 334 6. $y = (y^\lambda)^\lambda$ for all $y \in Q$
 335 7. $L_x R_{(x^\alpha)^\lambda}^{-1} = \lambda J L_{(x^\alpha)^\lambda} M_x^{-1}$

336 **Proof.** 1. From equation (7) of Lemma 3.3, we have $y L_x /^{(13)} z^\alpha L_{x^\alpha}^{-1} = (z^\alpha /^{(13)} y) L_{x^\alpha}^{-1} M_x^{-1} \Rightarrow$

337
 338 $(L_x, L_{x^\alpha}^{-1}, L_{x^\alpha}^{-1} M_x^{-1}) \in AATP(Q, /^{(13)})$

339 2. Let $x = e \Rightarrow e^\alpha \rightarrow e$ the identity element in Q , in equation (7), we have

$$\begin{aligned} e y /^{(13)} z^\alpha &= e /^{(13)}(z^\alpha /^{(13)} y) \Rightarrow y /^{(13)} z^\alpha = \\ &(z^\alpha /^{(13)} y)^\lambda \Rightarrow y = (z^\alpha /^{(13)} y)^\lambda \circ_{(13)} z^\alpha \end{aligned} \quad (13)$$

340 Let $t = z^\alpha /^{(13)} y \Rightarrow z^\alpha = t \circ_{(13)} y$, put z^α and t in (13), give $y = t^\lambda \circ_{(13)}$
 341 $(t \circ_{(13)} y)$ for all $t \in Q$.

342 3. Let $z = e$ and $e^\alpha \mapsto e$ in equation (7), we have
 343 $(x \circ_{(13)} y) / (x^\alpha)^\rho = x /^{(13)}(x^\alpha \setminus^{(13)} y^\lambda) \Rightarrow y L_x R_{(x^\alpha)^\rho}^{-1} = y \lambda L_{x^\alpha}^{-1} M_x^{-1} \Rightarrow L_x R_{(x^\alpha)^\rho}^{-1} =$
 344 $\lambda J L_{x^\alpha}^{-1} M_x^{-1}$

345 4. Let $z = x$ in equation (7), we have $x \circ_{(13)} y = x /^{(13)}(x^\alpha \setminus^{(13)}(x^\alpha /^{(13)} y)) \Rightarrow$
 346 $y L_x = y M_{x^\alpha}^{-1} L_{x^\alpha}^{-1} M_x^{-1} \Rightarrow L_x M_x = L_{x^\alpha}^{-1} M_{x^\alpha}^{-1}$

347 5. Let $z = y$ and $y^\alpha \mapsto y$ in equation (7), give $x /^{(13)}(x^\alpha)^\rho = (x \circ_{(13)} y) /^{(13)}(x \setminus^{(13)} y)$.

348 6. Let $z = x = e$ in equation (7), we obtain $(y^\lambda)^\lambda = y$.

349 7. Apply 2 to equation (7), to get $(x \circ_{(13)} y) /^{(13)}((x^\alpha)^\lambda \circ_{(13)} z^\alpha) = x /^{(13)}((x^\alpha)^\lambda \circ_{(13)}$
 350 $(z^\alpha /^{(13)} y))$. Let $z^\alpha \mapsto e$ to get $x \circ_{(13)} y /^{(13)}(x^\alpha)^\lambda = x /^{(13)}((x^\alpha)^\lambda \circ_{(13)} y^\lambda) \Rightarrow$
 351 $y L_x R_{(x^\alpha)^\lambda}^{-1} = y \lambda J L_{(x^\alpha)^\lambda} M_x^{-1} \Rightarrow L_x R_{(x^\alpha)^\lambda}^{-1} = \lambda J L_{(x^\alpha)^\lambda} M_x^{-1}$
 352 ■

353 **Corollary 3.1.** In (13)–parastrophe of a generalised middle Bol loop $(Q, \cdot, /, \setminus)$,
 354 the following hold:

- 355 1. $(x^\alpha)^\rho = (x^\alpha)^\lambda \forall x \in Q$
 356 2. $x^\rho = x^\lambda \forall x \in Q$

357 **Proof.** From 7 of Theorem 3.2, we have $L_x R_{(x^\alpha)^\lambda}^{-1} = \lambda J L_{x^\alpha}^{-1} M_x^{-1}$. Recall from 3 of
 358 Theorem 3.2, $L_x R_{(x^\alpha)^\rho}^{-1} = \lambda J L_{x^\alpha}^{-1} M_x^{-1}$. This implies that $L_x R_{(x^\alpha)^\rho}^{-1} = L_x R_{(x^\alpha)^\lambda}^{-1} \Rightarrow$
 359 $R_{(x^\alpha)^\rho}^{-1} = R_{(x^\alpha)^\lambda}^{-1} \Rightarrow (x^\alpha)^\rho = (x^\alpha)^\lambda$. Since α is bijective, we have $x^\rho = x^\lambda \forall x \in Q$
 360 ■

361 **Remark 3.1.** The above Corollary shows that in (13)–parastrophe of a gener-
 362 alised middle Bol loop $(Q, \cdot, /, \backslash)$, the right and the left inverse properties coincide.
 363 So, the (13)–parastrophe satisfies IP if it is commutative. Also, if $|Q^{(13)}| = 2$,
 364 then $x^\rho = x^\lambda = x \forall x \in Q$. Thus, (13)-parastrophe of Q is a loop.

365 **Corollary 3.2.** A commutative (13)–parastrophe of a generalised middle Bol
 366 loop $(Q, \cdot, /, \backslash)$, satisfies AAIP if $|x| = 2 \forall x \in Q$

367 **Proof.** Based on the Remark (3.1), the identity (7) become $(x \circ_{(13)} y) \circ_{(13)}$
 368 $(x^\alpha)^{-1} \circ_{(13)} (z^\alpha)^{-1} = x \circ_{(13)} [(x^\alpha)^{-1} \circ_{(13)} (z^\alpha \circ_{(13)} y^{-1})]^{-1}$. Let $x = e$ to get
 369 $(e \circ_{(13)} y) \circ_{(13)} (e^\alpha)^{-1} \circ_{(13)} (z^\alpha)^{-1} = e \circ_{(13)} [(e^\alpha)^{-1} \circ_{(13)} (z^\alpha \circ_{(13)} y^{-1})]^{-1}$, then
 370 $y \circ_{(13)} (z^\alpha)^{-1} = (z^\alpha \circ_{(13)} y^{-1})^{-1} \Rightarrow y^{-1} \circ_{(13)} (z^\alpha)^{-1} = (z^\alpha \circ_{(13)} y)^{-1}$ ■

371 **Corollary 3.3.** A commutative (13)–parastrophe of a generalised middle Bol
 372 loop $(Q, \cdot, /, \backslash)$ is a Steiner loop if it is a loop of exponent two.

373 **Proof.** This is a consequence of 2 of theorem 3.2 and the Corollary 3.1. ■

374 **Theorem 3.3.** A commutative (13)–parastrophe, of exponent two, of a gener-
 375 alised middle Bol loop $(Q, \cdot, /, \backslash)$ is a Moufang loop.

376 **Proof.** From Remark (3.1), we have the identity (7) to be $(x \circ_{(13)} y) \circ_{(13)} (x^\alpha)^{-1} \circ_{(13)}$
 377 $(z^\alpha)^{-1} = x \circ_{(13)} [(x^\alpha)^{-1} \circ_{(13)} (z^\alpha \circ_{(13)} y^{-1})]^{-1}$. Since $|Q^{(13)}| = 2$, we have
 378 $(x \circ_{(13)} y) \circ_{(13)} (x^\alpha \circ_{(13)} z^\alpha) = x \circ_{(13)} [x^\alpha \circ_{(13)} (z^\alpha \circ_{(13)} y)] \Rightarrow z^\alpha L_{x^\alpha} L_{xy} =$
 379 $z^\alpha R_y L_{x^\alpha} L_x \Rightarrow z^\alpha R_{x^\alpha} L_{xy} = z^\alpha L_y L_{x^\alpha} L_x \Rightarrow (x \circ_{(13)} y) \circ_{(13)} (z^\alpha \circ_{(13)} x^\alpha) =$
 380 $x \circ_{(13)} ((y \circ_{(13)} z^\alpha) \circ_{(13)} x^\alpha)$ ■

381 **Corollary 3.4.** In (13)–parastrophe, of exponent two, of a generalised middle
 382 Bol loop $(Q, \cdot, /, \backslash)$ is a GMBL

383 **Proof.** Follow from Theorem 3.3, we have $(x \circ_{(13)} y) \circ_{(13)} [(x^\alpha)^{-1} \circ_{(13)} (z^\alpha)]^{-1} =$
 384 $x \circ_{(13)} [(x^\alpha)^{-1} \circ_{(13)} (z^\alpha \circ_{(13)} y^{-1})]^{-1}$. Use $y^{-1} = y$ and Corollary 3.2 to get
 385 $(x \circ_{(13)} y^{-1}) \circ_{(13)} ((z^\alpha)^{-1} \circ_{(13)} x^\alpha) = x \circ_{(13)} [(z^\alpha \circ_{(13)} y)]^{-1} \circ_{(13)} x^\alpha \Rightarrow (x /^{(13)} y) \circ_{(13)}$
 386 $(z^\alpha \backslash^{(13)} x^\alpha) = x \circ_{(13)} [(z^\alpha \circ_{(13)} y) \backslash^{(13)} x^\alpha]$
 387 ■

388 **Lemma 3.4.** Let $(Q, \cdot, /, \backslash)$ be a generalised middle Bol loop. Then, the (23)–parastrophe
 389 of Q is given by

$$(y /^{(23)} x) \backslash^{(23)} (z^\alpha \circ_{(23)} x^\alpha) = x \backslash^{(23)} [(z^\alpha \backslash^{(23)} y) \circ_{(23)} x^\alpha] \quad (14)$$

390 **Proof.** Let

$$a \cdot b = x(z^\alpha y \setminus x^\alpha) \quad (15)$$

391 in an identity (2), where

$$a = x/y \Rightarrow x = a \cdot y \quad \underbrace{\Rightarrow}_{(23)\text{-permutation}} \quad y = a \circ_{(23)} x \Rightarrow a = y / {}^{(23)}x \quad (16)$$

392 and

$$b = z^\alpha \setminus x^\alpha \quad \underbrace{\Rightarrow}_{(23)\text{-permutation}} \quad z^\alpha \circ_{(23)} b = x^\alpha \Rightarrow z^\alpha \circ_{(23)} x^\alpha = b \quad (17)$$

393 Let $c = z^\alpha y$ in identity (2), then $\underbrace{z^\alpha \circ_{(23)} c}_{(23)\text{-permutation}} = y \Rightarrow c = z^\alpha \setminus {}^{(23)}y$. Let

394 $d = c \setminus x^\alpha \Rightarrow c \circ_{(23)} d = x^\alpha \Rightarrow c \circ_{(23)} x^\alpha = d$, put c into d to get

$$d = (z^\alpha \setminus {}^{(23)}y) \circ_{(23)} x^\alpha. \quad (18)$$

395 Also, let $t = x \cdot d \quad \underbrace{\Rightarrow}_{(23)\text{-permutation}} \quad x \circ_{(23)} t = d \Rightarrow t = x \setminus {}^{(23)}d$. Substitute d into

396 t

$$t = x \setminus {}^{(23)}[(z^\alpha \setminus {}^{(23)}y) \circ_{(23)} x^\alpha] \quad (19)$$

397 Now, going by the identity (2), we have $a \cdot b = t \quad \underbrace{\Rightarrow}_{(23)\text{-permutation}} \quad a \circ_{(23)} t = b \Rightarrow$

398 $a \setminus {}^{(23)}b = t$. Then, substituting equations (16), (17) and (19) in the equality
399 $a \setminus {}^{(23)}b = t$, gives

$$(y / {}^{(23)}x) \setminus {}^{(23)}(z^\alpha \circ_{(23)} x^\alpha) = x \setminus {}^{(23)}[(z^\alpha \setminus {}^{(23)}y) \circ_{(23)} x^\alpha] \quad (20)$$

400 which is the (23)–parastrophe of Q . ■

401 **Theorem 3.4.** Let $(Q, \cdot, /, \setminus)$ be a generalised middle Bol loop. Then, the fol-
402 lowing holds in (23)–parastrophe of Q

- 403 1. $(L_x^{-1}, R_{x^\alpha}, R_{x^\alpha} L_x^{-1}) \in AATP(Q, \setminus {}^{(23)})$ for all $x \in Q$
- 404 2. $(z \circ_{(23)} t) \circ_{(23)} t = z$ for all $z, t \in Q$
- 405 3. if $Q^{(23)}$ is middle symmetric then, $x \circ_{(23)} (z^\alpha \circ_{(23)} x^\alpha) = (x \circ_{(23)} z^\alpha) \circ_{(23)} x^\alpha$
406 that is, super α –elastic
- 407 4. $R_x^{-1} M_{x^\alpha} = R_{x^\alpha} L_x^{-1}$
- 408 5. $\rho J R_{x^\alpha} L_x^{-1} = R_{x^\alpha} L_{x^\lambda}^{-1}$

409 6. $\rho J R_{x^\alpha}^{-1} M_{x^\alpha} = R_{x^\alpha} L_{x^\lambda}^{-1}$

410 **Proof.** 1. this follows from equation (14),

$$y R_x^{-1} \setminus^{(23)} z^\alpha R_{x^\alpha} = (z^\alpha \setminus^{(23)} y) R_{x^\alpha} L_x^{-1} \Rightarrow (R_x^{-1}, R_{x^\alpha}, R_{x^\alpha} L_x^{-1}) \in AATP(Q, \setminus)$$

411 2. Put $x = e$ such that $e^\alpha \mapsto e$ is the identity map in (14), give $y \setminus^{(23)} z^\alpha =$
 412 $z^\alpha \setminus^{(23)} y \Rightarrow y \circ_{(23)} (z^\alpha \setminus^{(23)} y) = z^\alpha$. Let $t = z^\alpha \setminus^{(23)} y \Rightarrow z^\alpha \circ_{(23)} t = y$. Put y
 413 into the last equality to get $(z^\alpha \circ_{(23)} t) \circ_{(23)} t = z^\alpha$ for any $t \in Q$.

414 3. Put $y = x$ in (14), we have

$$\begin{aligned} z^\alpha \circ_{(23)} x^\alpha &= x \setminus^{(23)} [(z^\alpha \setminus^{(23)} x) \circ_{(23)} x^\alpha] \Rightarrow \\ x^\alpha \circ_{(23)} (z^\alpha \circ_{(23)} x) &= (z^\alpha \setminus^{(23)} x) \circ_{(23)} x^\alpha \Rightarrow \\ z^\alpha R_{x^\alpha} L_x &= z^\alpha M_x R_{x^\alpha} \quad \underbrace{\Rightarrow}_{\text{Use middle symmetric as } L_x = M_x \text{ to get}} \quad z^\alpha R_{x^\alpha} L_x = z^\alpha L_x R_{x^\alpha} \end{aligned}$$

415 or $x \circ_{(23)} (z^\alpha \circ_{(23)} x^\alpha) = (x \circ_{(23)} z^\alpha) \circ_{(23)} x^\alpha$

416 4. Put $z = e$ and $e^\alpha \mapsto e$, the identity element in (14), we have
 417 $(y /^{(23)} x) \setminus^{(23)} x^\alpha = x \setminus^{(23)} (y \circ_{(23)} x^\alpha) \Rightarrow y R_x^{-1} M_{x^\alpha} = y R_{x^\alpha} L_x^{-1} \Rightarrow R_x^{-1} M_{x^\alpha} =$
 418 $R_{x^\alpha} L_x^{-1}$

419 5. $y = e$ in (14), we have

$$\begin{aligned} x^\lambda \setminus^{(23)} (z^\alpha \circ_{(23)} x^\alpha) &= x \setminus^{(23)} ((z^\alpha)^\rho \circ_{(23)} x^\alpha) \Rightarrow \\ z^\alpha \rho J R_{x^\alpha} L_x^{-1} &= z^\alpha R_{x^\alpha} L_{x^\lambda}^{-1} \Rightarrow \rho J R_{x^\alpha} L_x^{-1} = R_{x^\alpha} L_{x^\lambda}^{-1} \end{aligned}$$

420 6. Use 4 and 5.

421 ■

422 **Corollary 3.5.** A commutative (23)–parastrophe of a generalised middle Bol
 423 loop $(Q, \cdot, /, \setminus)$ is totally symmetric.

424 **Proof.** This is a consequence of the right symmetric property 2 of Theorem 3.4.

425 ■

426 **Theorem 3.5.** Let the (23)–parastrophe of a generalised middle Bol loop $(Q, \cdot, /, \setminus)$
 427 be commutative and of exponent two, then $L_{x^\alpha} L_x = R_x R_{x^\alpha}$ for all $x \in Q$.

428 **Proof.** Recall (6) in Theorem 3.4, we have $\rho J R_x^{-1} M_{x^\alpha} = R_{x^\alpha} L_{x^\lambda}^{-1}$. Since $Q^{(23)}$
 429 is commutative, then it implies that it has a middle symmetric property as $L_x =$

430 M_x . Applying the middle symmetric identity gives $\rho J R_x^{-1} L_{x^\alpha} = R_{x^\alpha} L_{x^\lambda}^{-1}$. Then,
431 for all $t \in Q$, we have

$$\begin{aligned} t^\rho R_x^{-1} L_{x^\alpha} &= t R_{x^\alpha} L_{x^\lambda}^{-1} \Rightarrow x^\alpha \circ_{(23)} (t^\rho/x) = x^\lambda \setminus^{(23)} (t \circ_{(23)} x^\alpha) \Rightarrow \\ &x^\lambda \circ_{(23)} [x^\alpha \circ_{(23)} (t^\rho/x)] = t \circ_{(23)} x^\alpha \end{aligned}$$

432 Let $t^\rho /^{(23)} x = s \Rightarrow t^\rho = s \circ_{(23)} x$. Then, $x^\lambda \circ_{(23)} (x^\alpha \circ_{(23)} s) = (s \circ_{(23)} x) \circ_{(23)} x^\alpha$.
433 Using the fact that $|Q^{(23)}| = 2$ for all $x \in Q$, one obtains $s L_{x^\alpha} L_x = s R_x R_{x^\alpha} \Rightarrow$
434 $L_{x^\alpha} L_x = R_x R_{x^\alpha}$ for all $x \in Q$ ■

435 **Corollary 3.6.** If (23)–parastrophe of a generalised middle Bol loop $(Q, \cdot, /, \setminus)$
436 is commutative and $x^\alpha \mapsto x$, then $L_x^2 = R_x^2$ for all $x \in Q$.

437 **Proof.** Consequence of Theorem 3.5. ■

438 **Lemma 3.5.** Let $(Q, \cdot, /, \setminus)$ be a generalised middle Bol. Then, the (123)–parastrophe
439 of Q is given by

$$(z^\alpha /^{(123)} x^\alpha) \setminus^{(123)} (y \circ_{(123)} x) = [(y \setminus^{(123)} z^\alpha) /^{(123)} x^\alpha] \setminus^{(123)} x \quad (21)$$

440 **Proof.** Let $a \cdot b = x \cdot (z^\alpha y \setminus x^\alpha)$ in equation (2) where

$$a = x/y \Rightarrow a \cdot y = x \quad \underbrace{\Rightarrow}_{(123)\text{-permutation}} \quad y \circ_{(123)} x = a \quad (22)$$

$$441 \quad b = z^\alpha \setminus x^\alpha \Rightarrow z^\alpha \circ b = x^\alpha \quad \underbrace{\Rightarrow}_{(123)\text{-permutation}} \quad b \circ_{(123)} x^\alpha = z^\alpha \Rightarrow b = z^\alpha /^{(123)} x^\alpha \quad (23)$$

442 Let $c = z^\alpha \cdot y$ in equation (2), then, we have $y \circ_{(123)} c = z^\alpha \quad \underbrace{\Rightarrow}_{(123)\text{-permutation}} \quad c =$

443 $y \setminus^{(123)} z^\alpha$. Also, let $d = c \setminus x^\alpha \Rightarrow c \cdot d = x^\alpha \Rightarrow d \circ_{(123)} x^\alpha = c \Rightarrow d = c /^{(123)} x^\alpha$.
444 Substitute c into d , give

$$d = (y \setminus^{(123)} z^\alpha) /^{(123)} x^\alpha \quad (24)$$

445 Next, let $t = x \cdot d \quad \underbrace{\Rightarrow}_{(123)\text{-permutation}} \quad d \circ_{(123)} t = x \Rightarrow t = d \setminus^{(123)} x$. Substitute (24)

446 into t give

$$t = [(y \setminus^{(123)} z^\alpha) /^{(123)} x^\alpha] \setminus^{(123)} x \quad (25)$$

447 Going by the identity (2), we have $a \cdot b = t \quad \underbrace{\Rightarrow}_{(123)\text{-permutation}} \quad b \circ_{(123)} t = a \Rightarrow$

448 $b \setminus^{(123)} a = t$. Substitute (22), (23) and (25) into the equality $b \setminus^{(123)} a = t$,
449 gives the (123)–parastrophe as

$$(z^\alpha /^{(123)} x^\alpha) \setminus^{(123)} (y \circ_{(123)} x) = [(y \setminus^{(123)} z^\alpha) /^{(123)} x^\alpha] \setminus^{(123)} x$$

451 **Theorem 3.6.** Let $(Q, \cdot, /, \backslash)$ be a generalised middle Bol loop. Then, the fol-
 452 lowing hold in (123) -parastrophe of Q

- 453 1. $(L_x^{-1}, R_x, R_x^{-1}M_x) \in AATP(Q, \backslash^{(123)})$
- 454 2. $(y \circ_{(123)} t) \circ_{(123)} t^\rho = y$, i.e right inverse property
- 455 3. $(z^\alpha /^{(123)} x^\alpha)[(x^\alpha)^\lambda \backslash^{(123)} x] = z^\alpha \circ_{(123)} x$
- 456 4. $R_x L_{(x^\alpha)^\lambda}^{-1} = \rho J R_{x^\alpha}^{-1} M_x$
- 457 5. $R_x M_x^{-1} = M_{x^\alpha} R_{x^\alpha}^{-1}$
- 458 6. $(x \circ_{(123)} t) \circ_{(123)} x = (x \backslash^{(123)} t) \backslash^{(123)} x$ for all $x, t \in Q$

459 **Proof.** 1. From equation (21), we have

$$z^\alpha R_{x^\alpha}^{-1} \backslash^{(123)} y R_x = (y \backslash^{(123)} z^\alpha) R_{x^\alpha}^{-1} M_x \Rightarrow \\ (R_{x^\alpha}^{-1}, R_x, R_{x^\alpha}^{-1} M_x) \in AATP(Q, \backslash^{(123)})$$

460 2. Let $x^\alpha \mapsto x$ and put $x = e$, the identity element in equation (21), we have

$$((z^\alpha /^{(123)} e^\alpha) \backslash^{(123)} (y \circ_{(123)} e)) = ((y \backslash^{(123)} z^\alpha) /^{(123)} e) \backslash^{(123)} e^\alpha \Rightarrow z^\alpha \backslash^{(123)} y = \\ (y \backslash^{(123)} z^\alpha)^\rho \Rightarrow z^\alpha \circ_{(123)} (y \backslash^{(123)} z^\alpha)^\rho = y$$

461 Let $t = y \backslash^{(123)} z^\alpha \Rightarrow y \circ_{(123)} t = z^\alpha$ for any $t \in Q$, this implies that $(y \circ_{(123)}$
 462 $t) \circ_{(123)} t^\rho = y$.

463 3. Set $y = z^\alpha$ in equation (21), we have $(z^\alpha /^{(123)} x^\alpha) \backslash^{(123)} (z^\alpha \circ_{(123)} x) =$
 464 $(x^\alpha)^\lambda \backslash^{(123)} x \Rightarrow (z^\alpha /^{(123)} x^\alpha)[(x^\alpha)^\lambda \backslash^{(123)} x] = z^\alpha \circ_{(123)} x$

465 4. Put $z \rightarrow e$ in equation (21), to get $(x^\alpha)^\lambda \backslash^{(123)} y \circ_{(123)} x = (y^\rho /^{(123)} x^\alpha) \backslash^{(123)} x \Rightarrow$
 466 $y R_x L_{(x^\alpha)^\lambda}^{-1} = y \rho J R_{x^\alpha}^{-1} M_x \Rightarrow R_x L_{(x^\alpha)^\lambda}^{-1} = \rho J R_{x^\alpha}^{-1} M_x$

467 5. Set $z = x$ in equation (21), give $y \circ_{(123)} x = ((y \backslash^{(123)} x^\alpha) /^{(123)} x^\alpha) \backslash^{(123)} x \Rightarrow$
 468 $y R_x = y M_{x^\alpha} R_{x^\alpha}^{-1} M_x \Rightarrow R_x M_x^{-1} = M_{x^\alpha} R_{x^\alpha}^{-1}$

469 ■

470 **Corollary 3.7.** A commutative (123) -parastrophe of a generalised middle Bol
 471 loop $(Q, \cdot, /, \backslash)$ has an inverse property.

472 **Proof.** This is a consequence of 2 of Theorem 3.6. ■

473 **Corollary 3.8.** A commutative (123) -parastrophe of a generalised middle Bol
 474 loop $(Q, \cdot, /, \backslash)$ has AAIP if $|Q^{(123)}| = 2$

475 **Proof.** Applying Corollary 3.7 to (21) $(z^\alpha \circ_{(123)} (x^\alpha)^{-1})^{-1} \circ_{(123)} (y \circ_{(123)} x) =$
 476 $[(y^{-1} \circ_{(123)} z^\alpha) \circ_{(123)} (x^\alpha)^{-1}]^{-1} \circ_{(123)} x$. Put $x = e$ and $y = y^{-1}$ to get $(z^\alpha)^{-1} \circ_{(123)}$
 477 $y^{-1} = (y^{-1} \circ_{(123)} z^\alpha)^{-1}$ ■

478 **Corollary 3.9.** A commutative (123)–parastrophe, of exponent 2, of a gener-
 479 alised middle Bol loop $(Q, \cdot, /, \backslash)$ is Steiner loop.

480 **Proof.** Follows from Corollary 3.7. ■

481 **Theorem 3.7.** Let $Q^{(123)}$ be a commutative (123)–parastrophe of a generalised
 482 middle Bol loop $(Q, \cdot, /, \backslash)$ of exponent two, then $Q^{(123)}$ is a Moufang loop.

483 **Proof.** Using the Corollary 3.7 on identity (21), we have $(z^\alpha \circ_{(123)} (x^\alpha)^{-1})^{-1} \circ_{(123)}$
 484 $(y \circ_{(123)} x) = [(y^{-1} \circ_{(123)} z^\alpha) \circ_{(123)} (x^\alpha)^{-1}]^{-1} \circ_{(123)} x$. Since $|Q^{(123)}| = 2$, we have
 485 $(z^\alpha \circ_{(123)} x^\alpha) \circ_{(123)} (y \circ_{(123)} x) = [(y \circ_{(123)} z^\alpha) \circ_{(123)} (x^\alpha)] \circ_{(123)} x \Rightarrow z^\alpha L_{x^\alpha} R_{yx} =$
 486 $z^\alpha L_y R_{x^\alpha} R_x \Rightarrow z^\alpha L_{x^\alpha} R_{yx} = z^\alpha L_y L_{x^\alpha} R_x \Rightarrow (x^\alpha \circ_{(123)} z^\alpha) \circ (y \circ_{(123)} x) = (x^\alpha \circ_{(123)}$
 487 $(z^\alpha \circ_{(123)} y)) \circ_{(123)} x \Rightarrow (x^\alpha \circ_{(123)} z^\alpha) \circ (y \circ_{(123)} x) = x^\alpha \circ_{(123)} ((z^\alpha \circ_{(123)} y) \circ_{(123)} x)$
 488 ■

489 **Corollary 3.10.** A commutative (123)–parastrophe of a generalised middle Bol
 490 loop $(Q, \cdot, /, \backslash)$ is a GMBL of exponent two.

491 **Proof.** Follow from Corollaries 3.7 and 3.8 and (21), we get $(z^\alpha \circ_{(123)} (x^\alpha)^{-1})^{-1} \circ_{(123)}$
 492 $(y \circ_{(123)} x) = [(y^{-1} \circ_{(123)} z^\alpha) \circ_{(123)} (x^\alpha)^{-1}]^{-1} \circ_{(123)} x$. So, use $y^{-1} = y$ and take
 493 the following steps: $x \leftrightarrow x^\alpha$, $z^\alpha \leftrightarrow y$, one obtains

$$(x /^{(12)} y) \circ_{(12)} (z^\alpha \backslash^{(12)} x^\alpha) = (x /^{(12)} (z^\alpha \circ_{(12)} y)) \circ_{(12)} x^\alpha$$

494 which is the same as (3) ■

495 **Lemma 3.6.** Let $(Q, \cdot, /, \backslash)$ be a generalised middle Bol loop. Then, the (132)–parastrophe
 496 of Q is given by

$$(x^\alpha \circ_{(132)} z^\alpha) /^{(132)} (x \backslash^{(132)} y) = [x^\alpha \circ_{(132)} (y /^{(132)} z^\alpha)] /^{(132)} x \quad (26)$$

497 **Proof.** Let $a \cdot b = x \cdot (z^\alpha y \backslash x^\alpha)$ in equation (2) where

$$x / y = a \Rightarrow x = a \cdot y \quad \underbrace{\Rightarrow}_{(132)\text{-permutation}} \quad x \circ_{132} a = y \Rightarrow a = x \backslash^{(132)} y \quad (27)$$

498 and

$$z^\alpha \backslash x^\alpha = b \Rightarrow z^\alpha \cdot b = x^\alpha \quad \underbrace{\Rightarrow}_{(132)\text{-permutation}} \quad x^\alpha \circ_{(132)} z^\alpha = b \quad (28)$$

499 Let $c = z^\alpha \cdot y \xRightarrow{(132)\text{-permutation}} c \circ_{(132)} z^\alpha = y \Rightarrow c = y /^{(132)} z^\alpha$. Also, let $d =$
 500 $c \setminus x^\alpha \Rightarrow c \cdot d = x^\alpha \xRightarrow{(132)\text{-permutation}} x^\alpha \circ_{(132)} c = d$. Substitute c into d to get
 501 $d = x^\alpha \circ_{(132)} (y /^{(132)} z^\alpha)$. Let $t = x \cdot d \xRightarrow{\text{taking (132)-permutation}} t \circ_{(132)} x = d \Rightarrow t =$
 502 $d /^{(132)} x$. Hence, putting d into t , we have

$$t = [x^\alpha \circ_{(132)} (y /^{(132)} z^\alpha)] /^{(132)} x \quad (29)$$

503 Now, going by the identity (2), we have $a \cdot b = t \xRightarrow{\text{taking (132)-permutation}} t \circ_{(132)} a =$
 504 $b \Rightarrow b /^{(132)} a = t$. Substitute equations (27), (28) and (29) into the equality
 505 $b /^{(132)} a = t$, we have

$$(x^\alpha \circ_{(132)} z^\alpha) /^{(132)} (x \setminus^{(132)} y) = [x^\alpha \circ_{(132)} (y /^{(132)} z^\alpha)] /^{(132)} x$$

506 which is the (132)–parastrophe of Q . ■

507 **Theorem 3.8.** Let $(Q, \cdot, /, \setminus)$ be a generalised middle Bol loop. Then, the fol-
 508 lowing holds in (132)–parastrophe of Q

- 509 1. $(L_{x^\alpha}, L_x^{-1}, L_{x^\alpha} R_x^{-1}) \in AATP(Q, /^{(132)})$ for all $x \in Q$
- 510 2. $z^\alpha = t \circ_{132} (t \circ_{132} z^\alpha)$ i.e α –left symmetric property
- 511 3. $(x^\alpha \circ_{(132)} z^\alpha) \circ_{(132)} x = x^\alpha \circ_{(132)} (x /^{(132)} z^\alpha)$ or $M_x^{-1} L_{x^\alpha} = L_{x^\alpha} R_x$
- 512 4. $L_{x^\alpha} R_{x^\rho}^{-1} = \lambda J L_{x^\alpha} R_x^{-1}$
- 513 5. $L_x^{-1} M_{x^\alpha}^{-1} = L_{x^\alpha} R_x^{-1}$

514 **Proof.** 1. From equation (26), we have $z^\alpha L_{x^\alpha} /^{(132)} y L_x^{-1} = (y /^{(132)} z^\alpha) L_{x^\alpha} R_x^{-1} \Rightarrow$
 515 $(L_{x^\alpha}, L_x^{-1}, L_{x^\alpha} R_x^{-1}) \in AATP(Q, /^{(132)})$ for all $x \in Q$

516 2. Let $x^\alpha \mapsto e$ in (26), give $z^\alpha /^{(132)} y = y /^{(132)} z^\alpha$, by setting $t = y /^{(132)} z^\alpha \Rightarrow$
 517 $y = (z^\alpha \circ_{(132)} t) \Rightarrow z^\alpha = t \circ_{(132)} (t \circ_{(132)} z^\alpha)$

518 3. Put $y = x$ in (26), to get $(x^\alpha \circ_{(132)} z^\alpha) \circ_{(132)} x = x^\alpha \circ_{(132)} (x /^{(132)} z^\alpha) \Rightarrow$
 519 $z^\alpha M_x^{-1} L_{x^\alpha} = z^\alpha L_{x^\alpha} R_x \Rightarrow M_x^{-1} L_{x^\alpha} = L_{x^\alpha} R_x$ for all $x \in Q$

520 4. Put $y = e$ in (26), we have $(x^\alpha \circ_{(132)} z^\alpha) /^{(132)} x^\rho = (x^\alpha \circ_{(132)} (z^\alpha)^\lambda) /^{(132)} x \Rightarrow$
 521 $z^\alpha L_{x^\alpha} R_{x^\rho}^{-1} = (z^\alpha)^\lambda J L_{x^\alpha} R_x^{-1} \Rightarrow L_{x^\alpha} R_{x^\rho}^{-1} = \lambda J L_{x^\alpha} R_x^{-1}$.

522 5. Put $z = e$, we have $x /^{(132)} (x \setminus^{(132)} y) = (x \circ_{(132)} y) /^{(132)} x \Rightarrow y L_x M_x^{-1} =$
 523 $y L_x R_x^{-1} \Rightarrow L_x^{-1} M_x^{-1} = L_{x^\alpha} R_x^{-1}$ for all $x \in Q$.

524 ■

525 **Corollary 3.11.** Let $(Q, \cdot, /, \backslash)$ be a generalised middle Bol loop. Then, a com-
526 mutative (132)–parastrophe of Q is totally symmetric.

527 *Proof.* This is a consequence, of 2, of Theorem 3.8. ■

528 3.3 Holomorphic Structure of Generalised Middle Bol Loop

529 **Theorem 3.9.** $(Q, \cdot, /, \backslash)$ is a generalised middle Bol loop if and only if
530 $(JM_x^{-1}, JM_{x^\alpha}, JM_{x^\alpha}L_x)$ is an autotopism.

531 *Proof.* Suppose (Q, \cdot) is a generalised middle Bol loop, then

$$\begin{aligned} x(y^\alpha z \backslash x^\alpha) &= (x/z)(y^\alpha \backslash x^\alpha) \Leftrightarrow zM_x^{-1} \cdot y^\alpha M_{x^\alpha} = (y^\alpha \cdot z)M_{x^\alpha}L_x \\ &\Leftrightarrow zM_x^{-1} \cdot y^\alpha M_{x^\alpha} = (zJ \cdot y^\alpha J)JM_{x^\alpha}L_x \\ &\Leftrightarrow zJM_x^{-1} \cdot y^\alpha JM_{x^\alpha} = (z \cdot y^\alpha)JM_{x^\alpha}L_x \end{aligned}$$

532 Thus, $(JM_x^{-1}, JM_{x^\alpha}, JM_{x^\alpha}L_x) \in ATP(Q, \cdot)$ ■

533 **Theorem 3.10.** Let $(Q, \cdot, /, \backslash)$ be a loop with holomorph $(H(Q), *)$. Then,
534 $(H(Q), *)$ is a generalised middle Bol loop if and only if $(x\tau) \cdot (y \cdot z^\alpha \tau) \backslash x^\alpha =$
535 $(x^\alpha \tau / z^\alpha \tau) \cdot (y \backslash x)$ for all $x, y, z \in Q, \tau \in A(Q)$.

536 *Proof.* We need to show the necessary and sufficient condition for the holomorph
537 of a generalised middle Bol loop to be a generalised middle Bol loop.

$$(x^\alpha / z^\alpha)(y \backslash x) = x((y \cdot z^\alpha) \backslash x^\alpha) \tag{30}$$

538

$$\text{Let } (\phi, x) * (\psi, y) = (\theta, z) \Rightarrow (\phi, x) = (\theta, z) / (\psi, y), \text{ so ,} \tag{31}$$

$$(\phi\psi, x\psi \cdot y) = (\theta, z)$$

$$\Rightarrow \phi = \theta\psi^{-1}, x = (z/y)\psi^{-1}.$$

$$\Rightarrow (\theta, z) / (\psi, y) = (\theta\psi^{-1}, (z/y)\phi^{-1}) = (\phi, x). \tag{32}$$

$$\text{Also, } (\phi, x) * (\psi, y) = (\theta, z) \Rightarrow (\psi, y) = (\phi, x) \backslash (\theta, z).$$

$$\text{Thus, } (\phi\psi, x\psi \cdot y) = (\theta, z) \Rightarrow \psi = \phi^{-1}\theta, y = (x\phi^{-1}\theta) \backslash z$$

$$\Rightarrow (\psi, y) = (\phi^{-1}\theta, (x\phi^{-1}\theta) \backslash z) = (\phi, x) \backslash (\theta, z) \tag{33}$$

539

$$\begin{aligned}
 ((\phi, x)/(\psi, y)) * ((\theta, z^\alpha)\backslash(\phi, x^\alpha)) &= (\phi, x) * [((\psi, y) * (\theta, z^\alpha))\backslash(\phi, x^\alpha)] \\
 \text{RHS} &= (\phi, x) * [((\psi, y) * (\theta, z^\alpha))\backslash(\phi, x^\alpha)] \\
 &= (\phi, x) * \left((\psi\theta)^{-1}\phi, (y\theta \cdot z^\alpha)\theta^{-1}\psi^{-1}\phi\backslash x^\alpha \right) \\
 &= (\phi, x) * \left((\psi\theta)^{-1}\phi, y\psi^{-1}\phi \cdot z^\alpha\theta^{-1}\psi^{-1}\phi\backslash x^\alpha \right) \\
 &= (\phi\theta^{-1}\psi^{-1}\phi, (x\theta^{-1}\psi^{-1}\phi) \cdot (y\psi^{-1}\phi \cdot z^\alpha\theta^{-1}\psi^{-1}\phi\backslash x^\alpha)) \\
 \text{LHS} &= ((\phi, x)/(\psi, y)) * ((\theta, z^\alpha)\backslash(\phi, x^\alpha)) \\
 &= (\phi\theta^{-1}, (x^\alpha/z^\alpha)\theta^{-1}) * (\psi^{-1}\phi, (y\psi^{-1}\phi)\backslash x) \\
 &= (\phi\theta^{-1}\psi^{-1}\phi, (x^\alpha/z^\alpha)\theta^{-1}\psi^{-1}\phi \cdot (y\psi^{-1}\phi)\backslash x) \\
 \text{RHS} &= \text{LHS} \\
 \Leftrightarrow ((x\theta^{-1}\psi^{-1}\phi) \cdot (y\psi^{-1}\phi \cdot z^\alpha\theta^{-1}\psi^{-1}\phi\backslash x^\alpha)) &= ((x^\alpha/z^\alpha)\theta^{-1}\psi^{-1}\phi \cdot (y\psi^{-1}\phi)\backslash x)
 \end{aligned}$$

 540 Let $\tau = \theta^{-1}\psi^{-1}\phi$, then $(x\tau) \cdot (y\theta\tau \cdot z^\alpha\tau)\backslash x^\alpha = (x^\alpha/z^\alpha)\tau \cdot (y\theta\tau)\backslash x$.

 541 Replacing y by $y(\theta\tau)^{-1}$, we have

$$\begin{aligned}
 (x\tau) \cdot (y(\theta\tau)^{-1}\theta\tau \cdot z^\alpha\tau)\backslash x^\alpha &= (x^\alpha/z^\alpha)\tau \cdot (y(\theta\tau)^{-1}\theta\tau)\backslash x \\
 \Leftrightarrow (x\tau) \cdot (y \cdot z^\alpha\tau)\backslash x^\alpha &= (x^\alpha/z^\alpha)\tau \cdot (y\backslash x) \\
 \Leftrightarrow (x\tau) \cdot (y \cdot z^\alpha\tau)\backslash x^\alpha &= (x^\alpha\tau/z^\alpha\tau) \cdot (y\backslash x)
 \end{aligned}$$

 542 ■

543 **Corollary 3.12.** Let $(Q, \cdot, /, \backslash)$ be a loop with holomorph $H(Q, \cdot)$. Then, $H(Q, \cdot)$
 544 is a commutative generalised middle Bol loop if and only if $(\tau^{-1}M_{x^\alpha}^{-1}\tau, M_x, M_x^\alpha L_{x\tau}) \in$
 545 $ATP(Q, \cdot)$

 546 **Proof.** From the consequence of Theorem 3.10, we have

$$z^\alpha\tau^{-1}M_{x^\alpha}^{-1}\tau \cdot yM_x = (y \cdot z^\alpha)M_x^\alpha L_{x\tau} \quad (34)$$

$$\Leftrightarrow (\tau^{-1}M_{x^\alpha}^{-1}\tau, M_x, M_x^\alpha L_{x\tau}) \in ATP(Q, \cdot) \quad (35)$$

 547 ■

548 **Theorem 3.11.** Let $(Q, \cdot, /, \backslash)$ be a commutative generalised middle Bol loop
 549 with a holomorph $(H, *) = H(Q, \cdot)$. If :

- 550 1. $\tau = \tau(a, b) = R_{(b\backslash a)}R_b^{-1}$ for each $\tau \in A(Q)$ and for any $a, b \in Q$
 551 2. $M_x^{-1}R_{s\tau} = R_sR_x^{-1}R_{x\tau}$ for all $s, x \in Q$ and $\tau \in A(Q)$, then $H(Q, \cdot)$ is a
 552 GMBL.

553 **Proof.** From Corollary 3.12, observe that $(\tau^{-1}M_x^{-1}\tau, M_x, M_x^\alpha L_{x\tau}) = (\tau^{-1}, M_x, I_e) \circ$
 554 $(M_x^{-1}, M_x, M_x^\alpha L_x) \circ (\tau, M_x^{-1}, L_x^{-1}L_{x\tau})$. Where I_e is an identity map.

$$\begin{aligned} \text{Consider one hand } , (\tau^{-1}, M_x, I_e) \in ATP(Q, \cdot) &\Leftrightarrow a\tau^{-1} \cdot bM_x = ab \\ &\Leftrightarrow a\tau^{-1} \cdot b \setminus x = ab \\ &\Leftrightarrow a\tau^{-1}R_{b \setminus x} = aR_b \\ \Leftrightarrow \tau^{-1}R_{b \setminus x} = R_b &\Leftrightarrow \tau = \tau(a, b) = R_{b \setminus a}R_b^{-1} \end{aligned}$$

555 Also,

$$\begin{aligned} (\tau, M_x^{-1}, L_x^{-1}L_{x\tau}) \in ATP(Q, \cdot) \\ \Leftrightarrow s\tau \cdot yM_x^{-1} = (sy)L_x^{-1}L_{x\tau} \\ \Leftrightarrow yM_x^{-1}L_{s\tau} = yL_sL_x^{-1}L_{x\tau} \Leftrightarrow M_x^{-1}R_{s\tau} = R_sR_x^{-1}R_{x\tau} \end{aligned}$$

556 ■

557 **Corollary 3.13.** Let $(Q, \cdot, /, \setminus)$ be a commutative loop such that $M_x^{-1}R_{s\tau} =$
 558 $R_sR_x^{-1}R_{x\tau}$ for all $x, s \in Q$ and $\tau \in A(Q)$. $(H, *) = H(Q, \cdot)$ is commutative
 559 GMBL if and only if

- 560 1. (Q, \cdot) is a generalised middle Bol loop
- 561 2. $\tau = \tau(a, b) = R_{b \setminus a}R_b^{-1}$ for arbitrarily fixed $a, b \in Q$ and for each $\tau \in A(Q)$

562 **Proof.** It is straightforward. ■

563

CONCLUSION

564 In this research, we have been able to shown that the two identities of GMBL
 565 are equivalent if the generalising map α is bijective such that it fixes the identity
 566 element. Also, among all the five parastrophes of GMBL, (12)–parastrophe of
 567 GMBL is a GMBL and (13)– and (123)– parastrophes of Q are GMBL of expo-
 568 nent two. In line with Lemma 3.2, it can be seen that (13)– and (123)–parastrophes
 569 of GMBL of exponent two are loop. It is noted that (23)– and (132)–parastrophes
 570 of GMBL with commutative property are totally symmetric. The work further
 571 reveals that, in (13)–parastrophe of Q , the right inverse element coincides with
 572 left inverse element if α is bijective such that $\alpha : e \rightarrow e$ which is one of the general
 573 property of middle Bol loop revealed by Kuznetsov in [15].

574

REFERENCES

- 575 [1] J. O. Adeniran, T. G. Jaiyéolá and K. A. Idowu *Holomorph of generalized*
 576 *Bol loops*, Novi Sad Journal of Mathematics, 44, (1), 37–51, 2014.

- 577 [2] A. O. Abdulkareem, J. O. Adeniran, A. A. A. Agboola and G. A. Ade-
578 bayo *Universal α -elasticity of generalised Moufang loops*. Annals of Mathe-
579 matics and Computer Science. 14, 1–11, 2023.
- 580 [3] A. O. Abdulkareem, J. O. Adeniran *Generalised middle Bol loops*. Journal
581 of the Nigerian Mathematical Society 39 (3), 303-313, 2020.
- 582 [4] V. D. Belousov *Foundations of the theory of quasigroups and loops*, (Russian)
583 Izdat. “Nauka”, Moscow 223pp,1967.
- 584 [5] R. O. Fisher and F. Yates *The 6x6 latin squares*, Proc. Soc 30, 429-507, 1934.
- 585 [6] A. Gvaramiya *On a class of loops* (Russian), Uch. Zapiski MAPL. 375, 25-
586 34,1971.
- 587 [7] T. G. Jaiyéḡlá, S. P. David and Y. T. Oyebo New algebraic properties of
588 middle Bol loops. ROMAI J. 11 (2), 161–183, 2015.
- 589 [8] T. G. Jaiyéḡlá, S. P. David, E. Ilojide and Y. T. Oyebo Holomorphic
590 structure of middle Bol loops. Khayyam J. Math. 3(2), 172–184. 2017
591 <https://doi.org/10.22034/kjm.2017.51111>
- 592 [9] T. G. Jaiyéḡlá, S. P. David and O. O. Oyebola *New algebraic properties*
593 *of middle Bol loops II*. Proyecciones Journal of Mathematics 40(1), 85–106,
594 2021. <http://dx.doi.org/10.22199/issn.0717-6279-2021-01-0006>
- 595 [10] T. G. Jaiyéḡlá *Some necessary and sufficient conditions fro parastrophic in-*
596 *varance in the associative law in quasigroups*, Fasciculi Mathematici, 40 25–
597 35, 2008.
- 598 [11] T. G. Jaiyéḡlá *Basic Properties of Second Smarandache Bol Loops*, In-
599 ternational Journal of Mathematical Combinatorics, 2, 11–20, 2009.
600 <http://doi.org/10.5281/zenodo.32303>.
- 601 [12] T. G. Jaiyéḡlá *Smarandache Isotopy of Second Smaran-*
602 *dache Bol Loops*, Scientia Magna Journal, 7(1), 82–93, 2011.
603 <http://doi.org/10.5281/zenodo.234114>.
- 604 [13] T. G. Jaiyéḡlá and B. A. Popoola *Holomorph of generalized Bol loops II*,
605 Discussiones Mathematicae-General Algebra and Applications, 35(1), 59-
606 –78, 2015. doi:10.7151/dmgaa.1234.
- 607 [14] T. G. Jaiyéḡlá, B. Osoba and A. Oyem *Isostrphy Bryant-Schneider Group-*
608 *Invariant of Bol Loops*, Buletinul Academiei De S, Tiinte, A Republicii
609 Moldova. Matematica, 2(99), 3–18, 2022.

- 610 [15] E. Kuznetsov *Gyrogroups and left gyrogroups as transversals of a special*
611 *kind*, Algebraic and discrete Mathematics 3, 54–81, 2005.
- 612 [16] Grecu, I *On multiplication groups of isostrophic quasigroups*, Proceedings of
613 the Third Conference of Mathematical Society of Moldova, IMCS-50, 19-23,
614 Chisinau, Republic of Moldova, 78–81, 2014.
- 615 [17] I. Grecu and P. Syrbu *On Some Isostrophy Invariants of Bol Loops*, Bulletin
616 of the Transilvania University of Brasov, Series III: Mathematics, Informat-
617 ics, Physics, 54(5), 145–154,2012.
- 618 [18] A. Drapal and V. Shcherbacov *Identities and the group of isostrophisms*,
619 Comment. Math. Univ. Carolin, 53(3), 347–374, 2012.
- 620 [19] Syrbu, P. and Grecu, I *On some groups related to middle Bol loops*, Studia
621 Universitatis Moldaviae (Seria Stiinte Exacte si Economice), 7(67), 10–18,
622 2013.
- 623 [20] P. Syrbu *Loops with universal elasticity*, Quasigroups Related Systems,1,
624 57–65, 1994.
- 625 [21] P. Syrbu *On loops with universal elasticity*, Quasigroups Related Systems,3,
626 41–54, 1996.
- 627 [22] P. Syrbu *On middle Bol loops*, ROMAI J., 6(2), 229–236, 2010.
- 628 [23] P. Syrbu and I. Grecu *Loops with invariant flexibility under the isostrophy*,
629 Bul. Acad. Stiinte Repub. Mold. Mat. 92(1), 122-128, 2020.
- 630 [24] I. Grecu and P. Syrbu *Commutants of middle Bol loops*, Quasigroups and
631 Related Systems, 22, 81–88, 2014.
- 632 [25] B. Osoba and Y. T. Oyebo *On Multiplication Groups of Middle Bol Loop*
633 *Related to Left Bol Loop*, Int. J. Math. And Appl., 6(4), 149–155, 2018.
- 634 [26] Osoba. B and Oyebo. Y. T *On Relationship of Multiplication Groups and*
635 *Isostrophic quasigroups*, International Journal of Mathematics Trends and
636 Technology (IJMTT), 58 (2), 80–84, 2018. DOI:10.14445/22315373/IJMTT-
637 V58P511
- 638 [27] B. Osoba *Smarandache Nuclei of Second Smarandache Bol Loops*, Scientia
639 Magna Journal, 17(1), 11–21, 2022.
- 640 [28] B. Osoba and T. G. Jaiyéolá *Algebraic Connections between Right and Middle*
641 *Bol loops and their Cores*, Quasigroups and Related Systems, 30, 149-160,
642 2022.

- 643 [29] T. Y Oyebo, B. Osoba, and T. G. Jaiyéolá. *Crypto-automorphism Group of*
644 *some quasigroups*, *Discussiones Mathematicae-General Algebra and Appli-*
645 *cations*. Accepted for publication.
- 646 [30] Y. T. Oyebo and B. Osoba *More results on the algebraic properties of middle*
647 *Bol loops*, *Journal of the Nigerian mathematical society*, 41(2), 129-42, 2022.
- 648 [31] Pflugfelder, Hala O *Quasigroups and loops: introduction* . *Sigma Series in*
649 *Pure Mathematics*, 7. Heldermann Verlag, Berlin. viii+147, 1971.
- 650 [32] V. A. Shcherbacov *A-nuclei and A-centers of quasigroup*, *Institute of math-*
651 *ematics and computer Science Academy of Science of Moldova Academiei*
652 *str. 5, Chisinau, MD -2028, Moldova* ,2011
- 653 [33] A. R. T, Solarin, J. O. Adeniran, T. G. Jaiyéolá, A. O. Iseré and Y.
654 T. Oyebo. "Some Varieties of Loops (*Bol-Moufang and Non-Bol-Moufang*
655 *Types*)". In: Hounkonnou, M.N., Mitrović, M., Abbas, M., Khan, M. (eds)
656 *Algebra without Borders – Classical and Constructive Nonassociative Alge-*
657 *braic Structures*. STEAM-H: Science, Technology, Engineering, Agriculture,
658 *Mathematics & Health*. Springer, Cham. 2023. [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-3-031-39334-1_3)
659 [3-031-39334-1_3](https://doi.org/10.1007/978-3-031-39334-1_3)

660
661
662

Received
Revised
Accepted